

24 Février 1975



LRB 75/02

HIGH RESOLUTION LASER SPECTROSCOPY
COPY OF THE D LINES OF ON-LINE PRODUCED
21,22,24,25_{Na} USING A NEW HIGH SENSITIVITY
METHOD OF DETECTION OF OPTICAL RESONANCES

G. Huber, C. Thibault, R. Klapisch
H.T. Duong, J.L. Vialle, J. Pinard,
P. Juncar, P. Jacquinet

HIGH RESOLUTION LASER SPECTROSCOPY OF THE D LINES OF OH-LINE PRODUCED
21, 22, 24, 25 Na USING A NEW HIGH SENSITIVITY METHOD OF DETECTION OF
OPTICAL RESONANCES

G. Huber[†], C. Thibault, E. Klapisch

Laboratoire Ferné Bernas du Centre de Spectrométrie Nucléaire et de
Spectrométrie de Masse, 91406 ORSAY, France.

and

H.T. Duong, J.L. Vialle, J. Pinard, P. Juncar^X, P. Jacquinet

Laboratoire Aimé Cotton, C.N.R.S. II, Bâtiment 505, 91405 ORSAY, France.

ABSTRACT

A polyisotopic sodium beam of ²¹⁻²⁵Na, produced by spallation of Al, was illuminated with a tunable cw dye laser. The atomic beam, analysed by a simple magnet is then ionized and detected after a mass spectrometer.

The results are : isotopic shifts relative to ²³Na : ²¹Na = -53.5 mK, ²²Na = -25.3 mK, ²⁴Na = +23.1 mK, ²⁵Na = +44.6 mK ; $\nu_I(^{25}\text{Na}) = +3.685$ (22) m.m. ; b-factor in the $P_{3/2}$ state of ²⁵Na = -16 (8) MHz.

ANALYTIC SUBJECT INDEX : 11.4 ; 54.1

Half-lives ¹, masses ², and other properties ³, of short-lived sodium isotopes have recently been determined using a mass spectrometer on-line with an accelerator. As an extension of this work, we wished to determine other static properties of these nuclei. We have thus attempted to measure the hyperfine structure (hfs) and isotope shift (IS) of the atomic D lines of sodium isotopes. The purpose of this paper is to present the first results that have been obtained using a newly developed method that could eventually extend our knowledge on the shape and magnetic properties of nuclei far from stability.

The hfs of the ground state of ²⁰⁻²⁴Na has already been measured by different methods ⁴; recently a determination of the absolute value of the magnetic moment of ²⁵Na has been published ⁵. However the hfs of the excited 3P states are unknown for the unstable isotopes and there are no data available for isotope shifts.

This type of measurement on a series of isotopes can give interesting indications on nuclear shapes. The observation of irregularities of the hfs and IS of short lived Hg isotopes ⁶ has been especially fruitful in that context. In the case of sodium, however, the same method as in ref. 6 would fail because of the Doppler broadening, and it is necessary to use one of the new methods of the Doppler free high resolution spectroscopy ⁷⁻⁹.

The special requirements of our on-line experiment have in fact led to the development of a variant of one ⁷ of these methods of detection of optical transitions in an atomic beam ¹⁰. In essence this method rests upon the magnetic detection of the optical pumping which occurs when a laser beam is tuned to the frequency of one of the hyperfine components of the D lines.

This optical pumping changes the magnetic state ($m_j = \pm 1/2$) of the atoms of an atomic beam and this change is detected by means of a magnetic filter, a six pole magnet which focusses the atoms with $m_j = + 1/2$ and defocusses the atoms with $m_j = - 1/2$.

In the actual experiment, the isotopes of sodium are produced in the spallation of aluminum by 150 MeV protons from the Orsay synchrocyclotron. Since this nuclear reaction results in a mixture of the different isotopes and their optical resonances might overlap, it is necessary to separate them on-line by a mass spectrometer.

A highly efficient target has been developed in order to have the best sodium isotopes production. We have used a molten metal target in a graphite container. It was loaded with 10 g of high purity aluminum and heated by alternating current up to 900° C. The yield of the $^{27}\text{Al}(p, 3pxn)\text{Na}$ spallation reaction with 150 MeV protons was studied by directly connecting the target to the thermoionic source of the mass spectrometer. The counting rate after mass separation is shown in Fig. 1. The diffusion delay time in the molten metal is strongly temperature dependent and decreases from 1 mn to 10 s between 850° C and 900° C.

This target is used in the present experiment as the oven of the atomic beam. A schematic view is given in Fig. 2. The collimation of the beam is defined by a nitrogen-cooled diaphragm to 30/1. The sodium atoms effusing from the oven interact with the laser beam before passing the six-pole magnet. The outgoing atoms impinge on a hot rhenium surface acting as the ion source of a small magnetic mass spectrometer. Mass separated ions are counted after the exit slit with an electron multiplier. This device detects equally well atoms of stable or short lived isotopes.

The laser is a single mode tunable dye laser of high stability which has been used in other experiments ¹¹. A new frequency control and scanning system can provide a frequency change by increments of 7.5 MHz (0.25 mK) initiated by pulses from a pilot clock; this system, called "sigma-meter" will be described elsewhere ¹². A small part of the laser light is used to observe the fluorescence signal from an auxiliary ²³Na atomic beam which serves as a reference.

For each isotope, the hfs of its D₁ line is recorded in a single scan, together with the corresponding line of ²³Na. The scanning time is about 10 ms. The signals are directly monitored on a 2-channel chart recorder and fed simultaneously, by a specially built routing device, into the different subgroups of a multichannel analyzer.

The total beam leaving the oven contains up to 10⁸ atoms/s of ²⁵Na. Because of the different losses, the corresponding counting rate at the detector is then 3000 per second. The background of the electron multiplier due to nuclear reaction is only 30-50 counts/s. It should be noted however that tailing of ²³Na ions at the mass 22 results in up to 1000 counts/s. This could certainly be improved in the future by adding an electrostatic deviation after the magnetic mass spectrometer.

Fig. 3 shows the spectra obtained. Our results on the hfs of the ²S_{1/2} ground state are in agreement with the known values obtained by radio-frequency techniques for ²¹, ²², ²⁴Na within the limits of the statistical uncertainties of the signal and accidental irregularities of the scanning by the sigma-meter. Our value for the magnetic moment of ²⁵Na is $\mu_I = +3.685 (22) \mu_n$. The value $|\mu_I| = 3.710 (70)$ given by M. Deimling et al. ⁵ is in agreement with our number.

The measured values of the isotope shifts are given in table I.

It appears that the measured values follow the law of the normal plus specific mass effect within the experimental errors. This type of experiment can also lead to a determination of the nuclear electric quadrupole moment if the hyperfine structure of the $^2P_{3/2}$ state can be measured with a good precision. For that purpose, we have also observed the four (2,3 - 2,3) transitions in the D_2 line of ^{25}K , which were only partially resolved with the present collimation of the atomic beam. From this measurement we have estimated the b factor for the $^2P_{3/2}$ state to be $b = -16$ (8) MHz. An experiment with a better collimation for the measurement of b is in progress.

The authors would like to thank their colleagues of the Institut de Physique Nucléaire whose support and help made it possible to carry out this experiment at the Orsay synchrocyclotron. It is a pleasure to acknowledge the able technical assistance of R. Baronnet (atomic beam apparatus), R. Leroux (electronics of the Sigma-meter), B. Roserbaum (data acquisition system), J.F. Kepinski (assembly and testing of the apparatus) and M. Jacotin (electronics of the mass spectrometer).

FOOTNOTES AND REFERENCES

[†] on leave from University Mainz and G.S.I. Darmstadt, W. Germany.

^x present address : Institut für Angewandte Physik der Universität Heidelberg.

- 1 - R. Klapisch, C. Thibault, A.M. Poskanzer, R. Prieels, C. Rigaud and E. Roeckl, Phys. Rev. Lett., 29, 1254 (1972).
- 2 - R. Klapisch, R. Prieels, C. Thibault, A.M. Poskanzer, C. Rigaud and E. Roeckl, Phys. Rev. Lett., 31, 118 (1973).

- 3 - E. Roeckl, P.F. Dittner, C. Métraux, K. Klupinich, C. Thibault and C. Rigaud, *Phys. Rev.*, C 10, 1191 (1974).
- 4 - G.H. Fuller and V.W. Cohen, *Nuclear Data Tables*, 5 (1969)
and
J. Bonn, E.W. Otten, Ch. Von Platen, W. Linder, H. Müller and H. Schweickert, Proceedings of the International Conference on the Properties of Nuclei far from the Region of Beta-Stability, Leysin, 1970, CERN Report n° CERN 70-30, 363 (1970).
- 5 - M. Deinling, J. Dietrich, R. Neugart, H. Schweickert, Fourth International Conference on Atomic Physics, Heidelberg, 207 (1974).
- 6 - J. Bonn, G. Huber, H.J. Kluge, U. Köpf, L. Kugler and E.W. Otten, *Phys. Lett.* 36 B, 41 (1971) and *Phys. Lett.*, 38 B, 308 (1972).
- 7 - H.T. Duong, P. Jacquinet, S. Liberman, J.-L. Picqué, J. Pinard, and J.-L. Vialle, *Opt. Comm.*, 7, 371 (1973).
- 8 - P. Jacquinet, S. Liberman, J.-L. Picqué and J. Pinard, *Opt. Comm.*, 8, 163 (1973).
- 9 - H.T. Duong, P. Jacquinet, S. Liberman, J. Pinard, J.-L. Vialle, *C.R. Acad. Sci. Paris*, 275 B, 909 (1973).
- 10 - H.T. Duong and J.-L. Vialle, *Opt. Comm.*, 12, 71 (1974).
- 11 - S. Liberman and J. Pinard, *Appl. Phys. Lett.*, 24, 142 (1974).
- 12 - J. Pinard and P. Juncar, to be published.

TABLE 1

Relative isotopic shifts of $^{21-25}\text{Na}$

Isotopes	Isotopic shift: (mK)
$^{21}\text{Na} - ^{23}\text{Na}$	- 53,5 (2)
$^{22}\text{Na} - ^{23}\text{Na}$	- 25,3 (2)
$^{24}\text{Na} - ^{23}\text{Na}$	+ 23,1 (4)
$^{25}\text{Na} - ^{23}\text{Na}$	+ 44,6 (2)

FIGURE CAPTIONS

Figure 1 - Isotopic yield of the $^{27}\text{Al} (p, 3pxn)^{25-x}\text{Na}$ spallation reaction with 150 MeV protons.

Figure 2 - Schematic view of the experiment.

Figure 3 - hfs and IS of the D_1 lines of $^{21-25}\text{Na}$ isotopes. These lines, measured individually relative to ^{23}Na are here shown together on a common wavenumber scale.




Fig 1




Fig. 2




Fig. 3

