

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

ДУБНА



E4 - 8543

M.M.Musakhanov, A.L.Zubarev

EVALUATION OF FRESNEL'S CORRECTIONS
TO THE EIKONAL APPROXIMATION
BY THE METHOD OF SEPARABILIZATION

1975

RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statut. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and English.

The preprints and communications, which are distributed only to the Member States of the JINR, have no letter-index.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

1. I. Ivanov. JINR, P2-4985, Dubna, 1971.

Мусаханов М.М., Зубарев А.Л.

E4 - 8543

Исследование френелевских поправок к эйкональному приближению методом сепарабилизации

В работе предложен метод сепарабилизации потенциала по приближенным решениям уравнения Шредингера, приводящий к вариационному принципу Швингера для амплитуды рассеяния. Эти результаты применяются для вычисления френелевских поправок к приближению Глаубера.

Препринт Объединенного института ядерных исследований
Дубна 1975

Musakhanov M.M., Zubarev A.L.

E4 - 8543

Evaluation of Fresnel's Corrections to the Eikonal Approximation by the Method of Separabilization

A method is suggested for separabilization of the potential by means of the solutions of the Schrödinger equations resulting in the Schwinger variational principle for the scattering amplitude. The results are applied to the calculation of the Fresnel's corrections to the Glauber approximation.

Preprint of the Joint Institute for Nuclear Research
Dubna 1975

E4 - 8543

M.M.Musakhanov,* A.L.Zubarev*

**EVALUATION OF FRESNEL'S CORRECTIONS
TO THE EIKONAL APPROXIMATION
BY THE METHOD OF SEPARABILIZATION**

Submitted to *ЖФ*

* Permanent address: Tashkent
State University

The separabilization method is applied widely to solve the quantum-mechanical problem of 3 and more bodies as it leads ultimately to the solution of one-dimensional equations⁽¹⁻⁴⁾.

The content of the method is as follows. Let V be an interaction potential that always may be presented as

$$V = \sum_{i,j} V|c\rangle\langle i|V^{-1}|j\rangle\langle j|V.$$

Retaining a finite number of forms we obtain a separable potential

$$V^{\sim} = \sum_{i,j} V|c\rangle\langle i|V^{-1}|j\rangle\langle j|V. \quad (1)$$

Convergence of the approximation (1) is provided by the well-known theorem of the functional analysis that a quite continuous operator

V may be approximated according to the norm by the operator V^{\sim} with any accuracy. In this study we modify the presentation (1), viz, change V by the separable

$$V^s = \frac{V|\varphi_k^+\rangle\langle\varphi_{k'}^-|V}{\langle\varphi_{k'}^-|V|\varphi_k^+\rangle}. \quad (2)$$

Schrödinger's equation with the potential (2) is solved in explicit form and for the scattering amplitude we obtain

$$f^s = \langle k'|V|\varphi_k^+\rangle C(k, k'). \quad (3)$$

where ψ_{κ}^{\pm} are any approximated solutions of Schrödinger's equation

$$\begin{aligned} |\psi_{\kappa}^{+}\rangle &= |k\rangle + G V |\psi_{\kappa}^{+}\rangle \\ \langle \psi_{\kappa}^{-}| &= \langle k| + \langle \psi_{\kappa}^{-}| V G. \end{aligned} \quad (4)$$

Let us formulate now the iteration method of solution. In equation (4) we change V by V^S , and as ψ^{\pm} we choose any approximated solutions f^{\pm} . Then with the potential (2) the equation (4) is solved in explicit form and $\psi_0^{\pm}, \psi_0^{\pm}(f)$ are found. Thereafter supposing, in (2) $\varphi^{\pm} = \psi_0^{\pm}$ the equation is solved again, etc.

Without dwelling on the convergence of this process, we note that it leads at every step for the scattering amplitude to Schwinger's variation principle⁽⁵⁾ (2). Really, if

$$\varphi^{\pm} - \psi^{\pm} = \varepsilon^{\pm} \sim \varepsilon, \quad (5)$$

then $f(k', k) - f^S(k', k) \sim \varepsilon^2$.

We shall use further this property widely.

With decreasing particle energy or increasing scattering angle the role of corrections (called Fresnel ones) connected with deviations from the eikonal approximation increases.

It should be remembered that this approximation corresponds to the solution of the Schrödinger equation with approximated Green's function

$$\begin{aligned} G_0(\vec{r}) &= \frac{m}{i\kappa} e^{i\kappa z} \Theta(z) \delta(\vec{r}_{\perp}), \\ \text{here } \vec{r} &= (0, 0, k) \quad \vec{r}_{\perp} \perp \vec{K}. \end{aligned} \quad (6)$$

The solutions of the equation with such Green's function are known

$$\begin{aligned} |\psi_{\kappa}^{+}\rangle &= \exp\left\{i\kappa z - \frac{i\kappa}{\kappa} \int_{-\infty}^z V(\vec{r}_{\perp}, z') dz'\right\} \\ \langle \psi_{\kappa}^{-}| &= \exp\left\{-i\kappa z - \frac{i\kappa}{\kappa} \int_z^{\infty} V(\vec{r}_{\perp}, z') dz'\right\}. \end{aligned} \quad (7)$$

Now functions (7) are inserted into (2) and (3).

As a result the amplitude takes the form

$$f_1(\vec{k}, \vec{k}') = C(\vec{k}, \vec{k}') f_0(\vec{k}, \vec{k}'); \quad f_0 = \langle \vec{k}' | V | \varphi_{\vec{k}}^+ \rangle$$

$$C = f_0(\vec{k}, \vec{k}') [f_0(\vec{k}, \vec{k}') - \int f_0(\vec{k}, \vec{k} - \vec{\eta}) f_0(\vec{k} - \vec{\eta}, \vec{k}') g(\vec{k}, \vec{\eta}) \frac{d^3 \vec{\eta}}{(2\pi)^3}]. \quad (8)$$

Here $g = G - G_0$ is the difference between the exact and eikonal Green's functions.

It is clear that amplitude $f_1(\vec{k}, \vec{k}')$ in formula (8) differs exactly with consideration of Fresnel's corrections.

From the explicit expression for

$$g(\vec{k}, \vec{\eta}) = m \frac{\eta^2}{(\vec{k}\vec{\eta} + i\epsilon)(2\vec{k}\vec{\eta} - \eta^2 + i\epsilon)} \quad (9)$$

it follows that a) $C \rightarrow 1$ for $E \rightarrow \infty$

b) with increase of the scattering angle C increases

c) for the explicit calculation of the correction an off-mass shell amplitude in eikonal approximation is necessary.

For example let us view the particle scattering Gaussian potential

$$V(r) = A \exp(-\alpha^2 r^2).$$

Restricting ourselves to the lowest k/E term in the correction

factor C we obtain the result

$$C^{-1} = 1 - I,$$

where

$$I = m \int \frac{d^3 \eta}{(2\pi)^3} \frac{\eta^2}{(\vec{k}\vec{\eta} + i\epsilon)(2\vec{k}\vec{\eta} - \eta^2 + i\epsilon)} \frac{V(\vec{\Delta} + \vec{\eta})}{V_{\Delta}} V_{\eta}. \quad (10)$$

Here

$$\vec{\Delta} = \vec{k}' - \vec{k}; \quad \vec{\Delta}\vec{k} \approx 0; \quad V_{\Delta} = \langle \vec{k}' | V | \vec{k} \rangle = \frac{\pi^{3/2} A}{|\alpha|^3} e^{-\Delta^2/(2\alpha)^2}.$$

Calculation of integral leads to the functions of the parabolic cylinder. However in limit cases

a) $\Delta/a \ll 1$ and b) $\Delta/a \gg 1$

the result takes a simple form

$$a) \bar{I} = -\frac{m}{2\pi} \frac{A}{2^3 \kappa^2} \left(1 + \frac{1}{2} \frac{\Delta^2}{2a^2} \right), \quad (11)$$

$$b) \bar{I} = -\frac{m}{2\pi} \frac{A}{2^3 \kappa^2} e^{\Delta^2/8a^2} \frac{\Delta}{2\sqrt{2}a}. \quad (12)$$

It means that the correction may become essential for large scattering angles.

Calculating Glauber's expression in the framework of the potential theory for the scattering amplitude of a rapid particle on the nucleus the following approximations are made⁽⁸⁾.

1. The binding and kinetic energy of nucleus nucleons are neglected. This approximation is often referred to as adiabatic one ($G \rightarrow \tilde{G}$).

2. The eikonal approximation is used ($\tilde{G} \rightarrow \tilde{G}_0$). Here G is exact Green's function.

In the adiabatic approximation Schrödinger's equation may be written in integral form (4). Therefore the results may be applied.

Let us view an example of the elastic πd scattering.

In this case

$$\tilde{G}(\vec{r}, R) = \delta(\vec{r}) \int \frac{d^3 p}{(2\pi)^3} e^{i\vec{p}\vec{R}} \left[\frac{R^2 - \vec{p}^2}{2m} - \frac{(R - \vec{p})^2}{2M} + i\epsilon \right]^{-1}. \quad (13)$$

Here $\vec{k}, m(\vec{k}, \vec{m})$ are the momentum and mass of the incident (scattered) meson. M is deuteron mass. The difference between \tilde{G} and \tilde{G}_0 is

$$g(\vec{R}, \vec{r}) = \delta(\vec{r}) \frac{m^2}{2\mu} \int \frac{d^3 q}{(2\pi)^3} e^{i(\vec{R} - \vec{q})\vec{R}} g_0(\vec{R}, \vec{q})$$

$$g(\vec{R}, \vec{q}) = \frac{m^2}{2\mu} g_0(\vec{R}, \vec{q}), \quad g_0(\vec{R}, \vec{q}) = \frac{m^2}{2\mu} \frac{\vec{q}^2}{(\vec{R}\vec{q} + i\epsilon) \vec{R}\vec{q} - \vec{q}^2 + i\epsilon}, \quad (14)$$

where $\mu = \frac{mM}{m+M}$

The expression for the scattering amplitude on fixed centres that includes all Fresnel's correction is as follows

$$f_1(\vec{k}, \vec{k}', \vec{z}) = C(\vec{k}, \vec{k}', \vec{z}) f_0(\vec{k}, \vec{k}', \vec{z})$$

$$C = f_0(\vec{k}, \vec{k}', \vec{z}) \left[f_0(\vec{k}, \vec{k}', \vec{z}) - \int \frac{d^3 q}{(2\pi)^3} f_0(\vec{k}, \vec{k} - \vec{q}, \vec{z}) f_0(\vec{k} - \vec{q}, \vec{k}', \vec{z}) g(\vec{k}, \vec{q}) \right]^{-1} \quad (15)$$

Here f_0 is an amplitude in eikonal (Glauber's) approximation. Averaging then over initial and final wave functions we obtain an expression for π^d scattering amplitude. To compare with the results of other studies (7-10) (3) let us calculate expression (15) approximately. Supposing the correction to be small we obtain

$$f_1 = f_0 + \int \frac{d^3 q}{(2\pi)^3} f_0(\vec{k}, \vec{k} - \vec{q}, \vec{z}) f_0(\vec{k} - \vec{q}, \vec{k}', \vec{z}) g(\vec{k}, \vec{q}) \quad (16)$$

In the second term in (16) let us take the amplitude in the following approximation

$$f_0 \approx e^{i\vec{d}\vec{z}/2} f_{0\pi N}(\Delta) + e^{-i\vec{d}\vec{z}/2} f_{0\pi p}(\Delta) \quad (17)$$

Note that πN amplitudes are to be taken in the eikonal approximation ($f_{0\pi N}$)

As a result we have

$$f_1 = f_0 + e^{i\vec{d}\vec{z}/2} \int g(\vec{q}) f_{0\pi N}(\vec{q}) f_{0\pi N}(-\vec{q} + \vec{d}) \frac{d^3 q}{(2\pi)^3} +$$

$$+ e^{-i\vec{d}\vec{z}/2} \int g(\vec{q}) f_{0\pi p}(\vec{q}) f_{0\pi p}(-\vec{q} + \vec{d}) \frac{d^3 q}{(2\pi)^3} + \quad (18)$$

$$+ \int g(\vec{q}) f_{0\pi p}(\vec{q}) f_{0\pi N}(-\vec{q} + \vec{d}) e^{i(\vec{q} + \vec{d}/2)\vec{z}} \frac{d^3 q}{(2\pi)^3} +$$

$$+ \int g(\vec{q}) f_{0\pi N}(\vec{q}) f_{0\pi p}(-\vec{q} + \vec{d}) e^{i(\vec{q} - \vec{d}/2)\vec{z}} \frac{d^3 q}{(2\pi)^3}.$$

It is seen that the 2nd and 3rd terms simply renormalize the approximation (17) and correspond with Fresnel's corrections, to the amplitudes of the πN scattering. The 4th and 5th terms are Fres-

nel's corrections renormalizing the rescattering effects. These corrections have been studied in a number of works⁽⁷⁻¹⁰⁾.

By any calculation method an evident difficulty is the question of the one-mass shell amplitude parametrization.

The authors express their sincere gratitude to V.B. Belyaev and B.I. Zamarjev for support and interest in this study and to B.S. Kopeliovich, L.I. Lapidus, G.I. Likasov, A.V. Tarasov for useful discussions on the present study.

References:

1. L.D. Fadueyev. Doklady na V mezhduнарод. konf. po fizike elektronnykh i atomnykh stolknoveniy. "Nauka", 1967 (in Russian).
2. A.G. Sitenko, V.F. Kharchenko, K.L. Petrov. Phys.Lett. 28 B, 308, 1968.
3. V.B. Belyaev, A.L. Zubarev. Jadernaya fizika. 14,545, 1971 (in Russian).
4. A. N. Mitra. Nucl. Phys. 32, 529, 1962.
5. M.L. Goldberger, K.M. Watson. "Collision Theory" (Wiley, New York), 1964.
6. C.J. Joachain, C. Quigg. Rev. Mod. Phys. 46,279, 1974.
7. K. Gottfried. Ann.Phys. 66,368, 1971.
8. A.L. Kolybasov, L.A. Kondratyuk. Jadernaya fizika 18,316, 1973 (in Russian).
9. L.I. Koval, S.G. Matinyan. Jadernaya fizika 19,380, 1974 (in Russian).
10. A.D. Polozov. Ukrainskiy fizicheskiy zhurnal, 18,11, 1973 (in Russian).

Received by Publishing Department
on January 20, 1975

Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups and individual scientists of more than 50 countries.

We expect that the recipients of the JINR publications will have a possibility to display initiative in sending free of charge publications to Dubna. We receive, on the exchange basis, scientific books, journals, preprints and other types of publications on the subject categories of our Institute.

The only kind of publications that we do not require are the reprints of the articles already published in scientific journals.

In a number of cases we address to scientific institutions - the most prominent recipients of our publications - with the request to send us free of charge some books or subscribe for our library to the scientific journals published in their countries.

Requests

The Publishing Department fulfils annually about 3000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

Publishing Department
Joint Institute for
Nuclear Research
Head Post Office,
P.O. Box 79
101000 Moscow,
U.S.S.R.

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

Scientific-Technical Library
Joint Institute for
Nuclear Research
Head Post Office
P.O. Box 79
101000 Moscow,
U.S.S.R.



Издательский отдел Объединенного института ядерных исследований.
Заказ 19108. Тираж 620. Уч.-изд. листов 0,36.
Редактор Б.Б. Колесова. Подписано к печати 5.2.75 г.