

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА



E2 - 9803

2/11/76

J.Fischer, P.Kolář

**DERIVATIVE ANALYTICITY RELATIONS
AND ASYMPTOTIC ENERGIES**

1976

RANGE OF JINR PUBLICATIONS

The preprints and communications of the Joint Institute for Nuclear Research (JINR) are considered to be original publications. They are published in accordance with Article 4 of the JINR Statute. Difference between the preprints and communications consists in that text of the preprint will be published in future in one of scientific journals or in aperiodic collections.

Indexing

The preprints, communications and deposited publications of the JINR have a single numbering (four last figures of the index).

The first sign of the index - a letter, denotes the language the paper is published in:

"P" - published in Russian;

"E" - published in English;

"D" - published both in Russian and in English.

The figure following the letter denotes the subject category of the given publication. List of subject categories of the JINR publications is sent periodically to the recipients.

Above-described index is placed in the right upper corner of the cover and on the title-page of each publication.

References

In bibliographical references to the JINR preprints and communications we recommend to indicate: author's initials and name, an abbreviation of the name of the institute, index, place and year of publication.

An example of bibliographical reference:

I. Ivanov. JINR, P2-4985, Dubna, 1971.

E2 - 9803

J.Fischer,* P.Kolář

**DERIVATIVE ANALYTICITY RELATIONS
AND ASYMPTOTIC ENERGIES**

Submitted to "Physics Letters" and to the
XVIII International Conference on High
Energy Physics, Tbilisi, 1976.

* Institute of Physics, ČSAV, Prague,
Czechoslovakia.

In the present note we discuss some problems related to derivative analyticity relations which, under certain conditions, give point-to-point correlations between the real and the imaginary part of a scattering amplitude on the energy cut. In the case of the symmetric amplitude $F_S(s,t)$, for instance, Bronzan et al.^{1/} have derived the relation

$$\operatorname{Re} \tilde{F}_S(s,t) = s^\alpha \tan \left[\frac{\pi}{2} \left(\alpha - 1 + \frac{2}{2 \ln s} \right) \right] \frac{\operatorname{Im} \tilde{F}_S(s,t)}{s^\alpha} \quad (1)$$

(α being a real number), assuming that the imaginary part can be expanded in powers of $\ln s$ on the energy cut. Here $\tilde{F}_S(s,t)$ means the amplitude without poles and subtraction constants, s and t denoting the usual Mandelstam variables.

Suppressing higher branch points given by the unitarity condition, this derivation restricts severely the validity of the result to certain mathematical models. Indeed, it has been proved by Eichmann and Dronkers^{2/} that relation (1) is exactly valid only on some class of entire functions of $\ln s$. Further, Heidrich and Kazes^{3/} show that relation (1) is violated even in the energy intervals between two branch points.

In the present note we examine the derivative analyticity relations in the asymptotic regime.

Retaining only one derivative in (1), we obtain the following relation

$$\frac{\operatorname{Re} \tilde{F}_S(s,t)}{s} = \frac{\pi}{2} \frac{\partial}{\partial \ln s} \frac{\operatorname{Im} \tilde{F}_S(s,t)}{s} \quad (2)$$

which has been derived by Gribov and Migdal^{4/} in the Regge theory for the contribution of the Pomeron and its cuts. We shall show that relations of type (2) are, in the region of asymptotic energies, an immediate consequence of the general, model-independent principles of the S-matrix theory provided that the high-energy limits (finite or infinite) of certain quantities are assumed to exist.

The symmetric and the antisymmetric hadron-hadron elastic scattering amplitude will be denoted by $F_S(W,t)$ and $F_A(W,t)$, respectively, W being defined as $(s-u)/4M$, where M is

the mass of the target particle. Further, we use the following notation for convenience:

$F_{S,A}(E) \equiv F_{S,A}(E,0)$, $\sigma_S(E) = \frac{1}{E} \text{Im } F_S(E)$ (the optical theorem),

$$\rho_{S,A}(E,t) \equiv \frac{\text{Re } F_{S,A}(E,t)}{\text{Im } F_{S,A}(E,t)} \quad \text{and} \quad \rho_{S,A}(E) \equiv \rho_{S,A}(E,0).$$

We recall the following general properties^{/5/} of $F_{S,A}(E,t)$:

(i) There exist two positive numbers, $R(t)$ and t_0 , such that $F_{S,A}(E,t)$ are analytic functions of E in the upper half of the complex E -plane for $|E| > R(t)$, provided that $|t| < t_0$. This region will be denoted by \mathcal{D}_t .

(ii)

$$F_S(E,t) = F_S^*(-E^*,t)$$

$$F_A(E,t) = -F_A^*(-E^*,t)$$

for every $E \in \mathcal{D}_t$ and $|t| < t_0$.

(iii) $F_{S,A}(E,t)$ on the energy cut have been obtained by regularizing the corresponding distributions and are continuous in the closure of \mathcal{D}_t .

(iv) $F_{S,A}(E,t)$ are polynomially bounded for $|E| \rightarrow \infty$, $E \in \mathcal{D}_t$, $|t| < t_0$.

(v) For the real values of E , $F_{S,A}(E,t)$ satisfy the Froisart-Martin bound

$$|F_{S,A}(E,t)/E \ln^2 E| < \text{const}$$

for any $E > E_1$ (the bound for $t < 0$ being even stronger).
(vi)

$$\text{Im } F_S(E,t=0) \geq 0 \quad \text{for any } E > E_2.$$

These properties allow us to derive the following statements.

Statement 1. Let $F_S(E)$ be a function of complex E satisfying conditions (i) to (vi). Let further $\lim_{E \rightarrow \infty} \rho_S(E)$ exist and

$$\lim_{E \rightarrow \infty} \sigma_S(E) = \infty. \quad (3)$$

If the limit

$$\lim_{E \rightarrow \infty} \frac{I}{2} \frac{E \frac{d}{d \ln E} \sigma_f(E)}{\operatorname{Re} F_f(E)} \quad (4)$$

exists, then its value is 1.

In the remainder of the present paper, t is kept fixed in the interval $-t_0 < t \leq 0$.

The following statement includes both forward and non-forward scattering and allows $\operatorname{Im} F_S(E, t)/E$ to tend to infinity or to zero.

Statement 2. Let $F_S(E, t)$ satisfy the conditions (i) to (vi). Let $\operatorname{Im} F_S(E, t)$ not change the sign above some energy on the cut and the conditions

$$\left| \int_{E_0}^{\infty} \operatorname{Im} F_S(E, t) \frac{dE}{E} \right| = \infty, \text{ for some } E_0 \quad (5)$$

and

$$\left| \lim_{E \rightarrow \infty} \ln \left| \frac{F_S(E, t)}{E} \right| \right| = \infty \quad (6)$$

be satisfied. If the limit

$$\lim_{E \rightarrow \infty} \frac{I}{2} \frac{E \frac{\partial}{\partial \ln E} |F_S(E, t)/E|}{|F_S(E, t)| \arctan \beta_S(E, t)} \quad (7)$$

exists, then its value is 1.

The proof is given in the Appendix.

As a consequence, we find for the antisymmetric amplitude $F_A(E, t)$ the following result:

Corollary. Let $F_A(E, t)$ satisfy conditions (i) to (v) and let further

$$a) \lim_{E \rightarrow \infty} \frac{|F_A(E, t)|}{E} = 0; \quad (8)$$

b) $\text{Im } F_A(E, t)$ not change the sign above some energy;

c)
$$\left| \int_{E_0}^{\infty} \text{Im } F_A(E, t) dE \right| = \infty \quad \text{for some } E_0 \quad (9)$$

d)
$$\lim_{E \rightarrow \infty} \left| \text{Im } |F_A(E, t)| \right| = \infty \quad (10)$$

If the limit

$$\lim_{E \rightarrow \infty} \frac{\pi}{2} \frac{\frac{\partial}{\partial \ln E} |F_A(E, t)|}{|F_A(E, t)| \arctan \rho_A(E, t)} \quad (11)$$

exists, then its value is 1.

An interesting high-energy result can be obtained for the difference

$$D(E, t) = \frac{\pi}{2} \frac{\partial}{\partial \ln E} \left| \frac{F_S(E, t)}{E} \right| - \left| \frac{F_S(E, t)}{E} \right| \arctan \rho_S(E, t) \quad (12)$$

which, provided that $\rho_S(E, t)$ is sufficiently smooth and tends to zero, transforms into

$$\frac{\pi}{2} \frac{\partial}{\partial \ln E} \left(\frac{\text{Im } F_S(E, t)}{E} \right) - \frac{\text{Re } F_S(E, t)}{E} \quad (13)$$

Statement 3. Let $F_S(E, t)$ be a function satisfying (i) to (vi). Let $\text{Im } F_S(E, t)$ not change the sign above some energy on the cut and condition (5) be satisfied. If the limit $\lim_{E \rightarrow \infty} D(E, t)$ exists then its value is 0.

The proof is sketched in the Appendix.

Similar theorems for the difference (13) and for $F_A(E, t)$ can also be obtained; details will be published elsewhere.

We should like to draw the reader's attention to some interesting differences among the statements derived. The symmetric total cross-section $\sigma_S(E)$ is required to tend to infinity in Statement 1. If $\sigma_S(E)$ approaches zero in the high-energy limit, expression (4) is modified to (7) because $\rho_S(E)$ may be non-vanishing at $E \rightarrow \infty$ in this case^{16/}. In contra-

distinction to this no explicit restriction is imposed on the high-energy behaviour of $\sigma_S(E)$ in Statement 3.

Let us note that the results obtained have the form of high-energy limits and no a priori claim can be laid to their validity at finite energies. On the other hand, referring to the measurements of small-angle hadron-hadron scattering at the CERN ISR and at Fermilab above 100 GeV (lab), one can infer that the available energies would be already appropriate for the numerical verification of the Statements presented.

We are indebted to O. Dumbravs for stimulating discussions.

APPENDIX

Firstly, we give the proof of Statement 2. Statement 1 is proved in an analogous manner.

Since the limit (7) exists, the limit

$$\lim_{E \rightarrow \infty} \left[\left(\frac{2}{\pi} \int_{E_0}^E \arctan \rho_S(E', t) \frac{dE'}{E'} - \ln \left| \frac{F_S(E, t)}{E} \right| \right) / \ln \left| \frac{F_S(E, t)}{E} \right| \right] \quad (14)$$

also exists because of l'Hôpital's rule. The Cauchy theorem for $\ln(-F_S(E, t))/E$ implies ^{6,7)}

$$\ln \left| \frac{F_S(E, t)}{E} \right| = \frac{2}{\pi} \int_{E_0}^E \arctan \rho_S(E', t) \frac{dE'}{E'} + C - \mu(E, t) \quad (15)$$

where
$$\mu(E, t) = \frac{1}{\pi} \int_0^{\pi} \ln \left| \frac{F_S(Ee^{i\varphi}, t)}{F_S(E, t)} \right| d\varphi$$

and C is a constant depending on E_0 and t (t fixed $-t_0 < t \leq 0$). Combining (14) with (15) and using (6) we easily see that the limit (14) is equal to

$$\lim_{E \rightarrow \infty} \frac{\mu(E, t)}{\ln |F_S(E, t)/E|} \quad (16)$$

This limit can be found by using a method which is due to Vernov⁷⁾. Indeed, it follows that

$$\left| \int_{E_0}^E \mu(E', t) \frac{dE'}{E'} \right| < \text{const.} \quad (17)$$

for every $E > E_0$ (note that E_0 must be sufficiently large^{6/} and, obviously, $K_0 > \max(R(t), E_1, E_2)$). Consequently, (16) is equal to zero and (7) equals 1.

We prove now Statement 3. Since $\lim_{E \rightarrow \infty} D(E, t)$ is assumed to exist, it follows from (12) and (15) that

$$\lim_{E \rightarrow \infty} \frac{\partial \mu(E, t)}{\partial \ln E} \left| \frac{F_T(E, t)}{E} \right|$$

also exists and

$$\lim_{E \rightarrow \infty} D(E, t) = -\frac{\pi}{2} \lim_{E \rightarrow \infty} \frac{\partial \mu(E, t)}{\partial \ln E} \left| F_T(E, t)/E \right|$$

To prove that $\lim_{E \rightarrow \infty} D(E, t) = 0$, let us assume that this limit is different from zero. Then the integral

$$\int_{\ln E_0}^{\infty} \frac{\partial \mu(E', t)}{\partial \ln E'} \left| \frac{F_T(E', t)}{E'} \right| \frac{d \ln E'}{\ln E'}$$

diverges and, because of the Froissart-Martin bound, also

$$\int_{\ln E_0}^{\infty} \frac{\partial \mu(E', t)}{\partial \ln E'} \ln E' d \ln E'$$

is divergent. Integrating by parts, we transform it to

$$\lim_{E \rightarrow \infty} \left(\mu(E', t) \ln E' \Big|_{E_0}^E - \int_{E_0}^E \mu(E', t) \frac{dE'}{E'} \right).$$

But the divergence of this expression contradicts (17). Thus, $\lim_{E \rightarrow \infty} D(E, t) = 0$.

REFERENCES

1. J.B.Bronzan, Argonne Report ANL/HESP 7327 (1973),
J.B.Bronzan et al, Phys.Lett. 49B (1974) 272.
2. G.K.Eichmann, J.Dronkers, Phys.Lett. 52B (1974) 428.
3. J.Heidrich, E.Kazes, Lett.al Nuovo Cimento 12 (1975) 365.
4. V.N.Gribov, A.A.Migdal, Yad.Fiz. 8 (1968) 1002.
5. A.Martin, Scattering theory: unitarity, analyticity and crossing (Springer Verlag, Berlin-Heidelberg-New York 1969).
6. J.Fischer et al, Phys.Rev. D13 (1976) 133.
7. Yu.S.Vernov, Yad.Fiz. 10 (1969) 176.

Received by Publishing Department
on May 20, 1976.

**SUBJECT CATEGORIES
OF THE JINR PUBLICATIONS**

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

D10-6142	Proceedings of the International Symposium on Data Handling of Bubble and Spark Chambers. Dubna, 1971.	564 pg	9.96
D13-6210	Proceedings of the Vth International Symposium on Nuclear Electronics. Warsaw, 1971.	372 pg	6.36
D1-6349	Proceedings of the IVth International Conference on High Energy Physics and Nuclear Structure. Dubna, 1971.	670 pg	9.92
D-6840	Proceedings of the Second International Symposium on High Energy and Elementary Particle Physics (Strbske Pleso, CSSR, 1972).	398 pg	6.78
D2-7161	Proceedings of the III International Seminar on Non-Local Quantum Field Theory. Alushta, 1973.	280 pg	5.00
D1,2-7411	Deep Inelastic and Many-Body Processes. Dubna, 1973.	507 pg	8.76
D13-7616	Proceedings of the VII International Symposium on Nuclear Electronics. Budapest, 1973.	372 pg	5.62
P1,2-7642	Proceedings of the International School of Young Physicists on High Energy Physics. Gomel, 1973.	623 pg	10.53
D10-7707	Meeting on Programming and Mathematical Methods for Solving the Physical Problems. Dubna, 1973.	564 pg	8.23
D1,2-7781	Proceedings of the III International Symposium on High Energy and Elementary Particle Physics. Sinaia, Romania, 1973.	478 pg	7.29
D3-7991	Proceedings of the II International School on Neutron Physics. Alushta, 1974.	554 pg	4.35
D1,2-8405	Proceedings of the IV International Symposium on High Energy and Elementary Particle Physics. Varna, 1974.	376 pg	3.89
D10,11-8450	International School on Use of Computers in Nuclear Research. Tashkent, 1974.	465 pg	4.52

P1,2-8529	The International School-Seminar of Young Scientists. Actual Problems of Elementary Particle Physics. Sochi, 1974.	582 pg	4.70
D6-8846	XIV Symposium on Nuclear Spectroscopy and Nuclear Theory. Dubna, 1975.	180 pg	3.70
E2-9086	Proceedings of the 1975 JINR-CERN School of Physics. Alushta, 1975.	462 pg	16.00
D13-9164	Proceedings of the International Meeting on Proportional and Drift Chambers. Dubna, 1975.	344 pg	10.00
D1,2-9224	Proceedings of the IV International Seminar on High Energy Physics Problems. Dubna, 1975.	307 pg	4.80
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973, v. I.	666 pg	21.00
	Proceedings of the VI European Conference on Controlled Fusion and Plasma Physics. Moscow, 1973, v. II.	466 pg	15.00
D13-9287	Proceedings of the VIII International Symposium on Nuclear Electronics. Dubna, 1975	469 pg.	6.6
D1,2-9342	Proceedings of the V International Symposium on High Energy and Elementary Particle Physics. Warsaw, 1975.	339 pg.	6.6.

Orders for the above-mentioned books can be sent at the address:

Publishing Department, JINR
 Head Post Office, P.O. Box 79
 101000 Moscow, USSR

Conditions of Exchange

The preprints and communications of the Joint Institute for Nuclear Research are distributed free of charge on the mutual exchange basis to the universities, institutes, libraries, scientific groups of more than 50 countries.

Besides the regular distribution on the exchange basis, the Publishing Department fulfils annually about 4000 individual requests for our preprints and communications. The index of our publication must be obligatory indicated in such requests.

Addresses

Letters on all the questions concerning the exchange of publications as well as requests for individual publications are to be sent at the address:

*Publishing Department
Joint Institute for
Nuclear Research
Head Post Office,
P.O. Box 79
101000 Moscow,
U.S.S.R.*

We kindly ask to send all the publications on the exchange basis and also free of charge subscriptions to scientific journals at the address:

*Scientific-Technical Library
Joint Institute for
Nuclear Research
Head Post Office
P.O. Box 79
101000 Moscow,
U.S.S.R.*

Издательский отдел Объединенного института ядерных исследований.
Заказ 21498. Тираж 680. Уч.-изд. листов 0,47.
Редактор Э.В.Ивашевич. Подписано к печати 28.5.76 г.
Корректор Т.Е.Жильцова