


BR-801615



CBPF
CENTRO BRASILEIRO
DE PESQUISAS FÍSICAS

A0021/76

JUL, 1976

GHOST BASIS FOR NEUTRINO

M. Novello

Centro Brasileiro de Pesquisas Físicas

RIO DE JANEIRO

BRASIL

GHOST BASIS FOR NEUTRINO

M. Novello

*Centro Brasileiro de Pesquisas Físicas
Av. Wenceslau Brás, 71 - Rio de Janeiro, Brasil*

Recently some solutions of Dirac's equation in gravitational fields have been presented^[1] with the curious property of having a null energy-momentum tensor but a non-null current density. Some authors have interpreted this as being a sort of ghost neutrinos in the sense that they can be acted upon by a gravitational field although they cannot create gravitation.

One's first reaction to these solutions may be a rejection of them on physical grounds. However, the fact that they seem to be present in any kind of geometry makes worthwhile a less superficial investigation. In order to be able to study some details of these ghost we have to find not only one but a class of these solutions in a given geometry.

Let us restrict our comments on the present paper to the neutrino cosmological model recently founded by myself and I.D. Soares^[2].

This model contains arbitrary functions of the time-coordinate which permits to create such an infinity class of ghost solutions.

The line element has the form

$$(1) \quad ds^2 = dt^2 - 2A(t)dzdt - C^2(t)(dx^2 + dy^2)$$

In this geometry, for a spinor ψ which depends only on t , Dirac's equation assumes the form:

$$(2) \quad (\gamma^0 - \gamma^1)\dot{\psi} - \frac{1}{2} \frac{\dot{A}}{A} (\gamma^1 - \gamma^0)\psi + \frac{\dot{C}}{C} (\gamma^0 - \gamma^1)\psi = 0$$

This equation is identically satisfied by choosing ψ to have the form^(*)

$$(3) \quad \psi = \begin{pmatrix} \phi \\ \sigma' \phi \end{pmatrix}$$

ϕ is a 2-component spinor with arbitrary dependence on t and $\sigma' = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

The energy-momentum tensor of the neutrino is given by

$$(4) \quad T_{MN} = \bar{\psi} \gamma_{(M} D_{N)} \psi - D_{(M} \bar{\psi} \gamma_{N)} \psi$$

For the local tetrads we choose:

$$e^0_{(0)} = 1, \quad e^0_{(1)} = e^1_{(1)} = -A(t) \quad \text{and} \quad e^2_{(2)} = e^3_{(3)} = C(t)$$

(coordinate indices are written under parenthesis).

In this system of tetrads the only non-null components of T_{MN} are

(*) Our convention is the same as in ref [2].

$$\gamma^K = \begin{pmatrix} 0 & \sigma^K \\ -\sigma^K & 0 \end{pmatrix}; \quad \gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}; \quad \text{the } \sigma^K \text{ are Pauli matrices.}$$

$$(5) \quad T_{00} = T_{11} = -T_{01} = -8 \operatorname{Im}(\dot{\phi} \dot{\phi}^{\dagger})$$

(a dot means t-derivative).

Einstein's equation gives a unique relation between the metrical coefficients A, C and the spinor ψ :

$$(6) \quad \frac{\ddot{C}}{C} - \frac{\dot{A}}{A} \frac{\dot{C}}{C} = -2 \operatorname{Im}(\dot{\phi} \dot{\phi}^{\dagger})$$

From expression (5) the condition of ghostness is

$$(7) \quad \operatorname{Im}(\dot{\phi} \dot{\phi}^{\dagger}) = 0$$

We remark that given two ghost solution ψ_1 and ψ_2 the linear combination $\alpha\psi_1 + \beta\psi_2$ is still a solution of Dirac's equation but it is not necessarily a ghost for arbitrary complex numbers α and β . This is a direct consequence of linearity of Dirac's equation and non-linearity of the energy-momentum tensor.

There is two independent types of ghost spinor present in our geometry. They are

$$(8a) \quad U_K = T_K(t) U_I$$

$$(8b) \quad W_L = T_L(t) U_{II}$$

in which $U_I = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, $U_{II} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, T_K is an arbitrary real function of time.

By choosing a set of functions $\{T_K\}$ to constitute a basis for functions of t of a given class we can develop any spinor of our geometry into an infinity sum of ghost spinors. Indeed, we can write

$$(9) \quad \psi(t) = \sum_K \alpha_K U_K + \sum_K \beta_K W_K$$

in which α_K, β_K are complex numbers.

The energy of the field is given by

$$(10) \quad E = \sum_K \sum_L \text{Im}(\alpha_K^* \alpha_L + \beta_K^* \beta_L) T_K \dot{T}_L$$

It seems worthwhile to remark that these neutrinos do not have in general a definite helicity. However, if we insist that they must have helicity $\epsilon = \pm 1$ then the β 's and α 's are related by $\beta_K = \epsilon \alpha_K$.

Expression (10) tells us the following:

- (i) In general, by adding ghost neutrinos we do not obtain a ghost state.
- (ii) Any neutrino that generates the above geometry is a linear combination of infinity ghost states.

In a sense we could say that the state of the neutrino that creates curvature is composed of ghost-states each of which do not create curvature.

ACKNOWLEDGEMENTS

I would like to thank Dr. S. JOFFILY for many discussions on the subject.

REFERENCES

- [1] C. COLLINSON - P. MORRIS - Nuovo Cimento 16B - 273(1973).
T.M. DAVIS - J.R. RAY - Phys. Rev. D - (331), Vol. 9 ,
No. 2, 1974.
- [2] M. NOVELLO - I. DAMIÃO SOARES - Phys. Lett. Vol. 56A no.
6 (1976) 431.



O CBPF PUBLICA PERIODICAMENTE AS DUAS SÉRIES DE RELATÓRIO A SEGUIR.

SÉRIE A: PRÉ-PUBLICAÇÃO DE RELATÓRIOS DE TRABALHOS QUE NECESSITAM DE RÁPIDA CIRCULAÇÃO.

SÉRIE B TRABALHOS ORIGINAIS, MONOGRAFIAS, TABELAS E OUTROS DADOS NÃO DESTINADOS A PUBLICAÇÃO.

CBPF PUBLISHES PERIODICALLY THE FOLLOWING TWO SERIES OF REPORTS:

SERIES A: PRE-PUBLICATION OF REPORTS OF PAPERS NEEDING QUICK CIRCULATION.

SERIES B: ORIGINAL PAPERS, MONOGRAPHS, TABLES AND OTHER DATA WHICH ARE NOT INTENDED TO BE PUBLISHED ELSEWHERE.

PEDIDOS DE CÓPIA DESTES RELATÓRIOS FAVOR DIRIGIR-SE A:
REQUEST FOR COPY OF THIS REPORT PLEASE ADDRESS TO:

BIBLIOTECÁRIA - CHEFE
CBPF
AV. WENCESLAU BRAZ, 71 FUNDOS, ZC-82
RIO DE JANEIRO, RJ 20.000
BRASIL