

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА



5478.03392

E17 - 10550

T.D.Palev

A11

**LIE ALGEBRAICAL ASPECTS
OF QUANTUM STATISTICS.**

UNITARI QUANTIZATION (A-QUANTIZATION)

1977

E17 - 10550

T.D.Palev*

**LIE ALGEBRAICAL ASPECTS
OF QUANTUM STATISTICS.**

UNITARI QUANTIZATION (A-QUANTIZATION)

*Submitted to "Communications
in Mathematical Physics"*

*Permanent address: Institute for
Nuclear Research and Nuclear Energy,
Boul. Lenin 72, Sofia 13, Bulgaria.

Палев Ч.Д.

E17 - 10550

Ли-алгебраические аспекты квантовой статистики.
Унитарное квантование (A -квантование)

Показано, что аксиомы вторичного квантования могут в принципе удовлетворяться операторами рождения и уничтожения, порождающими (в случае n пар таких операторов) алгебру Ли A_n группы $SL(n+1)$. Введено понятие пространства Фока. Найлены матричные элементы этих операторов.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1977

Palev T.D.

E17 - 10550

Lie Algebraical Aspects of Quantum Statistics.
Unitary Quantization (A -Quantization)

It is shown that the second quantization axioms can, in principle, be satisfied with creation and annihilation operators generating (in the case of n pairs of such operators) the Lie algebra A_n of the group $SL(n+1)$. A concept of the Fock space is introduced. The matrix elements of these operators are found.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1977

1. INTRODUCTION

In the present paper we study some of the possible generalizations of the quantum statistics and more precisely of the second quantization procedure from a Lie algebraical point of view. The consideration is made in the framework of the Lagrangian field theory, however the results can easily be extended to other cases, e.g., to nuclear or solid state physics.

As is known^{/1/}, the ordinary quantum statistics can be considerably generalized if one quantizes the fields according to a weaker system of axioms, abandoning the usually accepted C-number postulate, i.e., the requirement for the commutator or the anticommutator of two fields to be a C-number. In this case the anticommutation relations between the fermi creation and annihilation operators f_i^+ and f_i^-*

$$\{f_i^\xi, f_j^\eta\} = \frac{1}{4}(\xi - \eta)^2 \delta_{ij} \quad (1)$$

can be replaced by a weaker system of double commutation relations for the so-

*Throughout the paper the indices $\xi, \eta, \epsilon, \delta$ take values \pm or ± 1 ; $\{x, y\} \equiv xy + yx$ and $[x, y] \equiv xy - yx$.

called parafermi operators b_i^\pm , namely

$$[[b_i^\xi, b_j^\eta], b_k^\epsilon] = \frac{1}{2}(\eta - \epsilon)^2 \delta_{jk} b_i^\xi - \frac{1}{2}(\xi - \epsilon)^2 \delta_{ik} b_j^\eta. \quad (2)$$

The commutation relations (2) exhibit some remarkable Lie algebraical properties. It turns out that the parafermi operators generate the algebra of the orthogonal group $^{/2/}$. To make the statement more precise, consider a finite number of operators b_1^\pm, \dots, b_n^\pm . Then the linear envelope over C of the operators

$$b_i^\xi, [b_j^\eta, b_k^\epsilon], \quad i, j, k = 1, \dots, n \quad (3)$$

is isomorphic to the classical Lie algebra B_n of the orthogonal group $SO(2n+1)^{/3/}$.

There exists one-to-one correspondence between the representations of B_n and the representations of n pairs of parafermi operators $^{/4/}$. Therefore the parafermi quantization is actually a quantization according to representations of the algebra of the orthogonal group in odd dimension and therefore may be called an odd-orthogonal quantization. This is an important point, a first hint that the group theory can in principle be relevant for the quantum statistics.

The algebras $B_n, n=1, 2, \dots$, constitute one of the four infinite series of the so-called classical Lie algebra. In the Cartan notation (which we follow) they are denoted as A_n, B_n, C_n and D_n for algebras of rank $n, n=1, 2, \dots$. The corresponding groups $SL(n), SO(2n+1), Sp(2n)$ and $SO(2n)$ are well known and therefore we do not define them here.

Once the Lie algebraical structure of the parafermi statistics is established, it is natural to ask whether one can quantize according to representations of the other classical Lie algebras. In the present paper we consider this question in connection with the algebra of the unimodular group.

In Sect. 3 we determine the concept of A statistics, i.e., statistics with creation and annihilation operators (a -operators) that generate the algebra of the unimodular group. Next (Sec. 4) we define the Fock spaces $W_p, p=1,2,\dots$, and the selection rules for the A-statistics. The integer p , called the order of the statistics, has well defined physical meaning: this is the maximal number of particles that can exist simultaneously (lemma 4). In Sec. 5 we calculate the matrix elements of the a -operators. In the limit $p \rightarrow \infty$ the a -operators reduce (up to a constant) to Bose operators.

The mathematics used in the paper is mainly of Lie algebraical nature. In order to introduce the notation and to make the exposition reasonably self-consistent, we collect in the next section some definitions and properties from the Lie algebra theory.

2. PRELIMINARIES AND NOTATIONS

Let A be a semi-simple complex Lie algebra of rank n , \mathcal{K} - its Cartan subalgebra. By $\omega_i, e_{\omega_i}, i=1,2,\dots,p$ we denote the roots and the root vectors of A . The roots ω_i

are vectors from the conjugate space \mathcal{K}^* of \mathcal{K} . Sometimes it is convenient to consider them also as vectors from \mathcal{K} using the fact that every linear functional $\lambda \in \mathcal{K}^*$ can be uniquely represented in the form

$$\lambda(h) = (h, \lambda) \quad \forall h \in \mathcal{K}. \quad (4)$$

Here $(,)$ is the Cartan-Killing form on \mathcal{A} and $\lambda \in \mathcal{K}^*$. The mapping

$$\theta: \lambda \rightarrow \lambda^* \equiv \theta\lambda \quad (5)$$

of \mathcal{K}^* on \mathcal{K} is one-to-one. From now on we consider the roots or any other linear functionals either as elements from \mathcal{K}^* or from \mathcal{K} , denoting them in both cases by the same symbol (i.e., for λ we write also λ).

With this agreement we can write

$$[h, e_{\omega_i}] = \omega_i(h) e_{\omega_i} = (h, \omega_i) e_{\omega_i} \quad \forall h \in \mathcal{K}. \quad (6)$$

The Cartan-Killing form defines a scalar product in the space \mathcal{K}^r which is the real linear envelope of all roots; $\mathcal{K} = \mathcal{K}^r + i\mathcal{K}^r$. Let h_1, \dots, h_n be an arbitrary covariant basis in \mathcal{K}^r (and hence a basis in \mathcal{K}). The root ω_i is said to be positive (negative) if its first non-zero coordinate is positive (negative). The simple roots, i.e., those positive roots which cannot be represented as a sum of other positive roots, constitute a basis in \mathcal{K} . Any positive (negative) root is a linear combination of simple roots with positive (negative) integer coefficients.

Consider an arbitrary finite-dimensional irreducible A -module W (i.e., a space where a finite-dimensional irreducible representation of A is realized). The basis x_1, \dots, x_N in W can always be chosen such that

$$hx_i = \lambda_i (h)x_i = (h, \lambda_i)x_i \quad \forall h \in \mathfrak{H}, i = 1, \dots, N. \quad (7)$$

Thus, to every basic vector $x_i \in W$ there corresponds an image $\lambda \in \mathfrak{H}$ (or \mathfrak{K}). The vectors x_i are the weight vectors and their images the weights of the A -module W . The mapping $\tau: x_i \rightarrow \lambda_i$ is surjective and the number of the vectors $\tau^{-1}(\lambda_i)$ is called multiplicity of the weight λ_i . Let e_ω be a root vector and λ_i be the weight of x_i . Then $e_\omega x_i$ is either zero or a weight vector with weight $\omega + \lambda_i$. The A -module W contains a unique (up to multiplicative constant) weight vector x_Λ with properties $e_{\omega_i} x_\Lambda = 0$ for all positive roots $\omega_i > 0$. The weight Λ of x_Λ is the highest weight of W . The multiplicity of Λ is unity and W is spanned over all vectors

$$e_{\omega_{i_1}} e_{\omega_{i_2}} \dots e_{\omega_{i_m}} x_\Lambda \quad m = 1, 2, \dots, \quad (8)$$

where $\omega_{i_1}, \dots, \omega_{i_m}$ are negative roots. Therefore an arbitrary weight λ is of the form

$$\lambda = \Lambda - \sum_{\omega_i > 0} k_i \omega_i \quad (9)$$

with k_i positive integers and sum over positive (or only simple) roots.

Let π_1, \dots, π_n be the simple roots of A . Then for an arbitrary weight λ the

n -tuple $[\lambda_1, \lambda_2, \dots, \lambda_n]$ has integer co-ordinates defined as

$$\lambda_i = \frac{2(\lambda, \pi_i)}{(\pi_i, \pi_i)} \quad i = 1, 2, \dots, n. \quad (10)$$

The n -tuple $[\Lambda_1, \dots, \Lambda_n]$ corresponding to the highest weight Λ has non-negative co-ordinates, and it defines the irreducible representation of A in W up to equivalence. On the contrary, to every vector $\Lambda \in \mathfrak{H}$, such that $\Lambda_1, \dots, \Lambda_n$ defined from (10) are non-negative integers, there corresponds an irreducible A -module. Thus, there exists a one-to-one correspondence between the irreducible (finite-dimensional) A -modules and the set $[\Lambda_1, \dots, \Lambda_n]$ of non-negative integers. We call $\lambda_1, \dots, \lambda_n$ canonical co-ordinates of λ .

Define an F -basis f_1, f_2, \dots, f_n in \mathfrak{H} (or in \mathfrak{K}) as follows

$$f_i = \frac{2}{(\pi_i, \pi_i)} \pi_i \quad i = 1, \dots, n \quad (11)$$

and let $K = \{f^1, f^2, \dots, f^n\}$ be the corresponding dual basis, i.e., $f^i(f_j) = (f^i, f_j) = \delta_j^i$. For an arbitrary $\lambda \in \mathfrak{H}$ we have

$$\lambda = \sum_i \lambda(f_i) f^i = \sum_i \frac{2(\lambda, \pi_i)}{(\pi_i, \pi_i)} f^i \quad (12)$$

and therefore in the K -basis the co-ordinates of every weight λ coincide with its canonical co-ordinates.

By means of the F -basis one can easily calculate the canonical co-ordinates of an arbitrary weight λ . Indeed

$$f_i x_\lambda = \lambda(f_i) x_\lambda = \frac{2(\lambda, \pi_i)}{(\pi_i, \pi_i)} x_\lambda \quad (12')$$

and therefore the i -th canonical co-ordinate λ_i of λ is an eigenvalue of f_i on x_λ . More generally, if h_1, \dots, h_n is an arbitrary covariant basis in \mathfrak{K} , then the covariant coordinates $(\lambda_1, \dots, \lambda_n)$ of the weight λ , i.e., the co-ordinates of λ in the dual (or contravariant) basis h^1, \dots, h^n are determined from the relation

$$h_i x_\lambda = \lambda(h_i) x_\lambda = \lambda_i x_\lambda. \quad (13)$$

An important property of the set Γ of all weights is its invariance under the Weyl group S which is a group of transformations of \mathfrak{K} . $S = \{S_{\omega_i} \mid \omega_i \text{ - roots of } \Lambda\}$ is a finite group, its elements labelled by the roots ω_i of Λ are defined as follows:

$$S_{\omega_i} \cdot h = h - \frac{2(h, \omega_i)}{(\omega_i, \omega_i)} \omega_i \quad \forall h \in \mathfrak{K}. \quad (14)$$

The set Γ of all weights is characterized by the following statement: if $\lambda \in \Gamma$, then

$$S_{\omega_i} \lambda = \lambda + j\omega_i \in \Gamma, \quad j \text{ - integer} \quad (15)$$

and Γ contains also the weights

$$\lambda, \lambda + \omega_i, \lambda + 2\omega_i, \dots, \lambda + j\omega_i. \quad (16)$$

All weights that can be connected by transformations of the Weyl group are called equivalent. They have the same multiplicity. Among the equivalent weights there exists only one weight, the dominant one, the canonical co-ordinates of which are nonnegative integers.

3. UNITARY QUANTIZATION (A -QUANTIZATION)

In the case of ordinary statistics the second quantization in the Lagrangian field theory can be performed in different equivalent ways. One can start, for instance, from the equal-time commutation relations. For generalizations we wish to consider, it is more convenient to follow the quantization procedure accepted by Bogolubov and Shirkov ^{6/}.

Apart from the fact that the fields become operators and the requirement for relativistic invariance, their approach is essentially based on what we call a main quantization postulate: the energy-momentum vector P^m and the angular-momentum tensor M^{mn} , $m,n=0,1,2,3$, are expressed in terms of the operator-fields by the same expressions as in the classical case.

It follows from this postulate, together with the requirement that the field transforms according to unitary representations of the Poincare group and the compatibility of the transformation properties of the field and the state vectors, that the field $\Psi(x)$ satisfies the commutation relation

$$[P^m, \Psi(x)] = -i\partial^m \Psi(x). \quad (17)$$

This relation expresses (in infinitesimal form) the translation invariance of the theory.

To proceed further, it is convenient to pass to the discrete notation in momentum space. Consider a field $\Psi(x)$ with mass m locked in a cube with edge L . For

the eigenvalues k_n^m of the 4-momentum P^m , $m=0,1,2,3$, one obtains

$$k_n^a = \frac{2\pi}{L} n^a, \quad k_n^0 = \sqrt{m^2 + \left(\frac{2\pi}{L}\right)^2 [(n^1)^2 + (n^2)^2 + (n^3)^2]}, \quad (13)$$

where $n = (n^1, n^2, n^3)$, $a=1,2,3$ and n^a runs over all non-negative integers. In momentum space the relation (17) reads as follows

$$[P^m, a_i^\pm] = \pm k_i^m a_i^\pm, \quad (19)$$

where $a_i^+(a_i^-)$ are the corresponding to $\Psi(x)$ creation and annihilation operators and the index i replaces all discrete indices (n spin, charge, etc.).

The commutation relations between the creation and annihilation operators are usually derived from the translation invariance law in momentum space (19). We call it the initial quantization equation (IQE). To determine the commutation relations one has to specify one more point. Up to now nothing was said about the order of the creation and annihilation operators that enter into P^m . In the ordinary theory it is usually accepted that the dynamical variables are written in a normal-product form and therefore for a fermi field this give

$$P^m = \sum_i k_i^m f_i^+ f_i^-, \quad (20)$$

where $f_i^+(f_i^-)$ are the Fermi creation (annihilation) operators (1). One can easily check that the initial quantization equation (with $a_i^\pm = f_i^\pm$) is compatible with the anticommutation relations (1). This is, however, not the case for the parafermi

operators (2), apart from the case of their Fermi representation. The parafermi statistics cannot be derived from the normal-product form of the dynamical variables. In order to fulfil (19) Green chose another ordering of the operators in P^m and in particular for spinor fields we wrote ^{/1/}:

$$P^m = \frac{1}{2} \sum_i k_i^m (b_i^+, b_i^-). \quad (21)$$

We see that the ordering of the operators in the 4-momentum is closely related to the corresponding statistics. It is natural to expect therefore that any other generalization of the statistics may require new expressions for P^m . In order to get a feeling as to how one can modify P^m , we now proceed to derive the parafermi statistics in such a way that later on it will be possible to generalize the idea to other cases.

We start with the expression (20). In order to use a proper Lie algebraical language (finite-dimensional Lie algebras), suppose that the sum in (10) is finite,

$$P^m = \sum_{i=1}^n k_i^m f_i^+ f_i^-. \quad (22)$$

This is only an intermediate step. In the final results we let $n \rightarrow \infty$.

As we have already mentioned, the set f_1^+, \dots, f_n^+ of Fermi creation and annihilation operators (1) generates one particular representation (we call it the Fermi representation) of the algebra B_n . We put now the question: can the expression (22) be written in such a form that the initial

quantization equation (19) will hold for the Fermi operators considered as generators of B_n , i.e., independently of the fact we are staying in one particular representation of B_n - the Fermi one. The Lie algebraical reason why (19) does not hold for the parafermi operators is clear. It is due to the fact that the 4-momentum (22) does not belong to B_n since it contains product of b_i^+ and b_i^- , which is not a Lie-algebraical operation. Therefore the IQR, considered as a commutation relation, is not preserved for other representation of B_n . If however, the 4-momentum together with the creation and annihilation operators can be embedded in a Lie algebra, so that in the Fermi case P^m reduces to (22), then the IQR (19) will hold for any other representation of this algebra.

For this purpose we rewrite the 4-momentum (22) in the following identical form

$$P^m = \sum_i k_i^m \left(\frac{1}{2} \{f_i^+, f_i^-\} + \frac{1}{2} \{f_i^-, f_i^+\} \right). \quad (23)$$

Consider the Lie algebra generated from f_i^\pm, \dots, f_n^\pm and $\{f_i^+, f_i^-\}$. Since $\{f_i^+, f_i^-\} = 1$, we obtain the algebra $B_n \otimes I$, where I is the one-dimensional commutative center. Now $P^m \in B_n \otimes I$ and therefore the commutation relation (19) holds for any other representation. In other words, if we substitute in (23) $f_i^\pm \rightarrow b_i^\pm$ and $\{f_i^+, f_i^-\} \rightarrow 1$, i.e., put

$$P^m = \sum_i k_i^m \left(\frac{1}{2} [b_i^+, b_i^-] + \frac{1}{2} I \right), \quad (24)$$

where I is the generator of the center of $B_n \otimes I$, then the initial quantization

condition (19) will be fulfilled for any representation of $B_n \otimes I$.

The operator

$$Q^m = \frac{1}{2} \sum_i k_i^m \quad (25)$$

commutes with all creation and annihilation operators and hence with all elements of $B_n \otimes I$. Therefore it is a constant within every irreducible representation and in the particular case of parafermi statistics the second term in (24) can be omitted. Thus, we obtain the expression (21) for P^m , postulated by Green from the very beginning.

We shall now apply a similar approach for the algebra A_n of the unimodular group $SL(n+1)$. The nontrivial part is to find an analogue of the Fermi operators, i.e., operators a_i^\pm that generate some representation of A_n and fulfil the initial quantization equation (19) with 4-momentum written (in this particular representation) in a normal-product form. Then we shall apply the above procedure to enlarge the class of admissible representation.

First we recall some properties of A_n . We consider A_n as a subalgebra of the algebra $gl(n+1)$ of the general linear group $GL(n+1)$. The algebra $gl(n+1)$ may be determined as a linear envelope of the generators e_{ij} , $i, j = 0, 1, \dots, n$, that satisfy the commutation relations

$$[e_{ij}, e_{kl}] = \delta_{jk} e_{i\ell} - \delta_{\ell i} e_{kj}, \quad i, j, k, \ell = 0, 1, \dots, n. \quad (26)$$

Let \mathcal{K} and $\tilde{\mathcal{K}}$ be the Cartan subalgebras of A and $gl(n+1)$, resp. Denote by $\text{env } X$ the

linear envelope of an arbitrary set X .
 In terms of the $gl(n+1)$ generators we have:

$$gl(n+1) = \text{env}\{e_{ij} \mid i, j = 0, 1, \dots, n\},$$

$$A_n = \text{env}\{e_{ii} - e_{jj}, e_{ij} \mid i \neq j = 0, 1, \dots, n\}, \quad (27)$$

$$\tilde{\mathcal{H}} = \text{env}\{h_i \mid h_i = e_{ii}, i = 0, 1, \dots, n\},$$

$$\mathcal{H} = \text{env}\{h_i - h_j \mid h_i = e_{ii}, i = 0, 1, \dots, n\}.$$

For a covariant basis in $\tilde{\mathcal{H}}$ we choose the vectors $(h_i = e_{ii})$

$$h_0, h_1, \dots, h_n. \quad (28)$$

The algebra $gl(n+1)$ is not semi-simple. Its Cartan-Killing form is degenerate and does not determine a scalar product on $\tilde{\mathcal{H}}$. It is convenient to introduce a metric in $\tilde{\mathcal{H}}$ with the relation

$$(h_i, h_j) = 2(n+1)\delta_{ij}. \quad (29)$$

Restricted on \mathcal{H} this metric coincides with the Cartan-Killing form of A_n .

From (26) and (29) one obtains

$$[h, e_{ij}] = (h_i h^i - h_j h^j) e_{ij} \quad \forall h \in \mathcal{H}, i \neq j = 0, 1, \dots, n, \quad (30)$$

where h^0, h^1, \dots, h^n is the contravariant (i.e., dual to h_0, h_1, \dots, h_n) basis in $\tilde{\mathcal{H}}$. Hence the generators $e_{ij}, i \neq j = 0, 1, \dots, n$ are the root vectors of A_n . The correspondence with their roots is

$$e_{ij} \rightarrow h^i - h^j \quad i \neq j = 0, 1, \dots, n. \quad (31)$$

In the basis (28) the generators

$$e_{ij}, \quad i < j \quad (i > j), \quad i, j = 0, 1, \dots, n \quad (32)$$

are the positive (negative) root vectors of A_n . The simple roots are

$$\pi_i = h^{i-1} - h^i, \quad i = 1, \dots, n. \quad (33)$$

Therefore the \mathbb{F} -basis (11) in this case reads as

$$f_i = \frac{2}{(\pi_i, \pi_i)} \pi_i = h_{i-1} - h_i, \quad i = 1, \dots, n. \quad (34)$$

We are now ready to define the analogue of the Fermi operators. Let E_{ij} , $i, j = 0, 1, \dots, n$ be $(n+1)$ -square matrix with 1 on the intersection of i -th row and j -th column and zero elsewhere. Clearly the mapping

$$\pi: e_{ij} \rightarrow E_{ij}, \quad i, j = 0, 1, \dots, n \quad (35)$$

determines a representation of $\mathfrak{gl}(n+1)$ and hence its restriction on A_n gives a representation of A_n .

The operators

$$A_i^+ = E_{i0}, \quad A_i^- = E_{0i} \quad i = 1, 2, \dots, n \quad (36)$$

generate the algebra A_n (in the above representation) since

$$\begin{aligned} [A_i^+, A_j^-] &= E_{ij}, \\ [A_k^+, A_k^-] &= E_{kk} - E_{00}, \quad i \neq j, \quad i, j, k = 1, \dots, n. \end{aligned} \quad (37)$$

Moreover for the commutation relations between A_1^\pm, \dots, A_n^\pm and the operator

$$P^m = \sum_i k_i^m A_i^+ A_i^- \quad (38)$$

we obtain the right expression:

$$[P^m, A_i^\pm] = \pm k_i^m A_i^\pm. \quad (39)$$

The operators A_1^ξ, \dots, A_n^ξ satisfy the initial quantization equation and can be considered as creation ($\xi = +$) and annihilation ($\xi = -$) operators.

The commutation relation (39) does not hold for other representations of A_n . In order to extend the class of admissible representation, we represent the 4-momentum (38) like in the Fermi case, in the form

$$P^m = \sum_i k_i^m (A_i^+ A_i^- + E_{00}). \quad (40)$$

Consider now the Lie algebra generated from the operators A_1^\pm, \dots, A_n^\pm and E_{00} . One can easily show, it is the algebra $gl(n+1) = A_n \oplus 1$. Since $P^m \in gl(n+1)$, the initial quantization equation (39) holds for any other representation of $gl(n+1)$. Hence we may define representation independent creation and annihilation operators as follows

$$a_i^+ = e_{i0}, \quad a_i^- = e_{0i}, \quad i = 1, \dots, n. \quad (41)$$

In this case we have to postulate for P^m the expression

$$P^m = \sum_i k_i^m (a_i^+ a_i^- + e_{00}). \quad (42)$$

The operators a_i^\pm are root vectors of A_n . The correspondence with their roots is

$$a_i^\pm \leftrightarrow \mp (h^0 - h^i), \quad i = 1, \dots, n, \quad (43)$$

and therefore the creation (annihilation) operators are negative (positive) root vectors. Since any other root $h^i - h^j$, $i \neq j = 1, \dots, n$

$$h^i - h^j = (h^0 - h^j) - (h^0 - h^i)$$

is a sum of the roots of a_j^- and a_i^+ , the creation and annihilation operators generate the algebra A_n .

The commutation relations of A_n can be written in terms of a_i^\pm only. From (26) we obtain

$$[[a_i^+, a_j^-], a_k^+] = \delta_{kj} a_i^+ + \delta_{ij} a_k^+, \quad (44)$$

$$[[a_i^+, a_j^-], a_k^-] = -\delta_{ki} a_j^- - \delta_{ij} a_k^-$$

$$[a_i^+, a_j^+] = [a_i^-, a_j^-] = 0.$$

Definition 1. The operators a_i^ξ , $i=1, 2, \dots$ satisfying the commutation relation (44) are called a -operators and the corresponding quantization (statistics) unitary or A -quantization (statistics).

We observe that the equal-frequency operators commute with each other. This property helps a lot in all calculations with the a -operators.

4. FOCK SPACES FOR THE a -OPERATORS

We now proceed to study those representations of the a -creation and annihilation operators that possess the main features of the Fock space representations in the ordinary quantum mechanics. We continue to consider a finite set of operators. The extension of the results to infinite (including continuum) number of a -operators will be evident.

Definition 2. Let a_1^ξ, \dots, a_n^ξ be a -creation ($\xi=+$) and annihilation ($\xi=-$) operators. The A_n -module W is said to be a Fock space of the algebra A_n if it fulfils the conditions:

1. Hermiticity condition

$$(a_i^+)^* = a_i^-, \quad i = 1, \dots, n. \quad (45)$$

Here $*$ denotes hermitian conjugation operation.

2. Existence of vacuum. There exists a vacuum vector $|0\rangle \in W$ such that

$$a_i^- |0\rangle = 0, \quad i = 1, \dots, n. \quad (46)$$

3. Irreducibility. The representation space W is spanned over all possible vectors

$$a_{i_1}^+ a_{i_2}^+ \dots a_{i_m}^+ |0\rangle, \quad m \in N_0. \quad (47)$$

By N_0 we denote the set of all non-negative integers. The Fock space of A_n is called also A_n -module of Fock, Fock module of the a -operators or simply Fock module.

Lemma 1. The hermiticity condition (45) can be satisfied if and only if the A_n -module W is a direct sum of irreducible finite-dimensional modules.

Proof

The generators of the compact form $su(n+1)$ of A_n read in terms of the a -operators as follows

$$\begin{aligned}
 E_{0j} &= i(a_j^+ + a_j^-) \\
 F_{0j} &= a_j^- - a_j^+ \\
 E_{jk} &= i|a_j^+, a_k^-| + i|a_k^+, a_j^-|, \\
 F_{jk} &= |a_j^+, a_k^-| - |a_k^+, a_j^-|.
 \end{aligned}
 \tag{48}$$

Evidently the generators are antihermitian if and only if (45) holds.

As is known, the antihermitian representations of the compact forms of the classical algebras are completely reducible. The irreducible components are finite-dimensional. This proves the sufficient part. The necessity follows from the observation that the metric in any irreducible $su(n)$ -module can be introduced so that the generators are antihermitian.

From the complete reducibility and the irreducibility condition (definition 1) we conclude.

Corollary 1. The Fock spaces are finite-dimensional irreducible A_n -modules.

In the remaining part of the paper by creation and annihilation operators we

always mean a -operators. Moreover we fix the ordering of the basis of \mathfrak{H} to be (28). Then the creation and annihilation operators a_i^+, a_i^- are negative and positive root vectors. In this case the operators a_1^-, \dots, a_n^- annihilate the highest weight vector x_Λ of the Fock space and hence x_Λ is one of the candidates for a vacuum state.

Lemma 2. Let W be a Fock space of Λ_n . Up to a multiplicative constant the vacuum state is unique and coincides with the highest weight vector x_Λ of W .

Proof

First suppose the vacuum is a weight vector $x_\lambda \neq x_\Lambda$. Then the corresponding weights are also different $\lambda \neq \Lambda$. Moreover $\Lambda - \lambda$ (i.e., the vector $\Lambda - \lambda$ is positive). The irreducibility condition says there exists a polynomial $P(a_1^+, \dots, a_n^+)$ of the creation operators such that

$$x_\Lambda = P(a_1^+, \dots, a_n^+) x_\lambda. \quad (49)$$

Denote by ω_i the root of a_i^+ . From (49) we have

$$\Lambda = \lambda - \sum_{i=1}^n k_i \omega_i \quad k_i \in \mathbb{N}_0.$$

This is, however, impossible, since $\Lambda - \lambda > 0$ and $\sum k_i \omega_i < 0$. We conclude that the vacuum cannot be a weight vector different from x_Λ .

More generally, suppose $|0\rangle \in W$ is a vacuum state different from x_Λ . An arbitrary vector $x \in W$ and in particular $|0\rangle$ can be represented uniquely as a sum of weight vectors x_{λ_i} with different weights λ_i :

$$|0\rangle = \sum_{j=0}^m x_{\lambda_j}, \quad \lambda_i \neq \lambda_j \quad \text{if} \quad i \neq j. \quad (50)$$

The vectors $x_{\lambda_0}, \dots, x_{\lambda_m}$ are linearly independent. The non-zero of the vectors $a_i^- x_{\lambda_0}, \dots, a_i^- x_{\lambda_m}$ are also linearly independent, since they correspond to different weights. Hence

$$a_i^- |0\rangle = 0 \quad \text{implies} \quad a_i^- x_{\lambda_j} = 0, \quad j=0,1,\dots,m. \quad (51)$$

Let for definiteness $\lambda_0 > \lambda_1 > \dots > \lambda_m$. The vector cannot be a vacuum state if $\lambda_0 \neq \Lambda$ since clearly there exists no polynomial $P(a_1^+, \dots, a_n^+)$ such that

$$x_{\Lambda} = P(a_1^+, \dots, a_n^+) |0\rangle. \quad (52)$$

Suppose $|0\rangle = x_{\Lambda} + x_{\lambda_1} + \dots + x_{\lambda_m}$. Then (52) can be satisfied only when there exists a monomial $(a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n}$ with the property

$$x_{\lambda_1} = (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} x_{\Lambda}.$$

This is, however, impossible since for $\ell_i \neq 0$ $a_i^- x_{\lambda_1} \neq 0$ and this contradicts (51).

In the following theorem we prove one convenient criterion for the A_n -module to be a Fock space.

Theorem 1. The A_n -module W is a Fock space if and only if it is an irreducible finite-dimensional module such that

$$a_i^- a_j^+ x_{\Lambda} = 0 \quad i \neq j = 1, \dots, n. \quad (53)$$

The highest weight vector x_{Λ} is a vacuum of W .

Proof

Let W be a Fock space. Then it is finite-dimensional irreducible A_n -module (corollary 1) and the vacuum $|0\rangle = x_\Lambda$ (lemma 2). The operator $[a_i^-, a_j^+]$, $i \neq j$ is a root vector of A_n . Its root $h^j - h^i$ cannot be represented as a linear combination of the roots $-h^0 + h^i$ of the creation operators a_1^+, \dots, a_n^+ . Hence there exists no polynomial $P(a_1^+, \dots, a_n^+)$ of a_1^+, \dots, a_n^+ such that

$$[a_i^-, a_j^+] x_\Lambda = P(a_1^+, \dots, a_n^+) x_\Lambda \neq 0.$$

Since $a_i^- a_j^+ x_\Lambda \in W$ it has to be zero, $a_i^- a_j^+ x_\Lambda = 0$, $i \neq j$.

The proof of the sufficient part of the theorem is based on the Poincare-Birkhoff-Witt theorem ^{7/}: Given a Lie algebra A with basis e_1, \dots, e_N . All ordered momentals $e_1^{j_1} e_2^{j_2} \dots e_N^{j_N}$ constitute a basis of the universal enveloping algebra U of A .

Let in the irreducible finite-dimensional A_n -module W the equality (53) holds. Divide the basis elements of A_n into three groups

$$I = \{a_i^+, [a_j^+, a_k^+] | j < k; i, j, k = 1, \dots, n\} = \{e_{-1}, e_{-2}, \dots, e_{-p}\}.$$

$$II = \{a_i^-, [a_j^-, a_k^-], [a_r^-, a_s^+] | j < k; r \neq s; i, j, k, r, s = 1, \dots, n\} = \{e_1, \dots, e_q\},$$

$$III = \{\omega_k | k = 1, \dots, n\},$$

where $\omega_1, \dots, \omega_n$ is a basis in the Cartan subalgebra \mathcal{H} . Order the elements within each group in an arbitrary way. From the

irreducibility and the Poincare-Birkhoff-Witt theorem it follows that W is linearly spanned on all vectors

$$e_{-1}^{i_1} \dots e_{-p}^{i_p} e_1^{j_1} \dots e_q^{j_q} \omega_1^{k_1} \dots \omega_n^{k_n} x_\Lambda. \quad (54)$$

Since x_Λ is an eigenvector of the Cartan subalgebra and the operators from II annihilate x_Λ , the vector (54) is non-zero only if $j_1 = j_2 = \dots = j_q = 0$. Hence W is spanned on all vectors

$$P(a_1^+, \dots, a_n^+) x_\Lambda,$$

where P is an arbitrary polynomial of the creation operators. This proves that W is a Fock space with vacuum $|0\rangle = x_\Lambda$.

Now it remains to determine the irreducible A_n -modules satisfying the condition (53). In order to solve this problem, we consider first some questions from the representation theory of A_n . As we mentioned, it is convenient to consider A_n as a subalgebra of $gl(n+1)$. This possibility is based on the circumstance that the irreducible $gl(n+1)$ -modules are also A_n -irreducible. On the other hand, every irreducible representation of A_n in W can be continued in infinitely many ways to an irreducible representation of $gl(n+1)$ in the same space. For this purpose it is enough to define the operator $f_0 = h_0 + h_1 + \dots + h_n$ in W where h_0, \dots, h_n is the covariant basis (28) in \mathcal{H} . Since f_0 commutes with $gl(n+1)$, f_0 has to be a constant in W , i.e.,

$$f_0 x = \Lambda_0 x \quad \forall x \in W \quad (55)$$

with Λ_0 being an arbitrary number. Let

f_1, \dots, f_n be the \mathbb{F} -basis (34) in \mathbb{K} . Then

$$\tilde{\mathbb{F}} = \{f_0, f_1, \dots, f_n\} \quad (56)$$

defines a basis in the Cartan subalgebra $\mathfrak{h} \subset \mathfrak{gl}(n+1)$.

The eigenvalues $\Lambda_0, \Lambda_1, \dots, \Lambda_n$ of $\tilde{\mathbb{F}}$ on the highest weight vector $x_\Lambda \in W$ characterize W as an irreducible $\mathfrak{gl}(n+1)$ -module. Let $x_{\lambda_1}, \dots, x_{\lambda_N}$ be a basis of weight vectors in W . In view of (55) the A_n -weights $\lambda_1, \dots, \lambda_N$ are naturally extended to linear functionals on \mathbb{K} from the requirement $\lambda_i(f_0) = \Lambda_0$. Then for any weight vector x_λ we have

$$hx_\lambda = \lambda(h)x_\lambda = (h, \lambda)x_\lambda \quad h \in \tilde{\mathbb{K}}. \quad (57)$$

The numbers $\Lambda_0, \Lambda_1, \dots, \Lambda_n$ are co-ordinates of the highest weight Λ in the basis

$$\tilde{\mathbb{K}} = \{f^0, f^1, \dots, f^n\} \quad (58)$$

dual to $\tilde{\mathbb{F}}$. We call $\tilde{\mathbb{K}}$ a canonical basis of $\mathfrak{gl}(n+1)$ and the co-ordinates $[\Lambda_0, \dots, \Lambda_n]$ canonical co-ordinates of the $\mathfrak{gl}(n+1)$ -module W . The properties of the Weyl group we shall often use read more simply in the orthogonal contravariant basis h^0, h^1, \dots, h^n . From the equality

$$\Lambda = \sum_{i=0}^n \Lambda_i f^i = \sum_{i=0}^n L_i h^i$$

we obtain for the orthogonal co-ordinate L_0, L_1, \dots, L_n of the highest weight W the following expressions

$$\begin{aligned}
L_0 &= \frac{1}{n+1} [\Lambda_0 + n\Lambda_1 + (n-1)\Lambda_2 + \dots + 1 \cdot \Lambda_n], \\
L_1 &= L_0 - \Lambda_1, \\
L_2 &= L_0 - \Lambda_1 - \Lambda_2, \\
&\dots \\
L_n &= L_0 - \Lambda_1 - \Lambda_2 - \dots - \Lambda_n.
\end{aligned}
\tag{59}$$

Since in the A_n -module W $\Lambda_1, \dots, \Lambda_n$ are non-negative integers and Λ_0 is an arbitrary constant, it can be chosen such that all orthogonal co-ordinates L_0, L_1, \dots, L_n are integers. Moreover

$$L_0 \geq L_1 \geq L_2 \geq \dots \geq L_n. \tag{60}$$

We pass now to the main problem of this section, classification of the Fock spaces. Unless otherwise stated, the roots and the weights are represented by their orthogonal co-ordinate in the contravariant orthogonal basis h^0, h^1, \dots, h^n in \tilde{H} i.e.,

$$\lambda = (\ell_0, \ell_1, \dots, \ell_n) \equiv \sum_{i=0}^n \ell_i h^i. \tag{61}$$

Theorem 2. The irreducible A_n -module is a Fock space if and only if its highest weight is $\lambda = (p, 0, \dots, 0)$; p is an arbitrary positive integer*.

Proof

As we know (theorem 1), the Fock spaces are those and only those irreducible A_n -modules whose highest weight vectors x_λ

*The case $p=0$ corresponds to the trivial one-dimensional representation.

are annihilated by all operators $a_i^- a_j^+$, $i \neq j = 1, \dots, n$, i.e.,

$$a_i^- a_j^+ x_\Lambda = 0 \quad i \neq j = 1, \dots, n. \quad (62)$$

Since $a_i^- x_\Lambda = 0$ and $[a_i^-, a_j^+] = e_{ij}$ (62) can be replaced by the requirement

$$e_{ij} x_\Lambda = 0 \quad i \neq j = 1, \dots, n. \quad (63)$$

The generators e_{ij} are root vectors of A_n with roots (31), i.e.,

$$e_{ij} \leftrightarrow h^i - h^j, \quad i \neq j = 1, \dots, n.$$

For $i < j$ e_{ij} is positive root vector and (63) holds from the definition of x_Λ . It remains to determine those A_n -modules with highest weights

$$\Lambda = (L_0, L_1, \dots, L_n) \quad (64)$$

for which the sums

$$\Lambda + h^i - h^j, \quad i < j = 1, \dots, n \quad (65)$$

are not weights.

We shall use the properties (14) and (15) of the Weyl group S . According to (14) if $S_{h^i - h^j} \in S$ and $\lambda = (\ell_0, \dots, \ell_i, \dots, \ell_j, \dots, \ell_n)$ is a weight, then $S_{h^i - h^j} \cdot \lambda$ is also a weight. Using the scalar product (29) we have

$$\begin{aligned} S_{h^i - h^j} \cdot \lambda &= \lambda - \frac{2(\lambda, h^i - h^j)}{(h^i - h^j, h^i - h^j)} (h^i - h^j) = \\ &= (\ell_0, \dots, \ell_j, \dots, \ell_i, \dots, \ell_n). \end{aligned} \quad (66)$$

Thus, the Weyl group in this case reduces to (all possible) permutations of the orthogonal co-ordinates. For the highest weight (64) we have

$$S_{h^i - h^j} \cdot \Lambda = (L_0, \dots, L_j, \dots, L_i, \dots, L_n) = \\ = (\dots, L_i, \dots, L_j, \dots) + (L_j - L_i) \cdot (0, \dots, 0, -1, 0, \dots, 0, 1, 0, \dots, 0).$$

According to (15) all vectors

$$(L_0, \dots, L_i, \dots, L_j, \dots, L_n) + k(0, \dots, 0, -1, 0, \dots, 0, 1, 0, \dots, 0) \quad (67)$$

with $0 \leq k \leq L_j - L_i$ are also weights. As we know, for $i < j$, $L_i \geq L_j$. Suppose $L_i > L_j$. Then k in (67) can be equal to one and

$$\lambda = \Lambda + h^i - h^j, \quad i < j$$

is a weight. Hence the A_n -module W is not a Fock space if in its orthogonal signature $\Lambda = (L_0, L_1, \dots, L_n)$ there exists $L_i > L_j$ for $0 < i < j$.

It remains to consider the modules with

$$\Lambda = (L_0, L_1, \dots, L_n), \quad L_0 \geq L_n. \quad (68)$$

Suppose for $0 < i < j$

$$\lambda = \Lambda + h^j - h^i = (L_0, L_1, \dots, L_i, L_i - 1, L_{i+1}, \dots, L_j, L_j + 1, L_{j+1}, \dots, L_n)$$

is a weight. Then $\lambda' = (L_0, L_1, \dots, L_i - 1, L_{i+1}, \dots, L_j, L_j + 1, L_{j+1}, \dots, L_n)$ is also a weight. This is, however, impossible since $\lambda' > \lambda$. Hence all A_n -modules with signatures (68) are Fock spaces.

We could have stopped the proof here since the signatures

$$(L_0, L_1, \dots, L_n) \quad \text{and} \quad (L_0 - 1, L_1, \dots, L_n) \quad (69)$$

describe one and the same A_n -module. This could have been done if all information

was carried by A_n , i.e., if the dynamical variables were functions of the generators of A_n only. This is however not the case. The 4-momentum (42) $P^m \in A_n$ although $P^m \in \mathfrak{gl}(n+1)$. Therefore physically the representations (69) are distinguishable.

We shall determine the orthogonal coordinates of Λ from the requirement for the energy of the vacuum $|0\rangle = x_\Lambda$ to be zero. In terms of the orthogonal basis (28) P^m can be written as

$$P^m = \sum_{i=1}^n k_i^m h_i. \quad (70)$$

Since for $\Lambda = (L_0, L, \dots, L)$ $h x_\Lambda = L x_\Lambda$, $i = 1, \dots, n$ we require

$$P^m |0\rangle = \sum_{i=1}^n k_i^m L |0\rangle = 0 \quad m = 1, 2, 3. \quad (71)$$

Here k_1^0, \dots, k_n^0 are analogs of the energy spectrum of the one-particle states, $k_j^0 \neq 0$ (see (18)). Therefore (71) implies $L = 0$.

Later on we shall see that h_i , $i = 1, \dots, n$ is a number operator for particles in a state i . This together with (71) also gives $L = 0$.

Consider the Fock space W_p with $\Lambda = (p, 0, \dots, 0)$. Using the definition (41) of the a -operators from (62) we have

$$a_i^- a_j^+ |0\rangle = \delta_{ij} p |0\rangle. \quad (72)$$

We obtain the same expression as in the case of parastatistics of order $p^{/8/}$. Therefore we call p an order of the A -statistics. We conclude that like in the parastatistics all Fock spaces are labelled with positive integers p , the order of the statistics.

The equation (72) together with the commutation relations (44) of the a -operators determines completely the representation space and the representation of the creation and annihilation operators of order p . The A -statistics can be defined by the relations (44). The representations of the statistics can be obtained from (72). In this case all calculations can be done without using any Lie algebraical properties of the a -operators. Clearly this point of view is convenient for generalization to the case of infinite and in particular to continuum number of operators. The Lie algebraical structure however helps a lot in all calculations. Therefore we shall continue to consider a finite number of pairs a_1^\pm, \dots, a_n^\pm of a -operators and on a later stage we shall let $n \rightarrow \infty$.

Let us consider some Lie-algebraical properties of the Fock spaces. In the A_n -module W with a highest weight $\Lambda = (L_0, \dots, L_n)$ an arbitrary weight $\lambda = (\ell_0, \dots, \ell_n)$ can be represented as

$$\lambda = \Lambda + \sum_i k_i \omega_i, \quad k_i \in N_0,$$

where

$$\omega_i \in \Sigma^- = \{h^i - h^j \mid i > j = 0, 1, \dots, n\}.$$

Since the sum of the first m co-ordinates, $m = 1, 2, \dots, n$ of an arbitrary negative root $\omega_i \in \Sigma^-$ is non-positive this is true also for the vector $\sum k_i \omega_i$ with k_i non-negative integers. Therefore for an arbitrary weight λ we have

$$\ell_0 + \ell_1 + \dots + \ell_m \leq L_0 + L_1 + \dots + L_m \quad m = 0, 1, \dots, n.$$

From this inequality and the circumstance that the weight system is invariant under permutations of the orthogonal coordinates we conclude that the vector $\lambda = (\ell_0, \ell_1, \dots, \ell_n)$ with integer co-ordinates is a weight if and only if

$$\ell_{i_0} + \ell_{i_1} + \dots + \ell_{i_m} \leq L_0 + L_1 + \dots + L_m \quad (73)$$

where $i_0 \neq i_1 \neq \dots \neq i_m = 0, 1, \dots, n$; $m = 0, 1, \dots, n$. Clearly (73) is equally for $m = n$.

Lemma 3. All weights of the A_n -module of Fock W_p with order of the statistics P are simple.

Proof

An arbitrary weight vector $x_\lambda \in W_p$ with weight λ is generated from x_Λ with polynomials of the creation operators,

$$x_\lambda = P(a_1^+, \dots, a_n^+) x_\Lambda. \quad (74)$$

Therefore the weight $\lambda = (\ell_0, \ell_1, \dots, \ell_n)$ of x_λ can be represented as

$$\lambda = \Lambda + \sum_{i=1}^n k_i (-h^0 + h^i) \quad k_i \in N_0. \quad (75)$$

In terms of the co-ordinate the last relation reads as

$$(\ell_0, \ell_1, \dots, \ell_n) = (p, 0, \dots, 0) + (-\sum_{i=1}^n k_i, k_1, k_2, \dots, k_n). \quad (76)$$

Hence $k_i = \ell_i$, $i = 1, \dots, n$ and an arbitrary weight λ is represented uniquely in the form (75). In terms of the weight vectors this gives that $P(a_1^+, \dots, a_n^+)$ in (74) is homogeneous with respect to every creation

operator a_i^+ :

$$P(a_1^+, \dots, a_i^+, \dots, a_n^+) = a_i^{\ell_i} P(a_1^+, \dots, a_i^+, \dots, a_n^+).$$

Since the creation operators commute,

$$P(a_1^+, \dots, a_n^+) = (a_1^+)^{\ell_1} (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n}.$$

Therefore every vector x_λ with weight $\lambda = (\ell_0, \ell_1, \dots, \ell_n)$ is collinear to the vector

$$(a_1^+)^{\ell_1} (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} x$$

and the corresponding weight space is one-dimensional

This lemma has no analogy in the para-statistics. For instance the states $|b_i^+ b_j^+ \rangle_0$ and $|b_j^+ b_i^+ \rangle_0$, $i \neq j$ have one and the same weight but in general are linearly independent.

In the following lemma we prove one important property of the Λ -statistics.

Lemma 4. Given A_n -module of Fock W_p with order of the statistics p . The vector

$$(a_1^+)^{\ell_1} (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle \quad (76')$$

is non-zero if and only if

$$(\ell_1 + \ell_2 + \dots + \ell_n) \leq p. \quad (77)$$

In particular in the Fock space W_p there can be no more than p particles.

Proof

In the previous lemma we saw that the vector (76) has a weight

$$\lambda = (p - \ell_1 - \dots - \ell_n, \ell_1, \dots, \ell_n). \quad (78)$$

If $\ell_1 + \dots + \ell_n \leq p$, then clearly (73) holds because $L_0 + \dots + L_m = p, m = 0, 1, \dots, n$. Therefore λ is a weight. There should exist at least one weight vector with weight λ . Since the multiplicity of λ is one, this is the vector (76) and hence this vector is not zero.

If $\ell_1 + \dots + \ell_n > p$, the weight (78) does not fulfil the inequality (73) for $m = n - 1$ and $\ell_{i_0} = \ell_1, \ell_{i_1} = \ell_2, \dots, \ell_{i_{n-1}} = \ell_n$ and the corresponding weight vector (76) is zero.

From (76) and (78) we conclude

$$\begin{aligned} h_i (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle &= \\ = \ell_i (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle & \quad i = 1, \dots, n. \end{aligned} \quad (79)$$

The operator h_i is a number operator of the particles in the state i . The number operator N is

$$N = N_1 + N_2 + \dots + N_n. \quad (80)$$

We obtain

$$N (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle = (\ell_1 + \dots + \ell_n) (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle \quad (81)$$

4. MATRIX ELEMENTS OF THE CREATION AND ANNIHILATION OPERATORS

The numbers ℓ_1, \dots, ℓ_n together with the order of the A -statistics p determine uniquely the state (76). We introduce the notation

$$|p; \ell_1, \ell_2, \dots, \ell_n\rangle = (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle. \quad (82)$$

The set of all vectors (82) constitute a basis of weight vectors in the Fock space W_p . The correspondence between the weight vectors and the weights written with their orthogonal co-ordinates reads as

$$|p; \ell_1, \ell_2, \dots, \ell_n \rangle \leftrightarrow (p - \sum_{i=1}^n \ell_i, \ell_1, \ell_2, \dots, \ell_n). \quad (83)$$

One has to remember that the notation $|p; \ell_1, \dots, \ell_n \rangle$ is defined only for $\ell_1 + \dots + \ell_n \leq p$.

We now proceed to calculate the matrix elements of n pairs of creation and annihilation operators a_1^\pm, \dots, a_n^\pm in the A_n -module of Fock W_p with order of statistics p .

We can write immediately

$$h_0 |p; \ell_1, \dots, \ell_n \rangle = (p - \sum_{i=1}^n \ell_i) |p; \ell_1, \dots, \ell_n \rangle, \quad (84)$$

$$h_i |p; \ell_1, \dots, \ell_n \rangle = \ell_i |p; \ell_1, \dots, \ell_n \rangle, \quad i = 1, \dots, n.$$

These equalities follow from the observation that the orthogonal co-ordinates of the weight (83) are eigenvalues of the operators (28) on the weight vector (82). Since

$$[a_i^-, a_i^+] = h_0 - h_i$$

we have

$$[a_i^-, a_i^+] |p; \ell_1, \dots, \ell_n \rangle = (p - L - \ell_i) |p; \ell_1, \dots, \ell_n \rangle, \quad (85)$$

where $L = \ell_1 + \ell_2 + \dots + \ell_n$.

First we calculate the matrix element of a_1^- .

$$\begin{aligned}
a_1^- |p; \ell_1, \dots, \ell_n\rangle &= [a_1^-, (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n}] |0\rangle = \\
&= [a_1^-, (a_1^+)^{\ell_1}] (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle + (a_1^+)^{\ell_1} a_1^- (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle.
\end{aligned}
\tag{86}$$

The second term in the last equality vanishes. Indeed the vector

$$a_1^- (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle$$

would have had a weight

$$(p - \sum_{i=2}^n \ell_i + 1, -1, \ell_2, \ell_3, \dots, \ell_n)$$

which is impossible since $\ell_0 + \ell_2 + \ell_3 + \dots + \ell_n = p + 1 > p$.

Using (84), for the first term we obtain

$$\begin{aligned}
&\sum_{i=0}^{\ell_1-1} (a_1^+)^i [a_1^-, a_1^+] (a_1^+)^{\ell_1-i-1} (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle = \\
&= \sum_{i=0}^{\ell_1-1} (p - \ell - \ell_1 + 2i + 2) \cdot (a_1^+)^{\ell_1-i-1} (a_2^+)^{\ell_2} \dots (a_n^+)^{\ell_n} |0\rangle.
\end{aligned}$$

This gives

$$a_1^- |p; \ell_1, \dots, \ell_n\rangle = \ell_1 (p - \sum_{j=1}^n \ell_j + 1) |p; \ell_1 - 1, \ell_2, \dots, \ell_n\rangle.$$

The generalization for a_i^- is evident:

$$a_i^- |p; \ell_1, \dots, \ell_i, \dots, \ell_n\rangle = \ell_i (p - \sum_{k=1}^n \ell_k + 1) |p; \ell_1, \dots, \ell_{i-1}, \dots, \ell_n\rangle. \tag{87}$$

Moreover

$$a_i^+ |p; \ell_1, \dots, \ell_i, \dots, \ell_n\rangle = |p; \ell_1, \dots, \ell_i + 1, \dots, \ell_n\rangle. \tag{88}$$

The metric in W_p is defined in a complete analogy with the scalar product in the Fock space of Bose (or Fermi) particles.

Postulate

a) $\langle 0|0\rangle = 1$

b) $\langle 0|a_i^+ = 0, \quad i = 1, \dots, n$

c) $((a_1^+)^{m_1} \dots (a_n^+)^{m_n} |0\rangle, (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle) =$
 $= \langle 0| (a_1^-)^{m_1} \dots (a_n^-)^{m_n} (a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n} |0\rangle. \quad (89)$

The vectors $|p; \ell_1, \dots, \ell_n\rangle$ constitute an orthogonal basis in W_p . To show this, suppose that in (89) some $m_i \neq \ell_i$ and let $m_i > \ell_i$. Then the vector

$$(a_i^-)^{m_i} (a_1^+)^{\ell_1} \dots (a_i^+)^{\ell_i} \dots (a_n^+)^{\ell_n} |0\rangle = 0$$

since otherwise there has to exist a weight

$$(p - \sum_{j=1}^m \ell_j + m_i, \ell_1, \dots, \ell_{i-1}, -(m_i - \ell_i), \ell_{i+1}, \dots, \ell_n)$$

which is impossible. For $m_i < \ell_i$ the same result can be obtained from the hermitian conjugate of (89). If $m_i = \ell_i, i = 1, \dots, n$ we obtain

$$(|p; \ell_1, \dots, \ell_n\rangle, |p; \ell_1, \dots, \ell_n\rangle) = \frac{p!}{(p-L)!} \prod_{i=1}^n \ell_i!, \quad (90)$$

where $L = \ell_1 + \dots + \ell_n$.

As an orthogonal basis in W_p one can accept the vectors

$$|p; \ell_1, \dots, \ell_n\rangle = \sqrt{\frac{(p-L)!}{p!}} \frac{(a_1^+)^{\ell_1} \dots (a_n^+)^{\ell_n}}{\sqrt{\ell_1! \ell_2! \dots \ell_n!}} |0\rangle. \quad (91)$$

In this basis we have for the matrix elements

$$a_i^+ |p; \ell_1, \dots, \ell_n\rangle = \sqrt{(\ell_i + 1)(p - \sum_{j=1}^n \ell_j)} |p; \ell_1, \dots, \ell_i + 1, \dots, \ell_n\rangle \quad (92)$$

$$a_i^- |p; \ell_1, \dots, \ell_n\rangle = \sqrt{\ell_i (p - \sum_{j=1}^n \ell_j + 1)} |p; \ell_1, \dots, \ell_i - 1, \dots, \ell_n\rangle \quad (93)$$

The matrix elements of the a -operators do not depend on n . Therefore the results can be extended in an evident way to the case of infinite number of operators.

Finally, we point out one interesting property of the A -statistics. Introduce the operators

$$A_i^\pm = \frac{a_i^\pm}{\sqrt{p}}, \quad i = 1, \dots, n \quad (94)$$

and consider the matrix elements of these operators on states with number of particles much less than p ,

$$\ell_1 + \ell_2 + \dots + \ell_n \ll p. \quad (95)$$

From (92-93) we obtain

$$A_i^- |p; \ell_1, \dots, \ell_n\rangle \approx \sqrt{\ell_i} |p; \ell_1, \dots, \ell_i - 1, \dots, \ell_n\rangle \quad (96)$$

$$A_i^+ |p; \ell_1, \dots, \ell_n\rangle \approx \sqrt{\ell_i + 1} |p; \ell_1, \dots, \ell_i + 1, \dots, \ell_n\rangle.$$

In a first approximation

$$\begin{aligned} [A_i^+, A_j^+] &= [A_i^-, A_j^-] = 0 && \text{exact commutators} \\ [A_i^-, A_j^+] &\approx \delta_{ij} \quad \text{if } \ell_1 + \dots + \ell_n \ll p. \end{aligned} \quad (97)$$

Moreover if (95) holds then

$$|p; \ell_1, \dots, \ell_n\rangle \approx \frac{(A_1^+)^{\ell_1} \dots (A_n^+)^{\ell_n}}{\sqrt{\ell_1! \dots \ell_n!}} |0\rangle. \quad (98)$$

We see that if the A-statistics allows a large number of particles p , then the commutation relations of the operators A_i^\pm on states with $\ell_1 + \dots + \ell_n \ll p$ coincide in a first approximation with the Bose creation and annihilation operators. In the limit $p \rightarrow \infty$ the operators A_i^\pm reduce to Bose operators.

This property has also an interesting Lie-algebraical consequence. It shows that the limit of certain representations (the Fock representations) of the simple algebra A_n leads to a representation of the solvable Lie algebra of Bose operators.

We have considered the statistics corresponding to the algebra of the unimodular group. In a similar way one can introduce a concept of C- and D-statistics^{/9/} or of statistics that correspond to other semi-simple Lie lagebras^{/5/}.

REFERENCES.

1. Green H.S. Phys.Rev., 1953, 90, p.270.
2. Kamerfuchi S., Takahashi Y. Nucl. Phys., 1960, 36, p.177.
3. Ryan C., Sudarshan E.C.G. Nucl.Phys., 1963, 47, p.207.
4. Palev T. Ann.Inst.Henri Poincare, 1975, 23, p.49.
5. Palev T. JINR, E2-10258, Dubna, 1976.

6. Bogolubov N.N., Shirkov D.V. Introduction to the Theory of Quantized Fields. Moscow, 1957 (English ed. Interscience Publishers, Inc., New York, 1959).
7. Jacobson N. Lie algebras. Interscience Publ., New York, 1962.
8. Greenberg O.W., Messiah A.M. Phys.Rev., 1965, 138, p.1155.
9. Palev T. Thesis. Institute for Nuclear Research and Nuclear Energy, Sofia (1976).

Received by Publishing Department
on March 31, 1977.

Subject Categories of the JINR Publications

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Gryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids.
15.	Experimental physics of nuclear reactions at low energies.
16.	Health physics. Shieldings
17.	Theory of condensed matter.



Издательский отдел Объединенного института ядерных исследований.
Заказ 23035. Тираж 370. Уч.-изд. листов 2,04.
Редактор Э.В. Пвашкевич. Подписано к печати 22.04.77 г.
Корректор Р.Д. Фомина.