

**MASTER**

CONF-790676-3

To be published in  
F. J. deSerres, W. M. Generoso  
and M. D. Shelby (eds.)  
DNA Repair and Mutagenesis  
in Eukaryotes, Plenum Press,  
New York, 1979

GENETIC AND PHYSIOLOGICAL FACTORS AFFECTING REPAIR AND MUTAGENESIS  
IN YEAST\*

Jeffrey F. Lemontt  
Biology Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830

Running Title: REPAIR AND MUTAGENESIS IN YEAST

---

\*Research sponsored by the Office of Health and Environmental Research,  
U. S. Department of Energy, under contract W-7405-eng-26 with the Union  
Carbide Corporation.

By acceptance of this article, the  
publisher or recipient acknowledges  
the U.S. Government's right to  
retain a nonexclusive, royalty-free  
license in and to any copyright  
covering the article.

— NOTICE —  
This report was prepared as an account of work  
sponsored by the United States Government. Neither the  
United States nor the United States Department of  
Energy, nor any of their employees, nor any of their  
contractors, subcontractors, or their employees, makes  
any warranty, express or implied, or assumes any legal  
liability or responsibility for the accuracy, completeness  
or usefulness of any information, apparatus, product or  
process disclosed, or represents that its use would not  
infringe privately owned rights.

**DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED**

cep

Send proof to: Dr. J. F. Lemontt  
Biology Division  
Oak Ridge National Laboratory  
Post Office Box Y  
Oak Ridge, Tennessee 37830

Abbreviations: UV, ultraviolet light; NA, nitrous acid; MMS, methylmethanesulfonate; EMS, ethylmethanesulfonate; MNNG, N-methyl-N'-nitro-N-nitrosoguanidine; HZ, hydrazine

## SUMMARY

Current views of DNA repair and mutagenesis in the yeast Saccharomyces cerevisiae are discussed in the light of recent data, and with emphasis on the isolation and characterization of genetically well-defined mutations that affect DNA metabolism in general (including replication and recombination). Various "pathways" of repair are described particularly in relation to their involvement in mutagenic mechanisms. In addition to genetic control, certain physiological factors such as "cell age", DNA replication and the regulatory state of the mating-type locus, are shown to also play a role in repair and mutagenesis.

## INTRODUCTION

Since the realization that DNA repair and mutagenesis in E. coli are closely related, enzymatically controlled cellular processes [8, 59, 172, 175], there has been a great deal of interest to determine whether repair mechanisms proposed for prokaryotes might also apply to eukaryotic cells. The axiomatic belief held by many that these more highly evolved cells would prove to be more complex in every way no doubt influenced many investigators to focus attention upon the unicellular uninucleate budding yeast Saccharomyces cerevisiae. This organism is among the simplest of all eukaryotes and offers several advantages over not only simpler prokaryotic systems but also more complex eukaryotic systems. These include: 1) ease of handling and low cost, 2) sophisticated genetic manipulation, 3) well-defined mitotic cell cycle in stable haploid, diploid and polyploid strains, 4) a relatively high tolerance of certain aneuploidies, 5) conjugation and meiotic development, 6) a nucleosome-dependent chromosomal organization, 7) a wealth of genetic and biochemical information concerning both nuclear-specific and mitochondrial-specific cellular processes [3, 5, 20, 51, 54, 118, 124, 149, 150, 154].

Attempts to explain the phenotypic traits of uvr, lex, rec and other mutants of E. coli, most notable of which is enhanced mutagen cytotoxicity, led inescapably to the hypothesis that this increased lethality is caused by the failure to repair DNA damage, providing the genomes suffer the same number of initial lesions [68, 172]. Moreover, the proposal by Evelyn Witkin in 1967 [173] that different pathways of repair could

represent either an accurate (error-proof or error-free) or an inaccurate (error-prone or mutagenic) molecular restoration of genetic information proved to be an important turning point in the way geneticists interpreted experiments concerning repair effects on induced mutagenesis. Today, error-prone repair of UV damage in bacteria is believed to have an inducible component involving alteration or inhibition of the editing (nucleolytic) function of a DNA polymerase [78, 176].

In spite of efforts to extend the concept of error-prone repair to eukaryotic organisms, we still know very little about the nature of this repair, much less how the errors are made. Thus, without direct error frequency assays involving defined damaged-DNA substrates and purified repair enzymes, it should be emphasized that the error-prone repair concept represents only a descriptive albeit useful construct, just as it did in 1967. The evidence for error-prone repair depends upon the identification of one or more mutant genes conferring both increased mutagen sensitivity and defective induced mutability, compared to a wild type. Similarly, error-free repair is inferred by virtue of its apparent absence in well-defined hypersensitive mutants that are also hypermutable (presumably because more lesions are repaired by error-prone repair). Obviously, these concepts cannot apply when chemical mutagens are subject to cellular metabolism unless numbers of DNA lesions are quantitated in both mutant and wild type cells. This is one reason why physical agents, particularly ultraviolet light and to a lesser extent ionizing radiations, have been so useful for quantitative studies of repair and mutagenesis.

There now exists a considerable body of information on repair and mutagenesis in Saccharomyces cerevisiae. Although certain aspects of this subject have been reviewed in the past [19, 56, 121, 164], the present summary will attempt to broaden our current understanding not only by including more recent genetic data, but also by emphasizing the interrelation between cellular processes that involve DNA metabolism, namely repair, mutagenesis, recombination and replication; also included are developmental systems regulating the sexual cycle which affect chromatin indirectly.

#### ISOLATION OF MUTANTS WITH ENHANCED MUTAGEN SENSITIVITY

In 1967 Nakai and Matsumoto [122] were the first to describe radiosensitive mutants in yeast. They identified two separate mutant loci, now called rad1 and rad2 [38], that caused a significant loss of UV resistance; another, now called rad51 [37, 42], led primarily to X-ray sensitivity with only a slight effect on UV sensitivity. Snow [157] then reported 6 UV-sensitive mutants, each carrying different mutant loci, four of which were also hypersensitive to NA (but with a different rank order). Snow hypothesized that although repair of lesions inflicted by UV and NA would likely occur by nonidentical enzymatic reactions, the different repair enzymes might represent components of one general repair system. Cox and Parry [22] then deliberately attempted to "saturate" the yeast genome with mutations conferring UV sensitivity in an effort to estimate the total number of independent genes responsible. Genetic analysis of 96 isolates revealed 22 separate mutant loci, 5 of which were also

(T-1) responsible for increased X-ray sensitivity. Subsequent isolations (Table 1) have revealed a large number of genetic loci controlling mutagen sensitivity, including 22 new genes detected in mutants selected for MMS sensitivity by Prakash and Prakash [135]. Of these, only 5 conferred MMS sensitivity alone, while others caused either UV (six) or X-ray (five) sensitivity as well, and 6 others led to cross sensitivity to both radiations.

Ananthaswamy *et al.* [1] recently attempted to "saturate" the yeast genome with mutations controlling X-ray sensitivity and found 15 new *complementation groups*, each complementing the existing rad set. Among such X-ray sensitive mutants all but four were cross sensitive to MMS, while none were cross sensitive to UV. Although allelism tests to mms strains have not been performed, it is likely that several new unique genes are represented because the combined X-ray-and MMS-sensitivity phenotype of these 11 is expressed by only 5 of 22 mms loci.

To summarize thus far, UV-sensitive mutants comprise at least 19 genetic loci (rad1,2,3,4,7,10,13,14,16,19,20,21,22 and mms3,6,10,13,18,19), X-ray-sensitive mutants 26-31 loci (rad50,51,52,53,54,55,56,57, xs3, xrs2, xrs4, mms8,9,14,16,20, and an estimated 10-15 others), UV- and X-ray sensitive mutants 15 loci (rad5,6,8,9,11,12,15,17,18 and mms7,11,12,15,17,21), and finally mutants at 5 loci (mms1,2,4,5,22) sensitive to MMS but not to radiations. Since many but not all of the above 65-70 mutations have been shown to recombine with one another in meiosis, it is important to bear in mind the distinction between allelism and genetic complementation in estimating numbers of genetic loci. On the other hand, complementing repair-deficient mutants at the same chromosomal locus have not been reported.

Does this very large number of genes represent a reasonably accurate estimate for all possible mutants of this type? Although the answer to this question is not known, it does not seem likely to be a resounding "yes" for several reasons. On statistical grounds (based on the distribution of allelic repeats observed at 22 loci), Cox and Parry [22] estimated an additional 8-15 undetected mutable rad loci. With respect to the 22 new mms loci, Prakash and Prakash [135] have calculated a maximum likelihood estimate of  $48 \pm 15$  (standard deviation) loci responsible for MMS sensitivity. Table I shows that mutants at some loci (e.g. rad1 or rad2) have been more readily detectable than others. Doubtless, this results from a variety of factors including the original strain employed (genetic background), the dose and type of mutagen used to enrich the mutant population prior to screening, the dose and type of mutagen used for screening hypersensitive strains, the conditions of treatment, and finally the level of effect arbitrarily chosen as a criterion for isolation. For example, the extreme UV sensitivity of rad1 and rad2 strains, a result of defective pyrimidine dimer excision repair [130, 162, 166] is probably a major reason such mutants have been repeatedly reisolated. Moreover, Prakash and Prakash [135] screened for the inability to grow in the continual presence of 0.5% MMS and found three loci (mms2,10,22) that did not confer MMS hypersensitivity when cells were exposed to brief MMS treatments in buffer. Thus, the permeability of MMS in such strains may be different under different conditions and in different genetic backgrounds. Since mms10 also enhances UV sensitivity, repair processes may be involved. Since mms2 and mms22 confer only enhanced MMS sensitivity (under certain conditions), MMS-specific repair processes may exist or alternatively, genetically altered MMS transport into the cell nucleus may be involved, again underscoring the need to compare strains having the same initial DNA damage.

## ISOLATION OF MUTANTS AFFECTED IN MUTAGENESIS AND RECOMBINATION

Early recognition of the interrelation between DNA repair, mutagenesis, recombination and replication in bacteria [68, 174] stimulated a number of yeast geneticists to ask whether these processes have anything in common in eukaryotic organisms. In particular, 1) Are mutagenesis and recombination in yeast genetically controlled?, 2) If so, how many genes are involved?, and 3) Do any of these genes function in repair-associated mechanisms? Table 2 shows that a number of genes have been identified in mutants selected for a variety of genetic endpoints other than enhanced mutagen sensitivity. These include spontaneous and induced mutation as well as spontaneous and induced mitotic recombination.

### Ultraviolet reversion-defective mutants

Using the vigorous UV-induced revertibility of the arg4-17 ochre allele to monitor induced mutability, Lemontt [93] screened clones (derived from cells surviving EMS treatment) for defective UV reversion. Upon genetic analysis 20 such isolates were found to comprise single recessive alleles of only 3 genes, called rev1, rev2 and rev3; rev2 was subsequently found to be allelic with UV-sensitive mutants isolated by Snow [157] and by Cox and Parry [22], and has since been renamed rad5 [38]. Mutations at any one of these 3 loci cause varying degrees of enhanced sensitivity to UV, X-rays and EMS, implicating their involvement in some form of DNA repair. This suggested that UV mutagenesis in yeast is genetically controlled by an error-prone repair process, as already proposed for E. coli at that time by Witkin [175].

T-2

The rev1 and rev3 genes were shown to cause large reductions in UV mutation frequencies compared to the wild type not only for reversion of arg4-17 (ochre), lys1-1 (ochre) and arg4-6 (putative missense) [93], but also for forward mutation at biosynthetic loci across the genome leading to auxotrophic requirements, and for forward mutation at 2 specific ADE loci (ade1 or ade2) [148] causing red-pigmented clones [95]; rev2 on the other hand, had much smaller effects at arg4-17 and lys1-1, and at biosynthetic loci. Yet rev2 had no significant effect at all on UV reversion of arg4-6 or on forward mutation at the ADE loci [93, 95]. Moreover, the average effect of rev1 across the genome (4% of the wild-type response) was much greater than at the 2 selected ade genes (19% of wild type), while the effect of rev3 was large in both cases (4% and 2% of wild type, respectively) [95]. This was one of the first indications that UV mutagenesis might not be acting uniformly at all genetic sites, i.e. a hot-spotting effect or a specificity of interaction between certain mutagenic (error-prone repair) enzymes and particular genomic regions or particular types of DNA damage. It was also suggested that the rev2 block is highly specific, perhaps affecting only UV reversion of ochre alleles.

More recently, extensive data of Lawrence and Christensen concerning the effect of rev genes on UV reversions of well-defined cyc1 (iso-1-cytochrome c) alleles has for the most part confirmed these earlier suggestions of specificity and nonrandomness of UV mutagenesis [88, 89, 91]. In addition, they have identified several other mutant rad loci that reduce UV

mutagenesis [87] -- rad6, rad8, rad9 and rad18. These all cause enhanced sensitivity to both UV and X-rays, like the rev genes and like recA and lex genes of E. coli [175]. Unlike lexA, however, which is dominant over the wild type allele [119], rad and rev genes involved in UV mutagenesis are all recessive in their effects on survival and induced mutation, suggesting the loss of required enzymatic steps in the mutagenic mechanism.

#### Forward mutation at the CAN1 locus

Unlike reversion, recessive forward mutation usually results in the loss of an essential cellular function and, in principle, can derive from a variety of mutational alterations. Many systems used to quantitate forward mutation (e.g. auxotrophy, pigmented clones, lethals) have limited utility mainly because they are nonselective and therefore relatively inconvenient. Many systems measure mutation at any one of a large number of genetic loci. Forward mutation at CAN1, however, represents a convenient, selective drug-resistance system that is sensitive to a variety of physical and chemical mutagens [11, 45, 49, 83, 84, 99, 100, 139, 168].

Recessive can1 mutants become resistant to the highly toxic arginine analogue, canavanine, by mutational alteration of CAN1 on chromosome V, believed to be the structural gene for the arginine-specific permease enzyme [49, 168]. Since this permease transports virtually all exogenous arginine (and canavanine) into the cell under normal conditions of ammonia

repression when general amino-acid permeases are inactive, all such canavanine-resistant mutants map at this one genetic locus [47, 48, 49]. Intragenic (interallelic) complementation has not been observed even among a large number of unique dihybrids, suggesting that arginine permease is functional as a single polypeptide [168]. Fine-structure mapping of alleles yielding the greatest recombination is suggestive of enough DNA to code for a protein as large as 260,000 daltons [168]. The molecular weight of arginine permease is not known, but it could be considerably less if a) the correspondence between gene and protein for this mapping method [81, 110, 123] is unreliable for CAN1 (the largest gene presumed to exist in yeast), as appears to be the case for very small genome intervals [115, 116], or if b) certain portions of the gene are non-structural and are subject to post-transcriptional or post-translational processing critical for functional integration of the permease into the cell membrane. Thus, mutations in noncoding but critical sequences could also result in inactive permease; and, unlike most mutable genes used in mutation studies, which usually affect soluble enzyme activities, CAN1 is responsible for the activity of an important membrane protein that must be synthesized (presumably on cytoplasmic ribosomes) and subsequently transported and integrated into the cell envelope in some specific way.

In wild-type yeast grown to stationary phase in a yeast extract-peptone-dextrose (YEPD) complex broth, it is observed that many mutagens including UV cause vigorous induction of can1 mutants and these are readily expressed on selective agar containing the drug, presumably before canavanine toxicity becomes too great [45, 99, 100, 101]. This is believed to be due in part to a relatively high turnover rate of the permease such that mutational

expression is strongly influenced by the cellular level of endogenous free arginine (dependent upon type of growth medium), rather than by the ability to undergo residual divisions on the plate [45]. This is consistent with the general observation that canavanine cytotoxicity is dependent upon the exogenous ratio of canavanine to arginine such that defective canavanyl proteins are eventually synthesized. Thus, for pre-growth in YEPD broth media the free arginine pool is presumably high enough to prevent significant toxicity during a period when the permease activity is decaying rapidly.

Ultraviolet forward mutation-defective mutants

In an effort to identify new genes controlling UV mutagenesis or its expression at CAN1, Lemontt screened for clones (YEPD pre-growth) with less than wild-type levels of UV mutation to canavanine resistance [96]. Such ultraviolet mutation-resistant isolates were subsequently characterized and found to carry one of 7 unlinked recessive umr alleles [100]. The umr loci did not cause canavanine resistance and were unlinked to can1, nor could they be explained by an extra (disomic) copy of chromosome V. Unlike rev or rad mutants, four of these genes (umr4, umr5, umr6, umr7) had no significant effect either on UV sensitivity or on the UV revertibility of 3 ochre mutations, his5-2, lys1-1, and ura4-1. Diploids homozygous for umr5, umr6 or umr7 all failed to sporulate, suggesting a meiotic defect [100]. The umr7 locus, known to be allelic with and mapping in the same region as tup1 and cyc9 on chromosome III, has an exceedingly rich pleiotropic phenotype with effects on conjugation ( $\alpha$ -specific poor mating ability), the cell surface (extreme flocculence or cell clumping and "self-shmooing"),

and membrane-associated functions (dTMP uptake and unusually high levels of iso-2-cytochrome c) [100, 104, 105, 151, 169]. The UMR4, UMR5, UMR6 and UMR7 genes may be more concerned with expression of can1 mutations rather than with mechanisms of mutagenesis directly [101]. On the other hand, umr1, umr2 and umr3 mutants were slightly more UV-sensitive than the wild type and were affected to varying degrees in UV revertibility of one or another of the 3 ochre alleles [100]. This is consistent with the idea that one or more of these latter 3 UMR genes are concerned with highly specific branches of mutagenic pathways, those contributing very little to the overall repair potential of the cell. Homozygotes of umr2 and umr3 failed to sporulate. However, since all umr loci except umr1 also led to increased canavanine toxicity, and since UV-induced can1 mutation frequencies may be boosted if selection is delayed and preceded by a period of cell division in growth medium [101], it is possible that there has been a genetic alteration either in the arginine pool size or in the rate or quality of arginine permease turnover processes. Thus, UMR1 seems most likely to be involved in mutagenic repair pathways and this is further supported by the finding that umr1 rad2 and umr1 rad6 double mutant haploids are much more UV-sensitive (synergistic) than the respective single rad strains [103]. On the other hand, the involvement of UMR2 and UMR3 in mutagenic pathways remains more tentative.

"Hypo-rec" and "hyper-rec" mutants

In 1964 Holliday [62] proposed a molecular model for gene conversion in fungi implicating a role for repair enzymes such as nucleases, polymerases and ligases. The model involves breakage and reannealing of complementary DNA strands of homologous chromatids to generate a "hybrid"

region. If the region includes a heterozygous mutation, this hybrid DNA will contain one or more mispaired bases (mismatched or heteroduplex DNA), a substrate for repair. Recombination-deficient mutants of *E. coli* were found to be radiation sensitive [18]. Radiation-sensitive mutants of *Ustilago* [63] were recombination-deficient [64]. Further, postreplication repair of daughter-strand gaps in DNA [152] appeared to require recA<sup>+</sup>-dependent recombination ability [156]. All these findings had the effect of intensifying the search for genes affecting recombination in yeast. It should be emphasized that recessive rec mutants are not easily selected by conventional genetic means since single mutational events in one chromosome would not be expressed in heterozygous diploids. Instead, radiation-sensitive mutants were routinely examined (in homozygous condition) for effects on either meiotic or mitotic recombination. Early indications were that many rad genes had no effect in meiosis, whereas rad/rad mitotic cells generally expressed higher levels of radiation-induced recombination than comparable RAD/RAD or RAD/rad diploids at equal exposures, suggesting that unrepaired radiation damage to DNA is recombinogenic [94, 158].

In a more direct approach Rodarte-Ramon and Mortimer [146, 147] selected rec mutants directly on the basis of defective X-ray-induced mitotic gene conversion at arg4. They constructed a strain disomic (n+1) and heteroallelic for arg4 on chromosome VIII, thereby permitting expression of rec genes on any of the other haploid chromosomes. Seven genes were identified, two of which conferred X-ray sensitivity, one of which caused UV and X-ray sensitivity, and four of which had no effect on radiation survival; rec2 (X-ray sensitive) was later found to be allelic with rad52 [42]. These mutants suggested that enzymatic steps required for induced

recombination in yeast might also be shared by certain repair pathways.

With a similar system Maloney and Fogel [108] screened heteroallelic arg4 disomics for enhanced spontaneous mitotic gene conversion. Several genes have been identified and are believed to be affected in a regulatory mechanism that normally keeps mitotic recombination at a low level. (There exists genetic evidence for such repression of mitotic intragenic recombination [33]). Some of these mutants exhibit enhanced sensitivity to MMS, UV or X-rays, implicating a role of DNA repair in the regulation of spontaneous mitotic recombination. This is supported by the results of Prakash and Prakash [136] who found that homozygotes of mms8, mms9, mms13 or mms21 exhibit a hyper-rec phenotype (increased spontaneous mitotic segregation from CAN1/can1 to can1/can1); rad18 also shows the hyper-rec phenotype [7].

As summarized in a review by Resnick [142], it should be emphasized that mitotic recombination of alleles within a gene is observed to be predominantly a nonreciprocal process (i.e. gene conversion), rather than a result of reciprocal crossing-over events, whereas the reverse is generally true of intergenic recombination, particularly after the frequency has been raised by exposure to external agents [122a]; and, most mutagens are good recombinogens. In either case, mutational alteration of one homolog occurs much too infrequently to account for mitotic recombination in yeast and other fungi. The implications for mammalian somatic cell mutagenesis should be clear: The induction of autosomal recessive "mutants" in cell lines believed to be already heterozygous [16] may in fact be induction of mitotic crossing-over anywhere between the genetic locus and its centromere, or to a lesser extent, gene conversion (assuming that chromosome loss, deletion and nondisjunction can be excluded).

Probing the potential relation between DNA repair and spontaneous mutability, von Borstel and his coworkers found that the spontaneous mutation rate is increased by several rad genes -- rad18, rad52, xs3 [163] and more recently rad3, rad6, rad51 [55]. In screening directly for such mutators [165], a minimum of 8 genetic loci have been identified [55]; one, MUT6, is dominant and without effect on UV, X-ray or MMS sensitivity; among the other 7 recessive mut loci, all but one (mut1) sensitizes cells to one or more of these mutagens; mut5 has been reported to be allelic with rad 51. These authors believe that spontaneous DNA lesions (including replication errors) are susceptible to repair by systems having several steps in common with systems that repair mutagen-induced DNA damage. The isolation of antimutator strains has also been reported [138]. In addition, rev3 has antimutator activity [131].

Previous work with bacteriophage T4 mutator [159] and antimutator [28] strains has suggested that the exonucleolytic activity associated with the polymerase has a proof-reading or editing function that normally corrects (repairs) spontaneous replicative errors, presumably mismatched bases [58, 153]. The inducible component of UV-induced mutagenesis in E. coli also appears to be associated with some process that permits the replicase (DNA polymerase III) to make errors at higher than normal frequency [9]. Whether yeast and higher eukaryotic cells have the same or a similar mechanism is not known. It does seem clear, however, that spontaneous mutability is genetically controlled in a complex way that is not entirely independent of repair mechanisms.

There is evidence that spontaneous mutagenesis and mitotic recombination are under joint genetic control in yeast. Using a procedure to select

dominant or recessive mutations affecting spontaneous forward mutability at CAN1, Golin and Esposito [46] have described a semi-dominant mutation, rem1-1, that elevates the spontaneous rates of both mutagenesis and mitotic recombination. Meiotic recombination is not affected by this mutation even when homozygous, but ascospore viability is reduced, suggesting a meiotic defect in chromosomal integrity or disjunction. These authors feel that spontaneous mutation and recombination are enhanced as a result of an increase in specific DNA structures, such as mismatched base pairs or single-stranded regions. A previously selected meiotic mutation spo7-1 isolated as sporulation deficient, has been shown to be responsible for both antimutator activity (mitotic) and defective premeiotic DNA synthesis [31]. In this regard, it should be added that there is now a good correlation between meiotic deficiencies and certain X-ray-sensitive rad mutants [42, 44]; sporulative ability is reduced in homozygotes of rad51 and rad55; rad50, rad52 and rad57 homozygotes do sporulate but nearly all meiotic products are inviable, analogous to mei mutants of Drosophila [4] and rec mutants of Ustilago [66, 67]. Sporulation is completely abolished by rad6-1. Recent results by Game et al [44] show that RAD50, RAD52 and RAD57 are not required for early and late meiotic events (namely pre-meiotic DNA synthesis and sporulation, respectively), but are required for successful meiotic recombination.

To summarize there are now a large number of genes in yeast believed to control various aspects of DNA repair, mutagenesis and recombination. Mutants selected on the basis of one altered property often turn out to be pleiotropic with respect to another phenotypic trait. To this extent

certain yeast mutants appear analogous to bacterial mutants affected in some aspect of DNA metabolism. Clearly, not all of these genes may be concerned with DNA repair directly, as discussed by previous authors [27, 56, 135]. The challenge to define in molecular terms cellular functions gone awry in nearly 100 (and potentially more) mutants underscores and provides evidence for the enormous complexity of eukaryotic DNA-related metabolism. It should also be pointed out that the existence of dominant and semi-dominant mutations that jointly affect repair, mutagenesis and recombination in yeast raises the possibility that induction of such mutations in somatic tissues of mammals, for example, might also serve to increase (by recombination) the overall rate of homozygosis of deleterious heterozygous recessive loci. In this way, induction of hyper-rec mutants might increase the cancer risk.

#### "PATHWAYS" OF REPAIR

It has been possible to characterize presumed repair-deficient mutations by their interactions in multiply-mutant haploid strains, as developed by Game and Cox [39, 40] and by Haynes [56]. Howard-Flanders et al [69] were among the first to demonstrate its usefulness for understanding DNA repair mechanisms by constructing a double mutant of E. coli carrying both uvrA and recA and showing that the two mutations interacted synergistically, with regard to UV sensitivity. That is, UV survival of this double mutant was very much less than what would have been expected on the basis of an additive effect of the two single mutants; further, the UV dose yielding an average of one lethal event (37% survival) corresponded to approximately one pyrimidine dimer per cell. This result

suggested two important hypothesis: first, that uvrA and recA each block very different repair pathways acting on UV-damaged DNA, and second, that these two major pathways could account for virtually all of the UV resistance exhibited by the wild type. This agreed with the finding that uvrA mutants lacked excision repair but recA mutants did not [68]. Two or more mutations blocking DNA repair along the same linear pathway are expected to interact epistatically, such that the multiple mutant is no more sensitive to the mutagen than the most sensitive single mutant.

Despite some inherent limitations of multiple mutant analysis, as discussed in detail elsewhere [39, 40, 87], it has been possible to gain a certain amount of information concerning mechanisms of repair in yeast from this kind of approach in conjunction with other phenotypic traits expressed by repair-deficient mutants. On this basis, there exist 3 so-called "epistasis groups" of rad loci, such that a strain carrying multiple mutations within a group exhibits epistasis, while a strain with mutations in different groups exhibits either an additive or synergistic interaction. The epistasis groups are suggestive of metabolic pathways [21].

#### Excision repair of UV damage

The epistasis group defined by rad1, rad2, rad3, rad4, rad10 and rad16 consists of mutants with a biochemical defect in excision repair of UV-induced pyrimidine dimers [39, 130, 133, 134, 144, 145, 162, 166]; furthermore, rad22 is epistatic to rad1 with respect to UV survival [87]. Thus, at least 7 genes appear to be required for excision repair in yeast. It should be pointed out that excision repair acts only on nuclear DNA and is not able to remove pyrimidine dimers from mitochondrial DNA [130, 167]. In general, none of the mutants in this pathway leads to X-ray

sensitivity, nor do they have any effect on recombination, meiosis, or sporulation. Additionally, like uvr mutants of E. coli [175], most if not all of these mutants exhibit enhanced frequencies of UV mutagenesis compared to the wild type at equal UV doses and a significant fraction of the induced mutability is photoreversible in both mutant and wild-type strains [2, 87, 92, 141, 177]. At equal survival levels induced mutabilities are approximately the same in RAD and rad2 strains [30]. Moreover, with respect to the observed spectrum of base-pair changes inferred from amino-acid replacements in iso-1-cytochrome c among UV revertants of ochre cyc1 alleles [92], the rad1 response is the same as that produced by the wild-type. All these findings have suggested that UV mutations are produced predominantly from unexcised pyrimidine dimers in DNA by a mutagenic process different from the excision repair pathway, which is considered to be essentially error-free.

#### Error-prone repair of UV damage

The epistasis group defined by rad6, rad8, rad9, rad18, rev1, rev2 (i.e. rad5), and rev3, consists of mutants with varying degrees of both UV and X-ray sensitivity [21, 22, 40, 87, 93]; none except rad9 significantly reduce mitotic recombination [82]; rad6 prevents sporulation [22] while the others do not apparently affect meiosis. The rad18 gene is synergistic with rad1, rad2, or rad3, but epistatic with rad6 [40]. All 7 mutants of this group are epistatic with rad6, suggesting that all are involved in one major pathway concerned with repair of UV damage [87]. Mutants carrying rad6 or rad9 are proficient in carrying out pyrimidine dimer excision [134]; the others are also excision-proficient (see [19]),

The most interesting property of mutants in this group is defective UV mutagenesis, suggesting that this single repair pathway is error-prone for UV damage, by analogy to recA and lexA mutants of E. coli [87, 96]. From this point on, the analogy to prokaryotic mechanisms of mutagenesis begins to break down. Lawrence and Christensen [88, 90] have pointed out how many of the observations in yeast are at best difficult to explain with the one-step unitary model proposed for E. coli [12, 176], according to which suppression of the editing function of DNA polymerase permits replication past a pyrimidine dimer while inserting two random, often incorrect bases opposite the lesion. Thus, recA and lexA mutations should prevent induction of mutations of all types and at all genetic sites; and, mutations induced in the wild type should involve double base-pair changes [88, 90].

Lawrence et al. [92] have demonstrated that rad6 and rad18 both affect UV mutagenesis not only quantitatively but also qualitatively by altering the spectrum of base changes observed among induced revertants of cyc1 nonsense alleles. Not all mutants defining error-prone repair in yeast block UV-induced mutational changes of all types and at all genetic sites; and, double base-pair changes are rare in yeast [90]. While rev3 and rad6 are nonspecific and prevent normal levels of UV mutagenesis at every genetic site tested, the remaining mutants of this pathway have strong allele-specific effects with respect to UV reversion (for examples see Table 3).

The REV2 gene product appears to be concerned only with UV reversion of ochre alleles, yet, clearly, this is not the case for all such alleles. While the REV1 product may not be required for frameshift mutagenesis by

T-3

UV, it is required for many but not all base-pair transitions and transversions. By inspection of the nearly complete base sequence information in the region of many cycl alleles and of their revertants, it has been possible to test the idea that the allele specificity of UV reversion (as typified by rev1) may be based upon one or more of the following factors: a) position within the gene, b) kind of DNA triplet altered, c) type of base-pair change (e.g. transition vs. transversion, or AT to GC vs. GC to AT), d) variable recovery of reversions, e) nonrandom formation of dimers in regions rich in adjacent pyrimidines, f) unusual kinds of premutational lesions, or g) different ratios of mutagenic to nonmutagenic repair at different genetic sites. Lawrence and Christensen have concluded that none of these factors alone can satisfactorily account for the nonrandomness of UV reversion [88, 90]. Even in a wild-type strain the same ochre triplet occurring at different sites reverts with entirely different patterns of base-pair change, suggestive of some form of site (or sequence) specificity [155]. Moreover, UV reversion of the cycl-131 allele by GC to AT transition, which does not require REV1 but which is nonetheless photoreversible, occurs within an alternating purine-pyrimidine nucleotide sequence, obviously a region where intrastrand pyrimidine dimers cannot be induced [88, 90]. Thus, we also need to understand how DNA damage at one site results in mutation at another.

Thus far, the best explanation for site-specific mutagenesis is a presumed nonrandom interaction between certain gene products (of error-prone repair) and DNA damage in particular genomic regions [90]. It is not known whether this surprising level of complexity is unique to yeast or to eukaryotes in general. Bacterial studies in the past have for the most part not been concerned with this question of specificity. Not until

very recently have mutation-resistant ("rev-like") mutants of E. coli been selected directly [74]. If such apparent site-specific regulation of mutagenesis is found to be unique to eukaryotes, the molecular environment of the chromatin is likely to play a role.

#### Minor repair pathway for UV damage

A third "minor pathway" for repair of UV damage appears to involve enzymatic steps whose major ostensible function is to repair ionizing radiation-damaged DNA. The rad50 and rad51 genes each confer slight UV sensitivity and are epistatic to one another, yet rad51 interacts synergistically with both rad3 and rad18 with respect to UV survival. One lethal event (37% survival) in rad3 rad18 double mutants corresponds to approximately 6 pyrimidine dimers per cell, while in rad3, rad18 rad51 triple mutants, there are only 1 or 2 needed to produce the same effect. This suggests that unrepaired dimers are lethal and that virtually all of the UV resistance expressed by the wild type can be accounted for by the action of these 3 pathways [21, 40]. The rad52 locus also acts in this minor pathway, but has little or no effect on UV reversion. In excision-defective strains, however, a rad52 rad1 strain is nearly 10-fold more UV hypermutable than a rad1 strain, suggesting that this minor pathway is essentially error-free for repair of UV damage [87].

#### Repair of ionizing radiation damage

The major pathway for repair of ionizing radiation damage is controlled by RAD50, RAD51, ..., RAD57; rad52 contributes the greatest gamma-ray hypersensitivity, which is also exhibited by all double mutants with rad52 [113]. Frequencies of gamma-ray reversion in mutants of this group are

similar to that expressed by the wild type, suggesting an error-free mode of repair [112]. Since rad52 strains are defective in gamma-ray-induced mitotic gene conversion [146, 147] and are also unable to repair double-strand DNA breaks [60, 143], this RAD52 pathway may involve recombinational repair; rad52 strains also have increased X-ray-induced dominant lethality, suggestive of a defect in the repair of chromosome breaks [61].

With respect to ionizing radiation survival and mutagenesis, McKee and Lawrence [112, 113] have found that the single mutagenic repair system for UV is also responsible for mutagenic repair of ionizing radiation damage and requires the functions of the RAD6, RAD8, REV1, REV2 and REV3 genes. Although both radiations produce very different kinds of premutational DNA damage, gamma-ray mutagenesis is efficiently blocked by mutations of these loci, all of which comprise a "rad6 epistasis group" for gamma ray survival; rad9 and rad18 also belong to this group but do not block gamma-ray mutagenesis significantly, suggesting that this RAD6 pathway consists of both error-free and error-prone repair processes. In addition, McKee and Lawrence [114] have observed in rev strains allele-specific gamma-ray reversion patterns that are very similar to those expressed after UV exposure. These authors have argued that the simple idea of an enzymatic pathway for mutagenic repair consisting of sequential gene-controlled steps, with separate branch points leading to mutational specificity, does not adequately explain the distinctive yet partially overlapping mutational phenotypes expressed by mutants of this "pathway".

Mutagenesis by several chemical agents also requires a functional repair system, specifically the RAD6 and RAD9 gene products [129, 131, 132].

McKee and Lawrence [114] argue that mutations of different kinds or at different sites that arise from potentially very different premutational lesions are produced by the coordinate action of a large number of partially independent sets of gene functions.

#### PHYSIOLOGICAL FACTORS AFFECTING REPAIR AND MUTAGENESIS

Early studies on the recovery of yeast from radiation and chemical damage showed that the degree of liquid-holding recovery, an indicator of repair activity, could be modified by different physiological conditions [127, 128]. More recent studies [126] not only have emphasized the importance of genetically controlled repair processes but also have expanded our view of the diversity of cellular factors that can affect repair. These include "cell age", DNA replication and the mating-type dependent regulatory system.

##### "Cell age"

The term "cell age", as developed by Parry and co-workers [125] encompasses two different phenomena -- either the position in the mitotic cell cycle of synchronous cultures, or the transition of exponentially growing (log-phase) asynchronous cultures to a nutrient-limited stationary phase. Early observations of the increased radiation resistance of the budding cell fraction of yeast cultures [6] is now understood to be a reflection of the hypersensitive G1 and early S (DNA replication) stages, compared to the more resistant late S and G2 periods [15, 26, 27]. In general, this pattern is similar to that first observed in mammalian cells [161]. The variations in UV resistance could be due to different amounts

of initial DNA damage induced at different times in the cell cycle, or to varying efficiencies of repair mechanisms throughout the cycle. Although S-phase cells suffer 30% fewer pyrimidine dimers per unit UV dose than do cells having minimal resistance [14], this factor is not likely to be responsible for the bulk of the observed variations in UV sensitivity [25]. An excision repair-deficient (rad1) strain exhibits cyclic variations in UV sensitivity very similar to those found in a wild type, suggesting that excision repair acts efficiently and uniformly throughout the cell cycle [13]. By contrast, a rec5 strain, defective in UV-induced mitotic recombination, exhibits the same UV sensitivity in G1 and G2, a level comparable to the wild-type G1 level. This suggests that the increased G2 (over G1) UV resistance expressed by the wild type is due to a recombinational repair process that requires the REC5 gene product [13].

Davies et al [25] used a zonal rotor centrifugation method to isolate large yeast populations on the basis of bud size (correlated with progress in the cell cycle), an obvious improvement over the use of perturbing treatments that induce cell synchronony. Their results confirm earlier observations that in the wild type, UV resistance is minimal in G1 and maximum in S and G2; yields of UV-induced mitotic recombination (intergenic and intragenic) were maximum in G1 and minimum in S and G2, again suggesting a relationship between cell survival and recombination. Fabre [32] has shown that UV-induced intragenic mitotic recombination can occur in G1 before chromosome replication, confirming earlier results [170] and suggesting that homologous chromosomal pairing does not require duplicated chromatids. By contrast to UV, nitrous acid survival exhibited a minimum during only one period, that of DNA replication, while induced mitotic recombination occurred at all stages but was maximum during S [25].

Radiation and chemical mutagen sensitivity have also been compared in log-phase versus stationary-phase cultures [125]. As asynchronous log-phase cultures enter a transition period before entering stationary phase, cells tend to complete their cycles and begin arresting as unbudded cells (in G1 or more properly G<sub>0</sub>) [54]. Stationary-phase cells are observed to be more sensitive to UV and X-rays than log-phase cells, whereas just the reverse is true of sensitivity to several chemical mutagens. During the transition period, UV resistance begins to decrease in cultures that have already begun to show a significant reduction in the frequency of budded cells. While excision repair-defective (rad1 or rad2) strains also become more UV sensitive, a rad50 strain failed to exhibit this effect, suggesting a requirement for the RAD50 gene product. By contrast to UV, cell survival following treatments with nitrous acid, mitomycin C and ethyl-methanesulphonate increased in cultures beginning to show loss of budded cells. A significant fraction of this differential chemical mutagen sensitivity appears to be due to different numbers of initial DNA lesions inflicted, since cellular uptake of tritium-labelled EMS was 7-fold less in stationary cells compared to log-phase cells [125].

### DNA Replication

Until recently very little information has been available on the role of DNA replication in repair and mutagenesis of yeast. The main reason for this has been that only a few temperature-sensitive mutants have been described [52, 53, 72] that have large effects in turning off DNA synthesis specifically and rapidly after temperature shift. Some mutants also affect RNA synthesis; the gene product of many well-defined mutations are

not known. It has been suggested that a "replication complex" with one defective protein component might undergo slight conformational changes and still have some polymerizing activity [72]. Another reason concerns the fact that most in vivo studies eventually depend upon the conventional endpoint of colony (or mutant colony) formation, which in turn is dependent upon genome replication and cell division.

Yeast strains carrying cdc8 are defective in DNA replication (elongation not initiation) at 36°C but not at 23°C [52]. Prakash et al. [134a] have reported recently that cdc8 reduces frequencies of UV reversion (at 25°C), even in rad1 or rad51 strains, and argue that CDC8 plays a role in error-prone repair.

There now exists evidence that the temperature-sensitive cdc9-1 mutation [24] is defective in DNA ligase activity [73]. At the restrictive temperature this mutant 1) accumulates many single-strand breaks in DNA, 2) exhibits enhanced UV sensitivity, and 3) produces enhanced frequencies of spontaneous mitotic recombination (hyper-rec phenotype) [41]. These recent findings underscore the multiple role of this enzyme in DNA replication, repair and recombination in yeast. These authors suggest that the excess single-strand gaps in DNA are themselves recombinogenic, either directly or by means of the induction of a recombination repair system.

An important question concerns the kinetics of induced mutagenesis: Does it occur before, during, or after DNA replication? One approach might involve the use of a probe that can monitor the appearance of mutant (or recombinant) gene product as soon as it becomes expressed, rather than the phenotypic scoring of mutant (or recombinant) clones many generations removed from the initial mutagenic (or recombinogenic) event. Such a system has been previously used in Ustilago maydis to allow in vivo enzymatic measurement of radiation-induced mutation [97, 98] or mitotic recombination [65]. Another approach involves unambiguous detection of the "strandedness" of induced mutations. That is, fixation of premutational

damage in one strand of unreplicated (G1- phase) DNA is expected to give rise to a mosaic colony because after completion of the first cell cycle there will be one mutant and one nonmutant cell. Damage fixed as mutation in both strands prior to replication should lead to a pure mutant clone. Premutational damage not fixed as mutation until after the first round of replication will also produce mosaic clones.

James and Kilbey [70] observed UV induction of recessive lethal mutations in mitotic pedigrees of irradiated G1 diploid yeast cells. With this technique they found that after low exposures to UV, induced mutations were produced in an excision repair-proficient strain prior to the first round of post-irradiation DNA replication, and most mutations were 2-stranded. In an excision repair-defective (rad1) strain, induced mutations affecting both strands were not observed; moreover, mosaics arose as frequently in the second post-UV generation as in the first [71]. In rad1 strains unexcised pyrimidine dimers were shown to be responsible for UV mutagenesis even after passing through several DNA replication cycles [76], as Bridges and Munson had shown many years ago for E. coli [10]. Hannan et al. [50] had previously shown that G1 RAD haploid cells produced exclusively pure mutant clones after a UV exposure leading to high survival (63%, on the survival curve "shoulder"). Mosaics, however, were produced with increasing frequency for UV doses corresponding to exponentially decreasing survival, and could not be explained by first-division lethal sectoring. These findings support the idea that pure mutant clones are associated with efficient heteroduplex repair activity, such that loss of efficiency at higher UV doses leads to the induction of mosaics.

Kilbey et al. [75] have proposed a dimer/gap model to account for the different kinetics of UV mutagenesis in RAD and rad1 strains. According to the model, mutation fixation by error-prone repair is presumed to be initiated in both strains by a structure consisting of a single-stranded gap opposite a pyrimidine dimer, although this structure is produced in different ways by the two strains. In RAD strains after UV exposure sufficient to induce dimers close together on opposite complementary strands, the excision of one may often leave a gap that exposes the other. Since prereplicative mutagenesis (presumably an error-prone gap-filling process) eventually affects both DNA strands [70], then excision repair must remove the dimer, or heteroduplex repair must recognize and repair the mismatched site. In rad1 strains excision cannot occur and replication presumably generates daughter-strand gaps opposite pyrimidine dimers, followed by gap-filling and heteroduplex repair. This model is consistent with the observation that UV mutagenesis in RAD strains exhibits dose-squared dependence (2 dimers required) [75, 89, 100] compared to a linear dependence at low doses in rad1 [75] or rad2 [29] strains. Yet, there exists at least one case of linear induction in a RAD strain [30]. It is not altogether clear just how gap-filling generates single base-pair changes, which are responsible for the majority of UV mutations in yeast [90] rather than double base substitutions, as predicted by the bacterial model [12, 176]. An error-prone gap-filling model must also accommodate in some way the observations of nonrandom action (site-specificity of repair) and "mutation at a distance" [90].

Recent studies by Lemontt [99] have suggested that exposure of broth-grown stationary-phase yeast to HZ (a carcinogen and mutagen in other

organisms [77]) results in premutational DNA damage that becomes fixed as mutation at the time of DNA replication, as appears to be the case in Haemophilus [79, 80]. Unlike several other mutagens, HZ mutability at CAN1 is entirely dependent upon post-treatment DNA replication and occurs over a dose range that leaves cell viability unaffected. HZ exposure does not extend the 3-4 h growth lag normally observed in post-treatment medium. Prokaryotic studies have suggested that N<sup>4</sup>-aminocytosine may represent an important premutational DNA lesion to HZ-exposed cells [77]. Thus, unlike nonpairable pyrimidine dimers, this cytosine analogue might be considered a pairable lesion (with perhaps less than complete fidelity) produced in situ. N<sup>4</sup>-aminocytosine is known to be mutagenic in lambda phage and in E. coli when used as a precursor for DNA replication [17]. Thus, HZ may induce mutations in yeast by a mechanism of base mispairing at replication in the absence of any ostensible inhibitory effect on the replicative process itself, as proposed for Haemophilus. Kimball has suggested that N<sup>4</sup>-aminocytosine may be an intrinsically more efficient base-analogue mutagen than 5-bromouracil because the hydrazino ( $--NH_2NH_2$ ) substitution for the 4-amino group occurs at a base-pairing position on the pyrimidine ring, while the bromine substitution does not [78].

It has also been possible to obtain indirect evidence for prereplicative error-free repair of hydrazine-induced premutational damage [102]. As is also observed in Haemophilus [80] if post-treatment DNA replication or its initiation is delayed in growth medium (in yeast with hydroxyurea or cycloheximide, respectively), the maximum level of replication-dependent mutagenesis attainable after removal of inhibitors decreases (Table 4). Excision repair-defective rad2-1 strains also exhibit such loss of HZ mutability. These findings have suggested that premutational lesions are

T-4

being removed by some error-free process different from excision repair of pyrimidine dimers. Since cycloheximide blocks protein synthesis (which is required for initiation of DNA replication in yeast [57, 171]), this repair process must be constitutive. It is possible to speculate that some form of mismatch repair may be operating. Assuming that N<sup>4</sup>-aminocytosine is a major premutational lesion, the duplex distortion (presumed to occur by virtue of a hydrazino rather than an amino proton donor in hydrogen bonding to guanine) might also be correctable by specific cleavage of the terminal amino group, restoring normal base pairing without the need for strand breaks in the backbone. Examining hydrazine-treated Haemophilus, Kimball and Hirsch [80] failed to detect single-strand breaks (or alkali-labile sites) in unreplicated DNA, nor did they observe gaps in newly synthesized DNA.

#### Mating type locus-dependent regulation

Normal conjugation in yeast occurs between cells of opposite mating type, either  $\bar{a}$  or  $\alpha$ . These two mating phenotypes segregate in meiosis as different alleles of the same locus, called the mating-type locus (MAT). There are now many lines of evidence supporting the idea that the genetic information at MAT has a regulatory function that plays a central role in controlling whether a cell may undergo sexual conjugation or pursue meiotic development. Diploids exhibit one of three possible functional states at MAT --  $\bar{a}/\bar{a}$   $\alpha/\alpha$ , or  $\bar{a}/\alpha$ . Like  $\bar{a}$  or  $\alpha$  haploids, homozygous  $\bar{a}/\bar{a}$  or  $\alpha/\alpha$  diploids (selected by mitotic crossing over) are able to mate normally (and produce and respond to mating pheromones), exhibit medial bud initiation in mitosis, but cannot initiate meiosis and sporulation when challenged to do so in the appropriate medium. Such homozygotes

exhibit  $\tilde{a}$ -specific or  $\alpha$ -specific functions expressed by haploids. Diploids heterozygous ( $\tilde{a}/\alpha$ ) at MAT are repressed in mating ability (and fail to produce or respond to mating pheromones), exhibit polar bud initiation, and have gained meiotic and sporulative capacity. The  $\tilde{a}/\alpha$  state appears not only to turn off certain haploid functions, but also to turn on certain new diploid functions. Moreover, the expression of these  $\tilde{a}$ -specific,  $\alpha$ -specific and  $\tilde{a}/\alpha$ -specific functions may be altered by a variety of mutations in several genes unlinked to MAT (for recent reviews of mating-type-specific functions, see [23, 107, 109]). Thus, in mitotic cells the existence of phenotypic differences expressed by MAT homozygotes compared to the "normal" (after normal  $\tilde{a} \times \alpha$  mating) MAT heterozygotes constitutes evidence for MAT regulation.

DNA repair, mitotic recombination and mutagenesis all appear to be modulated to some degree by MAT. Although diploid yeast is much more X-ray resistant than haploid (the ploidy effect [117]), MAT homozygotes are more sensitive than  $\tilde{a}/\alpha$  [85, 117] cells suggesting that a fraction of the extra diploid resistance is due to MAT heterozygosity. Liquid-holding recovery in buffer after X-ray exposure (which does not occur in haploids) is believed to be controlled in part by a MAT-dependent process (Hunnable and Cox, cited in [23]). Moreover, Game and Mortimer [43] have found that some mutants (rad50, rad57) in the RAD52 pathway for repair of ionizing radiation damage exhibited a MAT effect ( $\tilde{a}/\alpha$  more resistant than  $\tilde{a}/\tilde{a}$  or  $\alpha/\alpha$ ), while others (rad52, rad54) did not show the MAT effect. Thus, RAD52 and RAD54 may act prior to MAT-dependent repair steps, whereas RAD50 and RAD57 may control subsequent MAT-independent steps [43]. A MAT effect for MMS sensitivity has also been reported [106].

Although UV survival does not show the MAT effect [86] UV-induced mitotic recombination does [36]. Frequencies of induced mitotic gene conversion in  $\underline{a}/\underline{a}$  or  $\alpha/\alpha$  diploids were as much as 100-fold lower than in  $\underline{a}/\alpha$  strains, again suggesting that the  $\underline{a}/\alpha$  regulatory state is required for maximal expression of induced mitotic recombination.

There is evidence that MAT regulation can affect UV mutagenesis. Martin, Prakash and Prakash [11] have found that although mms3 causes UV sensitivity in haploids and diploids,  $\underline{a}/\alpha$  mms3/mms3 diploids exhibit defective UV reversion of arg4-17 or lys2-1, compared to  $\underline{a}/\alpha$  mms3/MMS and  $\underline{a}/\alpha$  MMS/MMS diploids; mms3 haploids have wild-type UV revertibility. In addition,  $\underline{a}/\underline{a}$  or  $\alpha/\alpha$  derivatives of the  $\underline{a}/\alpha$  mms3/mms3 strain were restored to normal UV mutability. This shows that the  $\underline{a}/\alpha$  genetic configuration is responsible for the diploid-specific defective UV mutability.

Finally,  $\alpha$  umr7-1 haploids fail to express several  $\alpha$ -specific haploid functions (such as mating ability,  $\alpha$ -factor production,  $\underline{a}$ -factor response), while at the same time, they have apparently turned on some  $\underline{a}$ -specific functions ("shmoo" morphology,  $\alpha$ -factor proteolysis [34, 35]); on the other hand,  $\underline{a}$  umr7 haploids express normal  $\underline{a}$ -specific functions [104, 105]. Both types of umr7 strains are enormously flocculent (clumpy) but can be dispersed by distilled-water washing [100]. Although these strains are defective in UV mutagenesis at CAN1 [100], it seems likely that this is due to an aberrant cell envelope, which interferes in some way with normal expression of mutant arginine permease. This is supported by the observation that nonclumpy revertant derivatives exhibit wild-type levels of UV mutability at CAN1 [104].

REFERENCES

1. Ananthaswamy, H. N., T. J. McKey and R. K. Mortimer, Isolation and characterization of additional X-ray sensitive mutants of *Saccharomyces cerevisiae* (abs.), Ninth International Conference on Yeast Genetics and Molecular Biology, Rochester, N.Y., 1978, p. 45.
2. Averbeck, D., W. Laskowski, E. Eckardt, E. Lehmann-Brauns, Four radiation sensitive mutants of *Saccharomyces*. Survival after UV- and X-ray-irradiation as well as UV-induced reversion rates from isoleucine-valine dependence to independence, *Molec. Gen. Genet.*, 107 (1970) 117-127.
3. Bacila, M., B. L. Horecker and A. O. M. Stoppani (eds.) *Biochemistry and Genetics of Yeast*, Academic Press, New York, 1978.
4. Baker, B., A. T. C. Carpenter, M. S. Esposito, R. E. Esposito and L. Sandler, The genetic control of meiosis, *Ann. Rev. Genet.*, 10 (1976) 53-134.
5. Bandlow, W., R. J. Schweyen, K. Wolf and F. Kaudewitz (eds.) *Mitochondria 1977. Genetics and Biogenesis of Mitochondria*, Walter de Gruyter, New York, 1977.
6. Beam, C. A., R. K. Mortimer, R. G. Wolfe, and C. A. Tobias, The relation of radioresistance to budding in *Saccharomyces cerevisiae*, *Arch. Biochem. Biophys.*, 49 (1954) 110-122.
7. Boram, W. R. and H. Roman, Recombination in *Saccharomyces cerevisiae*: A DNA repair mutation associated with elevated mitotic gene conversion, *Proc. Natl. Acad. Sci. (U.S.A.)*, 73 (1976) 2828-2832.
8. Bridges, B. A., *Mechanisms of radiation mutagenesis in cellular and subcellular systems*, *Ann. Rev. Nuc. Sci.*, 19 (1969) 139-178.
9. Bridges, B. A., R. P. Mottershead and S. G. Sedgwick, Mutagenic DNA repair in *Escherichia coli* III. Requirement for a function of DNA polymerase III in ultraviolet-light mutagenesis, *Molec. Gen. Genet.*, 144 (1976) 53-58.
10. Bridges, B. A. and R. J. Munson, The persistence through several replication cycles of mutation-producing pyrimidine dimers in a strain of *Escherichia coli* deficient in excision-repair, *Biochem. Biophys. Res. Commun.*, 30 (1968) 620-624.
11. Brusick, D. J., Induction of cycloheximide-resistant mutants in *Saccharomyces cerevisiae* with N-methyl-N'-Nitro-N-nitrosoguanidine and ICR-170, *J. Bacteriol.*, 109 (1972) 1134-1138.

12. Caillet-Fauquet, P., M. Defais and M. Radman, Molecular mechanisms of induced mutagenesis. Replication in vivo of bacteriophage  $\phi$ X174 single-stranded, ultraviolet light-irradiated DNA in intact and irradiated host cells, *J. Mol. Biol.*, 117 (1977) 95-112.
13. Chanet, R., M. Heude and E. Moustacchi, Variations in UV-induced lethality and "petite" mutagenesis in synchronous culture of *Saccharomyces cerevisiae*. II. Responses of radiosensitive mutants to lethal damage, *Molec. Gen. Genet.*, 132 (1974) 23-30.
14. Chanet, R., R. Waters and E. Moustacchi, The induction of pyrimidine dimers in nuclear DNA after UV-irradiation during the synchronous cycle of *Saccharomyces cerevisiae*, *Int. J. Radiat. Biol.*, 27 (1975) 481-485.
15. Chanet, R., D. H. Williamson and E. Moustacchi, Cyclic variations in killing and "petite" mutagenesis induced by ultraviolet light in synchronized yeast strains, *Biochim. Biophys. Acta*, 324 (1973) 290-299.
16. Chasin, L. A., Mutation affecting adenine phosphoribosyl transferase activity in Chinese hamster cells, *Cell*, 2 (1974) 37-41.
17. Chu, B. C. F., D. M. Brown and M. G. Burdon, Effect of nitrogen and of catalase on hydroxylamine and hydrazine mutagenesis, *Mutation Res.*, 20 (1973) 265-270.
18. Clark, A. J. and A. D. Margulies, Isolation and characterization of recombination-deficient mutants of *Escherichia coli* K12, *Proc. Natl. Acad. Sci. (U.S.A.)*, 53 (1965) 451-459.
19. Cox, B. S., Pathways of UV repair and mutagenesis in *Saccharomyces cerevisiae*, in A. Castellani (ed.), *Research in Photobiology*, Plenum Press, New York, 1976, pp. 689-697.
20. Cox, B. S. and E. A. Bevan, Aneuploidy in yeast, *New Phytol.*, 61 (1962) 342-355.
21. Cox, B. S. and J. Game, Repair systems in *Saccharomyces*, *Mutation Res.*, 26 (1974) 257-264.
22. Cox, B. S. and J. M. Parry, The isolation, genetics and survival characteristics of ultraviolet light-sensitive mutants in yeast, *Mutation Res.*, 6 (1968) 37-55.
23. Crandall, M., R. Egel and V. L. MacKay, Physiology of mating in three yeasts, *Adv. Microb. Physiol.*, 15 (1977) 307-398.
24. Culotti, J. and L. H. Hartwell, Genetic control of the cell division cycle. III Seven genes controlling nuclear division, *Exp. Cell Res.*, 67 (1971) 389-401.

25. Davies, P. J., R. S. Tippins and J. M. Parry, Cell-cycle variations in the induction of lethality and mitotic recombination after treatment with UV and nitrous acid in the yeast, *Saccharomyces cerevisiae*, *Mutation Res.*, 51 (1978) 327-346.
26. de Langguth, E. N. and C. A. Beam, Repair mechanisms and cell cycle dependent variations in X-ray sensitivity of diploid yeast, *Radiat. Res.*, 53 (1973) 226-234.
27. de Langguth, E. N. and C. A. Beam, The effects of ploidy upon cell cycle dependent changes in X-ray sensitivity of *Saccharomyces cerevisiae*, *Radiat. Res.*, 55 (1973) 501-506.
28. Drake, J. W. and E. F. Allen, Antimutagenic DNA polymerases of bacteriophage T4, *Cold Spring Harbor Symp. Quant. Biol.*, 33 (1968) 339-341.
29. Eckardt, F. and R. H. Haynes, Kinetics of mutation induction by ultraviolet light in excision-deficient yeast, *Genetics*, 85 (1977) 225-247.
30. Eckardt, F. and R. H. Haynes, Induction of pure and sectoried mutant clones in excision-proficient and deficient strains of yeast, *Mutation Res.*, 43 (1977) 327-338.
31. Esposito, M. S., M. Bolotin-Fukuhara and R. E. Esposito, Antimutator activity during mitosis by a meiotic mutant of yeast, *Molec. Gen. Genet.*, 139 (1975) 9-18.
32. Fabre, F., Induced intragenic recombination in yeast can occur during the G<sub>1</sub> mitotic phase, *Nature*, 272 (1978) 795-798.
33. Fabre, F. and H. Roman, Genetic evidence for inducibility of recombination competence in yeast, *Proc. Natl. Acad. Sci.(U.S.A.)*, 74 (1977) 1667-1671.
34. Finkelstein, D., personal communication.
35. Finkelstein, D. B. and S. Strausberg, Metabolism of  $\alpha$ -factor by a mating type cells of *Saccharomyces cerevisiae*, *J. Biol. Chem.*, 254 (1979) 796-803.
36. Friis, J. and H. Roman, The effect of the mating-type alleles on intragenic recombination in yeast, *Genetics*, 59 (1968) 33-36.
37. Game, J. C., Radiation-sensitive mutants of yeast, in P. C. Hanawalt and R. B. Setlow (eds.), *Molecular Mechanisms for Repair of DNA*, Plenum Press, New York, 1975, pp. 541-544.
38. Game, J. C. and B. S. Cox, Allelism tests of mutants affecting sensitivity to radiation in yeast and a proposed nomenclature, *Mutation Res.*, 12 (1971) 328-331.

39. Game, J. C. and B. S. Cox, Epistatic interactions between four rad loci in yeast, *Mutation Res.*, 16 (1972) 353-362.
40. Game, J. C. and B. S. Cox, Synergistic interactions between RAD mutations in yeast, *Mutation Res.*, 20 (1973) 35-44.
41. Game, J. C., L. H. Johnston and R. C. von Borstel, Enhanced mitotic recombination in a ligase-defective mutant of the yeast *Saccharomyces cerevisiae*, *Proc. Natl. Acad. Sci.*, submitted.
42. Game, J. C. and R. K. Mortimer, A genetic study of X-ray sensitive mutants in yeast, *Mutation Res.*, 24 (1974) 281-292.
43. Game, J. C. and R. K. Mortimer, personal communication.
44. Game, J. C., T. J. Zamb, R. J. Braun, M. Resnick and R. M. Roth, The role of radiation (rad) genes in meiotic recombination in yeast, *Genetics*, submitted.
45. Gocke, E. and T. R. Manney, Expression of radiation-induced mutations at the arginine permease (CAN1) locus in *Saccharomyces cerevisiae*, *Genetics*, 91 (1979) 53-66.
46. Golin, J. E. and M. S. Esposito, Evidence for joint genic control of spontaneous mutation and genetic recombination during mitosis in *Saccharomyces*, *Molec. Gen. Genet.*, 150 (1977) 127-135.
47. Grenson, M. and C. Hou, Ammonia inhibition of the general amino acid permease and its suppression in NADPH-specific glutamate dehydrogenaseless mutants of *Saccharomyces cerevisiae*, *Biochem. Biophys. Res. Commun.*, 48 (1972) 749-756.
48. Grenson, M., C. Hou and M. Crabeel, Multiplicity of the amino acid permeases in *Saccharomyces cerevisiae* IV. Evidence for a general amino acid permease, *J. Bacteriol.*, 103 (1970) 770-777.
49. Grenson, M., M. Mousset, J. M. Wiame, and J. Bechet, Multiplicity of the amino acid permeases in *Saccharomyces cerevisiae*. I. Evidence for a specific arginine-transporting system, *Biochem. Biophys. Acta.*, 127 (1966) 325-338.
50. Hannan, M. A., P. Duck and A. Nasim, UV-induced lethal sectoring and pure mutant clones in yeast, *Mutation Res.*, 36 (1976) 171-176.
51. Hartwell, L. H., Biochemical genetics of yeast, *Ann. Rev. Genet.*, 4 (1970) 373-396.
52. Hartwell, L. H., Genetic control of the cell division cycle in yeast II. Genes controlling DNA replication and its initiation, *J. Mol. Biol.*, 59 (1971) 183-194.
53. Hartwell, L. H., Three additional genes required for DNA synthesis in *Saccharomyces cerevisiae*, *J. Bacteriol.*, 115 (1973) 966-974.

54. Hartwell, L. H. *Saccharomyces cerevisiae* cell cycle, *Bacteriol. Rev.*, 38 (1974) 164-198.
55. Hastings, P. J., S.-K. Quah and R. C. von Borstel, Spontaneous mutation by mutagenic repair of spontaneous lesions in DNA, *Nature*, 264 (1976) 719-722.
56. Haynes, R. H., DNA repair and the genetic control of radiosensitivity in yeast, in P. C. Hanawalt and R. B. Setlow (eds.), *Molecular Mechanisms for Repair of DNA*, Plenum Press, New York, 1975, pp. 529-540.
57. Hereford, L. M. and L. H. Hartwell, Role of protein synthesis in the replication of yeast DNA, *Nature New Biol.*, 244 (1973) 129-131.
58. Hershfield, M. S. and N. G. Nossal, *In vitro* characterization of a mutator T4 DNA polymerase, *Genetics Suppl.*, 73 (1973) 131-136.
59. Hill, R. F., Ultraviolet-induced lethality and reversion to prototrophy in *Escherichia coli* strains with normal and reduced repair ability, *Photochem. Photobiol.*, 4 (1965) 563-568.
60. Ho., K., Induction of DNA double-strand breaks by X-rays in a radio-sensitive strain of the yeast *Saccharomyces cerevisiae*, *Mutation Res.*, 30 (1975) 327-334.
61. Ho, K. S. Y. and R. K. Mortimer, Induction of dominant lethality by X-rays in a radiosensitive strain of yeast, *Mutation Res.*, 20 (1973) 45-51.
62. Holliday, R., A mechanism for gene conversion in fungi, *Genet. Res. Camb.*, 5 (1964) 282-304.
63. Holliday, R., Radiation-sensitive mutants of *Ustilago maydis*, *Mutation Res.*, 2 (1965) 557-559.
64. Holliday, R., Altered recombination frequencies in radiation-sensitive strains of *Ustilago*, *Mutation Res.*, 4 (1967) 275-288.
65. Holliday, R., Biochemical measure of the time and frequency of radiation-induced allelic recombination in *Ustilago*, *Nature New Biol.*, 232 (1971) 233-236.
66. Holliday, R., *Ustilago maydis*, in R. C. King (ed.), *Handbook of Genetics*, vol. 1, Plenum Press, New York, 1974, pp. 575-595.
67. Holliday, R., R. E. Halliwell, M. W. Evans and V. Rowell, Genetic characterization of *rec-1*, a mutant of *Ustilago maydis* defective in repair and recombination, *Genet. Res. Camb.*, 27 (1976) 413-453.
68. Howard-Flanders, P., DNA repair, *Ann. Rev. Biochem.*, 37 (1968) 175-200.

69. Howard-Flanders, P., L. Theriot, and J. B. Stedeford, Some properties of excision-defective, recombination-defective mutants of *Escherichia coli* K12, *J. Bacteriol.*, 97 (1969) 1134-1141.
70. James, A. P. and B. J. Kilbey, The timing of UV mutagenesis in yeast: A pedigree analysis of induced recessive mutation, *Genetics*, 87 (1977) 237-248.
71. James, A. P., B. J. Kilbey and G. J. Prefontaine, The timing of UV mutagenesis in yeast: Continuing mutation in an excision-defective (rad-1) strain, *Molec. Gen. Genet.*, 165 (1978) 207-212.
72. Johnston, L. H. and J. C. Game, Mutants of yeast with depressed DNA synthesis, *Molec. Gen. Genet.*, 161 (1978) 205-214.
73. Johnston, L. H. and K. A. Nasmyth, *Saccharomyces cerevisiae* cell cycle mutant cdc9 is defective in DNA ligase, *Nature*, 274 (1978) 891-893.
74. Kato, T. and Y. Shinoura, Isolation and characterization of mutants of *Escherichia coli* deficient in induction of mutations by ultraviolet light, *Molec. Gen. Genet.*, 156 (1977) 121-131.
75. Kilbey, B. J., T. Brychcy and A. Nasim, Initiation of UV mutagenesis in *Saccharomyces cerevisiae*, *Nature*, 274 (1978) 889-891.
76. Kilbey, B. J. and A. P. James, The mutagenic potential of unexcised pyrimidine dimers in *Saccharomyces cerevisiae*, RAD1-1. Evidence from photoreactivation and pedigree analysis, *Mutation Res.*, 60 (1979) 163-171.
77. Kimball, R. F., The mutagenicity of hydrazine and some of its derivatives, *Mutation Res.*, 39 (1977) 111-126.
78. Kimball, R. F., The relation of repair phenomena to mutation induction in bacteria, *Mutation Res.*, 55 (1978) 85-120.
79. Kimball, R. F. and B. F. Hirsch, Tests for the mutagenic action of a number of chemicals on *Haemophilus influenzae* with special emphasis on hydrazine, *Mutation Res.*, 30 (1975) 9-20.
80. Kimball, R. F. and B. F. Hirsch, Fixation and loss of hydrazine-induced premutational damage in *Haemophilus influenzae*, *Mutation Res.*, 36 (1976) 39-48.
81. Korch, C. T. and R. Snow, Allelic complementation in the first gene for histidine biosynthesis in *Saccharomyces cerevisiae* I. Characteristics of mutants and genetic mapping in alleles, *Genetics*, 74 (1973) 287-305.
82. Kowalski, S. and W. Laskowski, The effect of three rad genes on survival, inter- and intragenic mitotic recombination in *Saccharomyces*, *Molec. Gen. Genet.*, 136 (1975) 75-86.

83. Larimer, F. W., personal communication.
84. Larimer, F. W., D. Ramey, W. Lijinsky, and J. L. Epler, Mutagenicity of methylated N-nitrosopiperidines in *Saccharomyces cerevisiae*, *Mutation Res.*, 57 (1978) 155-161.
85. Laskowski, W., Inaktivierungsversuche mit homozyoten Hefestämmen verschiedenen Ploidilgrades, *Z. Naturforsch.*, 15b (1960) 495-506.
86. Laskowski, W., Der  $\alpha$ -Effect, eine Korrelation Zwischen Paarungstypen-konstitution und Strahlenresistenz bei Hefen, *Zbl. Bakteriol. Parasitenkunde, Infektionskrankh, Hyg. 1 Orig.*, 184 (1962) 251-258.
87. Lawrence, C. W. and R. Christensen, UV mutagenesis in radiation-sensitive strains of yeast, *Genetics*, 82 (1976) 207-232.
88. \_\_\_\_\_, Ultraviolet-induced reversion of *cycl* alleles in radiation-sensitive strains of yeast. I. *rev1* mutant strains, *J. Mol. Biol.*, 122 (1978) 1-21.
39. \_\_\_\_\_, Ultraviolet-induced reversion of *cycl* alleles in radiation-sensitive strains of yeast II. *rev2* mutant strains, *Genetics*, 90 (1978) 213-226.
90. \_\_\_\_\_, Ultraviolet light induced mutagenesis in *Saccharomyces cerevisiae*, in P. C. Hanawalt and R. B. Setlow (eds.), *DNA Repair Mechanisms*, Academic Press, New York, 1978, pp. 437-440.
91. \_\_\_\_\_, Ultraviolet-induced reversion of *cycl* alleles in radiation-sensitive strains of yeast III. *rev3* mutant strains, *Genetics*, in press.
92. Lawrence, C. W., J. W. Stewart, F. Sherman and R. Christensen, Specificity and frequency of ultraviolet-induced reversion of an iso-1-cytochrome c ochre mutant in radiation-sensitive strains of yeast, *J. Mol. Biol.*, 85 (1974) 137-162.
93. Lemontt, J. F., Mutants of yeast defective in mutation induced by ultraviolet light, *Genetics*, 68 (1971) 21-33.
94. \_\_\_\_\_, Pathways of ultraviolet mutability in *Saccharomyces cerevisiae*. II. The effect of *rev* genes on recombination, *Mutation Res.*, 13 (1971) 319-326.
95. \_\_\_\_\_, Induction of forward mutations in mutationally defective yeast, *Molec. Gen. Genet.*, 119 (1972) 27-42.
96. \_\_\_\_\_, Genes controlling ultraviolet mutability in yeast, *Genetics Suppl.*, 73 (1973) 153-159.
97. \_\_\_\_\_, Induced mutagenesis in *Ustilago maydis*. I. Isolation and characterization of a radiation-revertible allele of the structural gene for nitrate reductase, *Molec. Gen. Genet.*, 145 (1976) 125-132.

98. Lemontt, J. F., Induced mutagenesis in *Ustilago maydis* II. An in vivo biochemical assay, *Molec. Gen. Genet.*, 145 (1976) 133-143.
99. \_\_\_\_\_, Mutagenesis of yeast by hydrazine: Dependence upon post-treatment cell division, *Mutation Res.*, 43 (1977) 165-178.
100. \_\_\_\_\_, Pathways of ultraviolet mutability in *Saccharomyces cerevisiae* III. Genetic analysis and properties of mutants resistant to ultraviolet-induced forward mutation, *Mutation Res.*, 43 (1977) 179-204.
101. \_\_\_\_\_, Pathways of ultraviolet mutability in *Saccharomyces cerevisiae* IV. The relation between canavanine toxicity and ultraviolet mutability to canavanine resistance, *Mutation Res.*, 43 (1977) 339-355.
102. \_\_\_\_\_, Loss of hydrazine-induced mutability in wild-type and excision-repair-defective yeast during post-treatment inhibition of cell division, *Mutation Res.*, 50 (1978) 57-66.
103. \_\_\_\_\_, unpublished data.
104. Lemontt, J. F., D. R. Fugit and V. L. MacKay, Pleiotropic mutations at the IUP1 locus that affect the expression of mating-type-dependent functions in *Saccharomyces cerevisiae*, *Genetics*, submitted.
105. Lemontt, J. F. and V. L. MacKay, A pleiotropic mutant of yeast expressing the mating-specific "shmoo" morphology during vegetative growth in the absence of exogenous mating hormone (abs.), *Genetics Suppl.*, 86 (1977) s38.
106. Livi, George P. and V. L. MacKay, Mating-type regulation of methyl methanesulfonate sensitivity in *Saccharomyces cerevisiae*, *Genetics*, submitted.
107. MacKay, V. L., Mating-type-specific pheromones as mediators of sexual conjugation in yeast, in J. Papaconstantinou and W. J. Rutter (eds.) *Molecular Control of Proliferation and Differentiation*, Academic Press, New York, 1978, pp. 243-259.
108. Maloney, D. and S. Fogel, High frequency mitotic conversion mutants in yeast (abs.), *Genetics Suppl.*, 83 (1976) s47.
109. Manney, T. R. and J. H. Meade, Cell-cell interactions during mating in *Saccharomyces cerevisiae*, in J. L. Reissig (ed.), *Microbiol Interactions, Receptors and Recognition*, Series B, vol. 3, Chapman and Hall, London, 1977, pp. 281-321.
110. Manney, T. R. and R. K. Mortimer, Allelic mapping in yeast by X-ray-induced mitotic reversion, *Science*, 143 (1964) 581-582.

111. Martin, P., S. Prakash and L. Prakash, personal communication.
112. McKee, R. H. and C. W. Lawrence, Genetic analysis of gamma ray mutagenesis in yeast. I. Reversion in radiation-sensitive strains, Genetics, submitted.
113. McKee, R. H. and C. W. Lawrence, Genetic analysis of gamma ray mutagenesis in yeast. Survival and reversion in double mutant strains, Mutation Res., submitted.
114. McKee, R. H. and C. W. Lawrence, Genetic analysis of gamma ray mutagenesis in yeast. II. Allele-specific control of mutagenesis. Genetics, submitted.
115. Moore, C. and F. Sherman, Role of DNA sequences in genetic recombination in the iso-1-cytochrome c gene of yeast. I. Discrepancies between physical distance and genetic distance determined by five mapping procedures, Genetics, 79 (1975) 397-418.
116. Moore, C. and F. Sherman, Role of DNA sequences in genetic recombination in the iso-1-cytochrome c gene of yeast. II. Comparison of mutants altered at the same and nearby base pairs, Genetics, 85 (1977) 1-22.
117. Mortimer, R. K., Radiobiological and genetic studies on a polyploid series (haploid to hexaploid) of *Saccharomyces cerevisiae*, Radiat. Res., 9 (1958) 312-326.
118. Mortimer, R. K. and D. C. Hawthorne, Yeast genetics, in A. H. Rose and J. S. Harrison (eds.) The Yeasts, vol. 1, Academic Press, New York, 1969, pp. 385-460.
119. Mount, D. W., K. B. Low, and S. J. Edmiston, Dominant mutations (lex) in *Escherichia coli* K-12 which affect radiation sensitivity and frequency of ultraviolet light-induced mutations, J. Bacteriol., 112 (1972) 886-893.
120. Moustacchi, E., Cytoplasmic and nuclear genetic events induced by UV light in strains of *Saccharomyces cerevisiae* with different UV sensitivities, Mutation Res., 7 (1969) 171-185.
121. Moustacchi, E., R. Chanet and M. Heude, Ionizing and ultraviolet radiations: Genetic effects and repair in yeast, in A. Castellani (ed.), Research in Photobiology, Plenum Press, New York, 1976, pp. 197-206.
122. Nakai, S. and S. Matsumoto, Two types of radiation-sensitive mutant in yeast, Mutation Res., 4 (1967) 129-136.
- 122a. Nakai, S. and R. K. Mortimer, Studies of the genetic mechanism of radiation-induced mitotic segregation in yeast, Molec. Gen. Genet., 103 (1969) 329-338.

123. Parker, J. H. and F. Sherman, Fine-structure mapping and mutational studies of gene controlling yeast cytochrome c, *Genetics*, 62 (1969) 9-22.
124. Parry, E. M. and B. S. Cox, The tolerance of aneuploidy in yeast, *Genet. Res. Camb.*, 16 (1970) 333-340.
125. Parry, J. M., P. J. Davies and W. E. Evans, The effects of "cell age" upon the lethal effects of physical and chemical mutagens in the yeast, *Saccharomyces cerevisiae*, *Molec. Gen. Genet.*, 146 (1976) 27-35.
126. Parry, J. M. and E. M. Parry, The effects of UV light post-treatments on the survival characteristics of 21 UV-sensitive mutants of *Saccharomyces cerevisiae*, *Mutation Res.*, 8 (1969) 545-556.
127. Patrick, M. H., R. H. Haynes and R. B. Uretz, Dark recovery phenomena in yeast. I Comparative effects with various inactivating agents, *Radiat. Res.*, 21 (1964) 144-163.
128. Patrick, M. H. and R. H. Haynes, Dark recovery phenomena in yeast. II Conditions that modify the recovery process, *Radiat. Res.*, 23 (1964) 564-579.
129. Prakash, L., Lack of chemically induced mutation in repair-deficient mutants of yeast, *Genetics*, 78 (1974) 1101-1118.
130. \_\_\_\_\_, Repair of pyrimidine dimers in nuclear and mitochondrial DNA of yeast irradiated with low doses of ultraviolet light, *J. Mol. Biol.*, 98 (1975) 781-795.
131. \_\_\_\_\_, Effect of genes controlling radiation sensitivity on chemically induced mutations in *Saccharomyces cerevisiae*, *Genetics*, 83 (1976) 285-301.
132. \_\_\_\_\_, The relation between repair of DNA and radiation and chemical mutagenesis in *Saccharomyces cerevisiae*, *Mutation Res.*, 41 (1976) 241-248.
133. \_\_\_\_\_, Defective thymine dimer excision in radiation-sensitive mutants *rad10* and *rad16* of *Saccharomyces cerevisiae*, *Molec. Gen. Genet.*, 152 (1977) 125-128.
134. \_\_\_\_\_, Repair of pyrimidine dimers in radiation-sensitive mutants *rad3*, *rad4*, *rad6*, and *rad9* of *Saccharomyces cerevisiae*, *Mutation Res.*, 45 (1977) 13-20.
- 134a. Prakash, L., D. Hinkle and S. Prakash, Decreased UV mutagenesis in *cdc8*, a DNA replication mutant of *Saccharomyces cerevisiae*, *Molec. Gen. Genet.*, 172 (1979) 249-258.
135. Prakash, L. and S. Prakash, Isolation and characterization of MMS-sensitive mutants of *Saccharomyces cerevisiae*, *Genetics*, 86 (1977) 33-55.

136. Prakash, S. and L. Prakash, Increased spontaneous mitotic segregation in MMS-sensitive mutants of *Saccharomyces cerevisiae*, *Genetics*, 87 (1977) 229-236.
137. Quah, S.-K., personal communication.
138. Quah, S.-K., R. C. von Borstel and P. J. Hastings, Antimutators in yeast (abs.), Cold Spring Harbor Laboratory Meeting on the Molecular Biology of Yeast, 1977, p. 82.
139. Rahn, R. O., N. P. Johnson, A. W. Hsie, J. F. Lemontt, W. E. Masker, J. D. Regan, W. C. Dunn, J. D. Hoeschele, and D. H. Brown, The interaction of platinum compounds with the genome: Correlation between DNA binding and biological effects, in H. R. Witschi (ed.), *The Scientific Basis of Toxicity Assessment*, Elsevier/North Holland, Amsterdam, in press.
140. Resnick, M. A., Genetic control of radiation sensitivity in *Saccharomyces cerevisiae*, *Genetics*, 62 (1969) 519-531.
141. \_\_\_\_\_, Induction of mutations in *Saccharomyces cerevisiae* by ultraviolet light, *Mutation Res.*, 7 (1969) 315-332.
142. \_\_\_\_\_, The induction of molecular and genetic recombination in eukaryotic cells, *Adv. Radiat. Biol.*, 8 (1978) 175-216.
143. Resnick, M. A. and P. Martin, The repair of double-strand breaks in the nuclear DNA of *Saccharomyces cerevisiae* and its genetic control, *Molec. Gen. Genet.*, 143 (1976) 119-129.
144. Resnick, M. A. and J. K. Setlow, Repair of pyrimidine dimer damage induced in yeast by ultraviolet light, *J. Bacteriol.*, 109 (1972) 979-986.
145. Reynolds, R. J., Removal of pyrimidine dimers from *Saccharomyces cerevisiae* nuclear DNA under nongrowth conditions as detected by a sensitive, enzymatic assay, *Mutation Res.*, 50 (1978) 43-56.
146. Rodarte-Ramon, U.S., Radiation-induced recombination in *Saccharomyces*: The genetic control of recombination in mitosis and meiosis, *Radiat. Res.*, 49(1972) 148-154.
147. Rodarte-Ramon, U. S. and R. K. Mortimer, Radiation-induced recombination in *Saccharomyces*: Isolation and genetic study of recombination-deficient mutants, *Radiat. Res.*, 49 (1972) 133-147.
148. Roman, H., A system selective for mutations affecting the synthesis of adenine in yeast, *Compt. Rend. Trav. Lab. Carls.*, 26 (1956) 299-314.
149. Rose, A. H. and J. S. Harrison (eds.), *The Yeasts*, Vol. 1, *Biology of Yeasts*, Academic Press, New York, 1969.
150. Rose, A. H. and J. S. Harrison (eds.) *The Yeasts*, Vol. 2, *Physiology and Biochemistry of Yeasts*, Academic Press, New York, 1971.

151. Rothstein, R. J. and F. Sherman, Genes affecting the expression of cytochrome *c* in yeast: Genetic mapping and genetic interactions, *Genetics*, submitted.
152. Rupp, W. D. and P. Howard-Flanders, Discontinuities in the DNA synthesized in an excision-defective strain of *Escherichia coli* following ultraviolet irradiation. *J. Mol. Biol.*, 131 (1968) 291-304.
153. Schnaar, R. L., N. Muzyczka and M. J. Bessman, Utilization of aminopurine deoxynucleoside triphosphate by mutator, anti-mutator and wild-type DNA polymerases of bacteriophage T4, *Genetics Suppl.*, 73 (1973) 137-140.
154. Sherman, F. and C. W. Lawrence, *Saccharomyces*, in R. C. King (ed.) *Handbook of Genetics*, vol. 1, Plenum Press, New York, 1974, pp. 359-393.
155. Sherman, F. and J. W. Stewart, Variation of mutagenic action on nonsense mutants at different sites in the iso-1-cytochrome *c* gene of yeast, *Genetics*, 78 (1974) 97-113.
156. Smith, K. C., and D. H. C. Meun, Repair of radiation-induced damage in *Escherichia coli*. I. Effect of *rec* mutations on postreplication repair of damage due to ultraviolet radiation, *J. Mol. Biol.*, 51 (1970) 459-472.
157. Snow, R., Mutants of yeast sensitive to ultraviolet light, *J. Bacteriol.*, 94 (1967) 571-575.
158. \_\_\_\_\_, Recombination in ultraviolet-sensitive strains of *Saccharomyces cerevisiae*, *Mutation Res.*, 6 (1968) 409-418.
159. Speyer, J. F., Mutagenic DNA polymerase, *Biochem. Biophys. Res. Commun.*, 21 (1965) 6-8.
160. Suslova, N. G. and I. A. Zakharov, The gene-controlled radiation sensitivity of yeast. VII Identification of the genes for the X-ray sensitivity, *Genetika*, 6 (1970) 158-163.
161. Terisima, T. and L. J. Tolmach, Change in X-ray sensitivities of Hela cells during the division cycle, *Nature*, 190 (1961) 1210-1211.
162. Unrau, P., R. Wheatcroft, and B. S. Cox, The excision of pyrimidine dimers from DNA of ultraviolet irradiated yeast, *Molec. Gen. Genet.*, 113 (1971) 359-362.
163. von Borstel, R. C., K. T. Cain and C. M. Steinberg, Inheritance of spontaneous mutability in yeast, *Genetics*, 69 (1971) 17-27.
164. von Borstel, R. C. and P. J. Hastings, Mutagenic repair pathways in yeast, in A. Castellani (ed.), *Research in Photobiology*, Plenum Press, New York, 1976, pp. 683-687.

165. von Borstel, R. C., S.-K. Quah, C. M. Steinberg, F. Flury and D. J. C. Gottlieb, Mutants of yeast with enhanced spontaneous mutation rates, *Genetics Suppl.*, 73 (1973) 141-151.
166. Waters, R. and E. Moustacchi, The disappearance of ultraviolet induced pyrimidine dimers from the nuclear DNA of exponential and stationary phase cells of *Saccharomyces cerevisiae* following various post-irradiation treatments, *Biochim. Biophys. Acta*, 353 (1974) 407-419.
167. Waters, R. and E. Moustacchi, The fate of ultraviolet-induced pyrimidine dimers in the mitochondrial DNA of *Saccharomyces cerevisiae* following various post-irradiation cell treatments, *Biochim. Biophys. Acta*, 366 (1974) 241-250.
168. Whelan, W. L., E. Gocke and T. R. Manney, The *CAN1* locus of *Saccharomyces cerevisiae*: Fine-structure analysis and forward mutation rates, *Genetics*, 91 (1979) 35-51.
169. Wickner, R. B., Mutants of *Saccharomyces cerevisiae* that incorporate deoxythymidine-5'-monophosphate into deoxyribonucleic acid in vivo, *J. Bacteriol.*, 117 (1974) 252-260.
170. Wildenberg, J., The relation of mitotic recombination to DNA replication in yeast pedigrees, *Genetics*, 66 (1970) 291-304.
171. Williamson, D. H., Replication of the nuclear genome does not require concomitant protein synthesis in yeast, *Biochem. Biophys. Res. Commun.*, 52 (1973) 731-740.
172. Witkin, E. M., Radiation-induced mutations and their repair, *Science*, 152 (1966) 1345-1353.
173. \_\_\_\_\_, Mutation-proof and mutation-prone modes of survival in derivatives of *Escherichia coli* B differing in sensitivity to ultraviolet light, *Brookhaven Symp. Biol.*, 20 (1967) 17-55.
174. \_\_\_\_\_, The role of DNA repair and recombination in mutagenesis, *Proc. XII Int. Cong. Genet.*, 3 (1969) 225-245.
175. \_\_\_\_\_, Ultraviolet-induced mutation and DNA repair, *Ann Rev. Genet.*, 3 (1969) 525-552.
176. \_\_\_\_\_, Ultraviolet mutagenesis and inducible DNA repair in *Escherichia coli*, *Bacteriol. Rev.*, 40 (1976) 869-907.
177. Zakharov, I. A., T. N. Kozina, I. V. Fedorova, Effets des mutations vers la sensibilité au rayonnement ultraviolet chez la levure, *Mutation Res.*, 9 (1970) 31-39.

Table 1 YEAST MUTANTS SELECTED FOR HYPERSENSITIVITY TO MUTAGENS

Author, year [ref.]	Enrichment (prescreen)	Isolation (screen)	Genetic Loci <sup>a</sup>	Cumulative Total (New Loci)
Nakai and Matsumoto, 1967 [122]	UV UV	-- <sup>b</sup> UV	<u>rad1</u> <u>rad2,rad51</u>	1 3
Snow, 1967 [157, unpublished]	EMS	UV	<u>rad1, ..., 5, 10, 14</u>	8
Cox and Parry, 1968 [22]	EMS, UV	UV	<u>rad1, ..., 17, 19... 22, 50</u>	23
Resnick, 1969 [140]	NA	UV, X <sup>c</sup>	<u>rad1, 2, 18, 52, 53, xs3</u>	27
Moustacchi, 1969 [120]	UV	UV	<u>rad1</u>	27
Zakharov et al., 1970 [177]	UV	UV	<u>rad2, rad4, uvs2</u>	28
Suslova and Zakharov, 1970 [156]	-	X	<u>rad50, 51, 54, xrs2, xrs4</u>	31
Averbeck et al., 1970 [2]	MNNG	UV	<u>rad2, rad9, r<sub>1</sub><sup>s</sup></u>	32
Snow [unpublished] Mortimer [unpublished] Game and Mortimer, 1974 [42]	-	X	<u>rad5, 9, 18, 50, 52, ..., 57</u>	35
Prakash and Prakash, 1977 [135]	UV	MMS	<u>rad1, 4, 6, 52, 55, 57, mms1, ..., 22</u>	57
Ananthaswamy et al., 1978 [1]	MNNG	X	<u>rad5, 17, 50, 53, 54, + 15 new isolates</u>	61-72

<sup>a</sup>Standardized rad locus assignments are based upon inter-lab allelism tests [37, 38, 39]; rad50 and higher confer only ionizing radiation sensitivity; rad1, ..., rad49 have been reserved for those that confer only UV sensitivity or sensitivity to both UV and ionizing radiation; others have not been tested for complementation or assigned to rad loci; ellipses (...) refer to consecutive locus numbers implied by the series.

<sup>b</sup>Identified in a survey of auxotrophic yeast stocks

<sup>c</sup>Screened for sensitivity to UV or X-rays, or both.

Table 2 YEAST MUTANTS SELECTED FOR ALTERED MUTAGENESIS OR MITOTIC RECOMBINATION

Author, year [ref.]	Selection Phenotype	Genetic Loci
Lemontt, 1971 [93]	Decreased UV reversion of <u>arg4-17</u> (i.e. UV hypo-mutable)	<u>rev1</u> , <u>rev2</u> , <u>rev3</u>
Lemontt, 1973 [96], 1976 [100]	Decreased UV forward mutation of <u>CAN1</u> (UV hypo-mutable)	<u>umr1</u> , ..., <u>umr7</u>
Rodarte-Ramon and Mortimer, 1972 [147]	Decreased X-ray-induced gene conversion at <u>arg4</u> (X-ray hypo-rec)	<u>rec1</u> , ..., <u>rec5</u> , 2D11, 2C16
von Borstel <u>et al.</u> , 1973[165] Hastings <u>et al.</u> , 1976 [55]	Increased spontaneous reversion of <u>lys1-1</u> (mutator)	<u>mut1</u> , ..., <u>mut5</u> , <u>MUT6</u> , <u>mut9</u> , <u>mut10</u>
Maloney and Fogel, 1976 [108]	Increased spontaneous gene conversion at <u>arg4</u> (hyper-rec)	-
Golin and Esposito, 1977 [46]	Increased spontaneous mutation of <u>CAN1</u> (mutator)	<u>rem1</u>
Quah <u>et al.</u> , 1978 [138]	Decreased spontaneous reversion of <u>lys1-1</u> (antimutator)	-

Table 3 ALLELE-SPECIFIC CONTROL OF UV MUTAGENESIS BY REV GENES<sup>a</sup>

Type	Allele	Codon			UV Revertibility			
		Mutant	Normal	Position	REV	rev1	rev2	rev3
ochre	<u>cycl-9</u>	UAA	GAA	2	+	-	-	-
	<u>-2</u>	UAA	CAA	21	+	-	+	-
	<u>-72</u>	UAA	GAA	66	+	-	+	-
amber	<u>cycl-179</u>	UAG	AAG	9	+	-	+	-
	<u>-84</u>	UAG	UGG	64	+	-	+	-
	<u>-76</u>	UAG	GAG	71	+	-	+	-
Initiation	<u>cycl-131</u>	GUG	AUG	-1	+	+	+	(-)
	<u>-133</u>	AGG	AUG	-1	+	-	+	-
	<u>-13</u>	AUP <sub>y</sub>	AUG	-1	+	-	-	-
	<u>-51</u>	CUG	AUG	-1	+	-	-	-
Frame-shift	<u>cycl-183</u>	+A	AAA	10	+	+	+	-
	<u>-239</u>	-G	AAG	4	+	+	+	-
	<u>-331</u>	-A	GAA	2	+	+	-	-
Proline missense	<u>cycl-115</u>	CCPy	CUP <sub>y</sub>	14	+	+	+	(-)
	<u>-6</u>	CCU	GCU	12	+	-	-	-

<sup>a</sup>Data from Lawrence and Christensen [87, 88, 89, 91]