

ANL/NDM--66

DE82 019477

ANL/NDM-66

**FAST-NEUTRON SCATTERING CROSS SECTIONS
OF ELEMENTAL SILVER***

by

A. B. Smith and P. T. Guenther

May 1982

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

***This work supported by the U. S. Department of Energy.**

**Applied Physics Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439
USA**

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

NUCLEAR DATA AND MEASUREMENTS SERIES

The Nuclear Data and Measurements Series presents results of studies in the field of microscopic nuclear data. The primary objective is the dissemination of information in the comprehensive form required for nuclear technology applications. This Series is devoted to: a) measured microscopic nuclear parameters, b) experimental techniques and facilities employed in measurements, c) the analysis, correlation and interpretation of nuclear data, and d) the evaluation of nuclear data. Contributions to this Series are reviewed to assure technical competence and, unless otherwise stated, the contents can be formally referenced. This Series does not supplant formal journal publication but it does provide the more extensive information required for technological applications (e.g., tabulated numerical data) in a timely manner.

OTHER ISSUES IN THE ANL/NDM SERIES ARE:

- ANL/NDM-1 Cobalt Fast Neutron Cross Sections--Measurement and Evaluation by P. T. Guenther, P. A. Moldauer, A. B. Smith, D. L. Smith and J. F. Whalen, July 1973.
- ANL/NDM-2 Prompt Air-Scattering Corrections for a Fast-Neutron Fission Detector: $E_n \leq 5$ MeV by Donald L. Smith, September 1973.
- ANL/NDM-3 Neutron Scattering from Titanium; Compound and Direct Effects by E. Barnard, J. deVilliers, P. Moldauer, D. Reitmann, A. Smith and J. Whalen, October 1973.
- ANL/NDM-4 ^{90}Zr and ^{92}Zr ; Neutron Total and Scattering Cross Sections by P. Guenther, A. Smith and J. Whalen, July 1974.
- ANL/NDM-5 Delayed Neutron Data - Review and Evaluation by Samson A. Cox, April 1974.
- ANL/NDM-6 Evaluated Neutronic Cross Section File for Niobium by R. Howerton, Lawrence Livermore Laboratory and A. Smith, P. Guenther and J. Whalen, Argonne National Laboratory, May 1974.
- ANL/NDM-7 Neutron Total and Scattering Cross Sections of Some Even Isotopes of Molybdenum and the Optical Model by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1974.
- ANL/NDM-8 Fast Neutron Capture and Activation Cross Sections of Niobium Isotopes by W. P. Poenitz, May 1974.
- ANL/NDM-9 Method of Neutron Activation Cross Section Measurement for $E_n = 5.5\text{--}10$ MeV Using the $\text{D}(d,n)\text{He-3}$ Reaction as a Neutron Source by D. L. Smith and J. W. Meadows, August 1974.
- ANL/NDM-10 Cross Sections for (n,p) Reactions on ^{27}Al , $^{46,47,48}\text{Ti}$, $^{54,56}\text{Fe}$, ^{58}Ni , ^{59}Co and ^{64}Zn from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, January 1975.
- ANL/NDM-11 Measured and Evaluated Fast Neutron Cross Sections of Elemental Nickel by P. Guenther, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, July 1975.
- ANL/NDM-12 A Spectrometer for the Investigation of Gamma Radiation Produced by Neutron-Induced Reactions by Donald L. Smith, April 1975.
- ANL/NDM-13 Response of Several Threshold Reactions in Reference Fission Neutron Fields by Donald L. Smith and James W. Meadows, June 1975.
- ANL/NDM-14 Cross Sections for the $^{66}\text{Zn}(n,p)^{66}\text{Cu}$, $^{113}\text{In}(n,n')^{113\text{m}}\text{In}$ and $^{115}\text{In}(n,n')^{115\text{m}}\text{In}$ Reactions from Near Threshold to 10 MeV by Donald L. Smith and James W. Meadows, July 1975.

- ANL/NDM-15 Radiative Capture of Fast Neutrons in ^{165}Ho and ^{181}Ta by W. P. Poenitz, June 1975.
- ANL/NDM-16 Fast Neutron Excitation of the Ground-State Rotational Band of ^{238}U by P. Guenther, D. Havel and A. Smith, September 1975.
- ANL/NDM-17 Sample-Size Effects in Fast-Neutron Gamma-Ray Production Measurements: Solid-Cylinder Samples by Donald L. Smith, September 1975.
- ANL/NDM-18 The Delayed Neutron Yield of ^{238}U and ^{241}Pu by J. W. Meadows January 1976.
- ANL/NDM-19 A Remark on the Prompt-Fission-Neutron Spectrum of ^{252}Cf by P. Guenther, D. Havel, R. Sjoblom and A. Smith, March 1976.
- ANL/NDM-20 Fast-Neutron-Gamma-Ray Production from Elemental Iron: $E_n \lesssim 2$ MeV by Donald L. Smith, May 1976.
- ANL/NDM-21 Note on the Experimental Determination of the Relative Fast-Neutron Sensitivity of a Hydrogenous Scintillator by A. Smith, P. Guenther and R. Sjoblom, June 1976.
- ANL/NDM-22 Note on Neutron Scattering and the Optical Model Near $A=208$ by P. Guenther, D. Havel and A. Smith, September 1976.
- ANL/NDM-23 Remarks Concerning the Accurate Measurement of Differential Cross Sections for Threshold Reactions Used in Fast-Neutron Dosimetry for Fission Reactors by Donald L. Smith, December 1976.
- ANL/NDM-24 Fast Neutron Cross Sections of Vanadium and an Evaluated Neutronic File by P. Guenther, D. Havel, R. Howerton, F. Mann, D. Smith, A. Smith and J. Whalen, May 1977.
- ANL/NDM-25 Determination of the Energy Scale for Neutron Cross Section Measurements Employing a Monoenergetic Accelerator by J. W. Meadows, January 1977.
- ANL/NDM-26 Evaluation of the $^{115}\text{In}(n,n')^{115m}\text{In}$ Reaction for the ENDF/B-V Dosimetry File by Donald L. Smith, December 1976.
- ANL/NDM-27 Evaluated (n,p) Cross Sections of ^{46}Ti , ^{47}Ti and ^{48}Ti by C. Philis and O. Bersillon, Bruyeres-le-Chatel, France and D. Smith and A. Smith, Argonne National Laboratory, January 1977.
- ANL/NDM-28 Titanium-II: An Evaluated Nuclear Data File by C. Philis, Centre d'Etudes de Bruyeres-le-Chatel, R. Howerton, Lawrence Livermore Laboratory and A. B. Smith, Argonne National Laboratory, June 1977.
- ANL/NDM-29 Note on the 250 keV Resonance in the Total Neutron Cross Section of ^6Li by A. B. Smith, P. Guenther, D. Havel and J. F. Whalen, June 1977.

- ANL/NDM-15 Radiative Capture of Fast Neutrons in ^{165}Ho and ^{181}Ta by W. P. Poenitz, June 1975.
- ANL/NDM-16 Fast Neutron Excitation of the Ground-State Rotational Band of ^{238}U by P. Guenther, D. Havel and A. Smith, September 1975.
- ANL/NDM-17 Sample-Size Effects in Fast-Neutron Gamma-Ray Production Measurements: Solid-Cylinder Samples by Donald L. Smith, September 1975.
- ANL/NDM-18 The Delayed Neutron Yield of ^{238}U and ^{241}Pu by J. W. Meadows January 1976.
- ANL/NDM-19 A Remark on the Prompt-Fission-Neutron Spectrum of ^{252}Cf by P. Guenther, D. Havel, R. Sjoblom and A. Smith, March 1976.
- ANL/NDM-20 Fast-Neutron-Gamma-Ray Production from Elemental Iron: $E_n \lesssim 2$ MeV by Donald L. Smith, May 1976.
- ANL/NDM-21 Note on the Experimental Determination of the Relative Fast-Neutron Sensitivity of a Hydrogenous Scintillator by A. Smith, P. Guenther and R. Sjoblom, June 1976.
- ANL/NDM-22 Note on Neutron Scattering and the Optical Model Near $A=208$ by P. Guenther, D. Havel and A. Smith, September 1976.
- ANL/NDM-23 Remarks Concerning the Accurate Measurement of Differential Cross Sections for Threshold Reactions Used in Fast-Neutron Dosimetry for Fission Reactors by Donald L. Smith, December 1976.
- ANL/NDM-24 Fast Neutron Cross Sections of Vanadium and an Evaluated Neutronic File by P. Guenther, D. Havel, R. Howerton, F. Mann, D. Smith, A. Smith and J. Whalen, May 1977.
- ANL/NDM-25 Determination of the Energy Scale for Neutron Cross Section Measurements Employing a Monoenergetic Accelerator by J. W. Meadows, January 1977.
- ANL/NDM-26 Evaluation of the $\text{In-}^{115}(\text{n},\text{n}')\text{In-}^{115\text{m}}$ Reaction for the ENDF/B-V Dosimetry File by Donald L. Smith, December 1976.
- ANL/NDM-27 Evaluated (n,p) Cross Sections of ^{46}Ti , ^{47}Ti and ^{48}Ti by C. Philis and O. Bersillon, Bruyeres-le-Chatel, France and D. Smith and A. Smith, Argonne National Laboratory, January 1977.
- ANL/NDM-28 Titanium-II: An Evaluated Nuclear Data File by C. Philis, Centre d'Etudes de Bruyères-le-Châtel, R. Howerton, Lawrence Livermore Laboratory and A. B. Smith, Argonne National Laboratory, June 1977.
- ANL/NDM-29 Note on the 250 keV Resonance in the Total Neutron Cross Section of ^6Li by A. B. Smith, P. Guenther, D. Havel and J. F. Whalen, June 1977.

- ANL/NDM-30 Analysis of the Sensitivity of Spectrum-Average Cross Sections to Individual Characteristics of Differential Excitation Functions by Donald L. Smith, March 1977.
- ANL/NDM-31 Titanium-I: Fast Neutron Cross Section Measurements by P. Guenther, D. Havel, A. Smith and J. Whalen, May 1977.
- ANL/NDM-32 Evaluated Fast Neutron Cross Section of Uranium-238 by W. Poenitz, E. Pennington, and A. B. Smith, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, October 1977.
- ANL/NDM-33 Comments on the Energy-Averaged Total Neutron Cross Sections of Structural Materials by A. B. Smith and J. F. Whalen, June 1977.
- ANL/NDM-34 Graphical Representation of Neutron Differential Cross Section Data for Reactor Dosimetry Applications by Donald L. Smith, June 1977.
- ANL/NDM-35 Evaluated Nuclear Data File of Th-232 by J. Meadows, W. Poenitz, A. Smith, D. Smith and J. Whalen, Argonne National Laboratory and R. Howerton, Lawrence Livermore Laboratory, February 1978.
- ANL/NDM-36 Absolute Measurements of the $^{233}\text{U}(n,f)$ Cross Section Between 0.13 and 8.0 MeV by W. P. Poenitz, April 1978.
- ANL/NDM-37 Neutron Inelastic Scattering Studies for Lead-204 by D. L. Smith and J. W. Meadows, December 1977.
- ANL/NDM-38 The Alpha and Spontaneous Fission Half-Lives of ^{242}Pu by J. W. Meadows, December 1977.
- ANL/NDM-39 The Fission Cross Section of ^{239}Pu Relative to ^{235}U from 0.1 to 10 MeV by J. W. Meadows, March 1978.
- ANL/NDM-40 Statistical Theory of Neutron Nuclear Reactions by P. A. Moldauer, February 1978.
- ANL/NDM-41 Energy-Averaged Neutron Cross Sections of Fast-Reactor Structural Materials by A. Smith, R. McKnight and D. Smith, February 1978.
- ANL/NDM-42 Fast Neutron Radiative Capture Cross Section of ^{232}Th by W. P. Poenitz and D. L. Smith, March 1978.
- ANL/NDM-43 Neutron Scattering from ^{12}C in the Few-MeV Region by A. Smith, R. Holt and J. Whalen, September 1978.
- ANL/NDM-44 The Interaction of Fast Neutrons with ^{60}Ni by A. Smith, P. Guenther, D. Smith and J. Whalen, January 1979.
- ANL/NDM-45 Evaluation of $^{235}\text{U}(n,f)$ Between 100 KeV and 20 MeV by W. P. Poenitz, July 1979.

- ANL/NDM-46 Fast-Neutron Total and Scattering Cross Sections of ^{107}Ag in the MeV Region by A. Smith, P. Guenther, G. Winkler and J. Whalen, January 1979.
- ANL/NDM-47 Scattering of MeV Neutrons from Elemental Iron by A. Smith and P. Guenther, March 1979.
- ANL/NDM-48 ^{235}U Fission Mass and Counting Comparison and Standardization by W. P. Poenitz, J. W. Meadows and R. J. Armani, May 1979.
- ANL/NDM-49 Some Comments on Resolution and the Analysis and Interpretation of Experimental Results from Differential Neutron Measurements by Donald L. Smith, November 1979.
- ANL/NDM-50 Prompt-Fission-Neutron Spectra of ^{233}U , ^{235}U , ^{239}Pu and ^{240}Pu Relative to that of ^{252}Cf by A. Smith, P. Guenther, G. Winkler and R. McKnight, September 1979.
- ANL/NDM-51 Measured and Evaluated Neutron Cross Sections of Elemental Bismuth by A. Smith, P. Guenther, D. Smith and J. Whalen, April 1980.
- ANL/NDM-52 Neutron Total and Scattering Cross Sections of ^6Li in the Few MeV Region by P. Guenther, A. Smith and J. Whalen, February 1980.
- ANL/NDM-53 Neutron Source Investigations in Support of the Cross Section Program at the Argonne Fast-Neutron Generator by James W. Meadows and Donald L. Smith, May 1980.
- ANL/NDM-54 The Nonelastic-Scattering Cross Sections of Elemental Nickel by A. B. Smith, P. T. Guenther and J. F. Whalen, June 1980.
- ANL/NDM-55 Thermal Neutron Calibration of a Tritium Extraction Facility using the $^6\text{Li}(n,t)^4\text{He}/^{197}\text{Au}(n,\gamma)^{198}\text{Au}$ Cross Section Ratio for Standardization by M. M. Bretscher and D. L. Smith, August 1980.
- ANL/NDM-56 Fast-Neutron Interactions with ^{182}W , ^{184}W and ^{186}W by P. T. Guenther, A. B. Smith and J. F. Whalen, December 1980.
- ANL/NDM-57 The Total, Elastic- and Inelastic-Scattering Fast-Neutron Cross Sections of Natural Chromium, Peter T. Guenther, Alan B. Smith and James F. Whalen, January 1981.
- ANL/NDM-58 Review of Measurement Techniques for the Neutron Capture Process by W. P. Poenitz, August 1981.
- ANL/NDM-59 Review of the Importance of the Neutron Capture Process in Fission Reactors, Wolfgang P. Poenitz, July 1981.
- ANL/NDM-60 Neutron Capture Activation Cross Sections of ^{94}Zr , ^{96}Zr , $^{98,100}\text{Mo}$, and $^{110,114,116}\text{Cd}$ at Thermal and 30 keV Energy, John M. Wyrick and Wolfgang P. Poenitz, (to be published).

- ANL/NDM-61 Fast-neutron Total and Scattering Cross Sections of ^{58}Ni by Carl Budtz-Jørgensen, Peter T. Guenther, Alan B. Smith and James F. Whalen, September 1981.
- ANL/NDM-62 Covariance Matrices and Applications to the Field of Nuclear Data, by Donald L. Smith, November 1981.
- ANL/NDM-63 On Neutron Inelastic-Scattering Cross Sections of ^{232}Th , ^{233}U , ^{235}U , ^{238}U , ^{239}U , and ^{239}Pu and ^{240}Pu by Alan B. Smith and Peter T. Guenther, January 1982.
- ANL/NDM-64 The Fission Fragment Angular Distributions and Total Kinetic Energies for $^{235}\text{U}(n,f)$ from .18 to 8.83 MeV by James W. Meadows, and Carl Budtz-Jørgensen, (to be published).
- ANL/NDM-65 Note on the Elastic Scattering of Several MeV Neutrons from Elemental Calcium by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-66 Fast-neutron Scattering Cross Sections of Elemental Silver by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-67 Non-evaluation Applications for Covariance Matrices by Donald L. Smith, (to be published).
- ANL/NDM-68 Fast-neutron Total and Scattering Cross Sections of ^{103}Rh by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-69 Fast-neutron Scattering Cross Sections of Elemental Zirconium by Alan B. Smith and Peter T. Guenther (to be published).
- ANL/NDM-70 Fast-neutron Total and Scattering Cross Sections of Niobium by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).
- ANL/NDM-71 Fast-neutron Total and Scattering Cross Sections of Elemental Palladium by Alan B. Smith, Peter T. Guenther and James F. Whalen (to be published).

TABLE OF CONTENTS

	<u>Page</u>
Abstract	ix
I. Introduction.	1
II. Outline of Experimental Methods	1
III. Experimental Results.	2
IV. Interpretation and Discussion	5
V. Comparisons with ENDF/B-V	9
VI. Summary Comment	9
Acknowledgements	9
References	11

FAST-NEUTRON SCATTERING CROSS SECTIONS
OF ELEMENTAL SILVER

by

A. B. Smith and P. T. Guenther
Applied Physics Division
Argonne National Laboratory
9700 South Cass Avenue
Argonne, Illinois 60439, USA

ABSTRACT

Differential neutron elastic- and inelastic-scattering cross sections of elemental silver are measured from 1.5 to 4.0 MeV at intervals of $\lesssim 200$ keV and at 10 to 20 scattering angles distributed between 20 and 160 deg. Inelastically-scattered neutron groups are observed corresponding to the excitation of levels at; 328 ± 13 , 419 ± 50 , 748 ± 25 , 908 ± 26 , 1150 ± 38 , 1286 ± 25 , 1507 ± 20 , 1623 ± 30 , 1835 ± 20 and 1944 ± 26 keV. The experimental results are used to derive an optical-statistical model that provides a good description of the observed cross sections. The measured values are compared with corresponding quantities given in ENDF/B-V.

I. INTRODUCTION

This is the second of a series of reports dealing with the fast-neutron properties of fission-product nuclei in the region of light-mass fission product yields.¹ The general objective is the derivation of a model suitable for the calculation of unmeasurable neutronic properties of the light-mass fission products. The primary applied relevance is to Fast-Breeder Reactor (FBR) systems.

Elemental silver consists of two isotopes; ^{107}Ag (51%) and ^{109}Ag (49%).² Their fission-product yields can be significant. For example, in fission-neutron-induced fission of ^{239}Pu , the yields are $\approx 3.7\%$ (^{107}Ag) and $\approx 1.6\%$ (^{109}Ag). Both isotopes are in the first island of isomerism with p-1/2 ground states and low-lying (≈ 100 keV) 7/2 and 9/2+ isomeric levels.³ The primary goal of the present measurements was the quantitative determination of the differential elastic-scattering cross sections, and from them the derivation of an appropriate optical-statistical model (OM). The measurements extended over a region where the model is sensitive to compound-nucleus (CN) contributions. A secondary goal was the determination of differential inelastic-scattering cross sections. The latter results are primarily of applied interest as the intentionally broad incident-energy spreads employed in the measurements precluded detailed resolution of excited-level structure.

Subsequent portions of this report deal with: i) a very brief outline of the experimental methods (Sec. II), ii) the experimental results (Sec. III), the derivation of an OM (Sec. IV), iii) and some comparisons with ENDF/B-V (Sec. V).

II. OUTLINE OF EXPERIMENTAL METHODS

All the measurements were made at the Argonne National Laboratory Fast-neutron Generator using the ten-angle time-of-flight apparatus.⁴ The neutron source was the $^7\text{Li}(p,n)^7\text{Be}$ reaction.⁵ The measurement sample was a cylinder 2 cm in diameter and 2 cm long, fabricated from high purity elemental silver. The average incident-neutron energies were known to ≈ 10 keV and the incident-neutron energy spreads were ≈ 70 to 100 keV. The ten scattered-neutron flight paths were ≈ 5.4 m long. Ten proton-recoil scintillation detectors were used to detect the scattered neutrons and an eleventh detector monitored the source intensity. The relative energy-dependent responses of the detectors were determined by observation of the fission neutrons emitted by the spontaneous fission of ^{252}Cf .⁶ These relative responses were normalized absolutely relative to the neutron total cross sections of carbon⁷ as described in Ref. 8. The resulting differential-scattering cross sections were corrected for angular-resolution, incident-neutron-attenuation and multiple-event effects as described in Ref. 9. The statistical uncertainties of the measured differential elastic-scattering cross sections were $\lesssim 2\%$ except near the extreme minima of the distributions where they were larger. The statistical uncertainties of the inelastic-scattering results varied from $\approx 10\%$ to 30% depending upon the particular inelastic-neutron group. Uncertainties associated with detector-calibration procedures were $\lesssim 3\%$. Neutron scattering angles were known to $\lesssim 0.75$ degree. Correction procedures introduced an

additional $\lesssim 1\%$ uncertainty, again excepting the extreme minima of the elastic distributions where they were larger. Thus the overall uncertainties for differential-elastic scattering cross sections were generally $\lesssim 5\%$, and for differential-inelastic scattering, $\approx 10\%$ to 30% .

III. EXPERIMENTAL RESULTS^a

A. Neutron-Elastic-Scattering Cross Sections

The differential-elastic-scattering measurements were made at incident energies extending from 1.5 to 4.0 MeV. Below 3.0 MeV the measurements were made at intervals of ≈ 50 keV (i.e., at intervals smaller than the incident-energy resolution) and at ten scattering angles distributed between ≈ 20 and 160 degrees. The above cited broad resolutions were intentionally used in order to enhance the statistical accuracies and to average possible cross-section fluctuations. These broad resolutions precluded the separation of the elastic- and first-two-inelastic-scattered neutron groups corresponding to similar levels in ^{107}Ag and ^{109}Ag ($E_x \equiv 0.0$, ≈ 0.09 and ≈ 0.125 keV). Thus, all the "elastic-scattering" results obtained in the present work are inclusive of two small inelastic-scattering components. The latter are dealt with in the interpretation as described below. The possible effect of fluctuations was further reduced by averaging differential cross sections obtained at adjacent energies up to incident energies of 3.0 MeV. Above 3.0 MeV the measurements were made at 200 keV intervals and at twenty scattering angles. At these higher energies the fluctuations should be negligible but the angular dependences of the cross sections are strong thus the additional angular detail was desirable.

The measured "elastic-scattering" results are illustrated in Fig. 1. In order to obtain the angle-integrated "elastic-scattering" cross sections, the measured values were least-square fitted with 6th-order Legendre-Polynomial series. The results of this fitting procedure are illustrated by the curves of Fig. 1 and the corresponding angle-integrated cross-section values are shown in Fig. 2. Of course, these angle-integrated results also include the two small inelastic-scattering components noted above. The uncertainties associated with the angle-integrated results are $\lesssim 5\%$. The present "elastic-scattering" values reasonably extrapolate to the lower-energy results of Ref. 10, as illustrated in Fig. 2. They are also consistent with the ^{107}Ag isotopic results of Ref. 11. The latter measurements fully resolved the elastic scattering and consequently the extreme minima of the distributions are slightly smaller than those of the present work at the lower energies where compound-nucleus cross sections are large. In addition, the ^{107}Ag and elemental level structures differ; resulting in different compound elastic-scattering (CE) contributions and thus in variations between elemental and isotopic scattering cross sections.

B. Neutron Inelastic-Scattering Cross Sections

The primary objective of the present measurement was the determination of energy-averaged elastic scattering cross sections and the energy resolution

a) Numerical values for all experimental results reported herein have been transmitted to the National Nuclear Data Center, Brookhaven National Laboratory.




Fig. 1. Differential-"Elastic-Scattering" Cross Sections of Elemental Silver. The present measured values are indicated by data symbols. Curves denote the results of fitting Legendre-Polynomial series to the measured values. Angle is expressed in the laboratory system.




Fig. 2. Neutron Total and "Elastic-Scattering" Cross Sections of Elemental Silver. The present experimental results are indicated by \square , those of Ref. 10 by \circ , those of Ref. 11 (^{107}Ag) by \diamond , and those of Ref. 12 by \times . Curves indicate the results of model calculations as described in the text.

was chosen accordingly. As a consequence of this and of the complex level structure of elemental silver the inelastic scattering cross sections were not well resolved. However, ten distinct inelastically-scattered neutron groups were observed that could be generally attributed to the excitation of two or more levels in ^{107}Ag and/or ^{109}Ag . The correlation of the observed neutron groups with reported levels in the two isotopes is outlined in Table 1.² The excitation energies derived from the measurements are averages of a number of observed values with the respective uncertainties defined as the RMS deviations from the simple averages.

The angle-integrated cross sections were determined by least-square fitting the measured differential values with low-order Legendre-Polynomial series (e.g., $\sum P_2$ as the observed differential-cross-section anisotropics were small). The uncertainties associated with the angle-integrated inelastic values vary from $\approx 10\%$ to 30% depending upon the particular measurements. The contributions to the uncertainties were as outlined above with an additional factor due to necessary corrections for the second-neutron group from the source reaction. The resulting inelastic-cross-section values are summarized in Fig. 3. As noted above, the measurements intentionally did not resolve the first two inelastic-neutron groups ($E_x \approx 90 \text{ keV}(7/2+)$ and $\approx 130 \text{ keV}(9/2+)$). Excitations of these two levels are indicated in Fig. 3, based upon previously reported work from this Laboratory.^{10,11} Cross sections for the excitation of the 328 keV ($3/2-$) level were determined at lower incident energies. As the energy increased this level was observed as a composite excitation with the 419 keV ($5/2-$) level due to the broad incident-neutron resolutions employed in the measurements. Furthermore, excitations in the region of 400-500 keV were particularly complicated by the presence of the elastically-scattered second neutron group from the source reaction. The appropriate corrections for this second-group perturbation considerably increased the uncertainty in this case. Cross sections corresponding to the excitation of a 748 keV level were attributed to contributions from ^{107}Ag and ^{109}Ag as outlined in Table 1. Cross sections for the remaining seven groups also appear to be due to levels in both ^{107}Ag and ^{109}Ag (see Table 1). These correlations are complex and, for $E_x \gtrsim 1300 \text{ keV}$, the reported levels of the two isotopes become increasingly uncertain.² Generally the present cross sections are consistent with the lower-energy elemental results of Ref. 10 and with the ^{107}Ag isotopic results of Ref. 11 where ^{109}Ag contributions do not significantly distort the comparisons. Moreover, the sum of the "eye-guides" of Fig. 3 are in good agreement with the non-elastic cross sections implied by the present elastic-scattering results and the neutron total cross sections of Ref. 12.

IV. INTERPRETATION AND DISCUSSION

It was assumed that the present results could be reasonably described by a spherical optical-statistical model (OM).^{13,14} The low-lying levels of the two isotopes are similar but those of ^{109}Ag are better known as the energy increases.² Therefore this interpretation was based upon the discrete levels of ^{109}Ag , as given in Ref. 2, to excitations of 1260 keV. Higher-energy levels were represented using the statistical parameterization of Gilbert and Cameron.¹⁵ The elemental mass was defined as $A \equiv 108$. The




Fig. 3. Neutron Inelastic-Scattering Cross Sections of Elemental Silver. Measured values are indicated by data symbols as follows; \circ = present work, \square = Ref. 10, and \times = Ref. 11. Heavy curves are "eyeguides", the light curves indicate the results of calculations as discussed in the text. Observed excitation energies are given in keV.

calculations used the OM code ABAREX with resonance fluctuation and correlation corrections as described by Moldauer.^{16,17} The OM parameters were derived by concurrently chi-square fitting all of the measured differential-elastic-scattering cross sections of Fig. 1. The fitting procedure combined the first two inelastic-neutron groups ($E_x \approx 0.09$ and 0.125 keV) with the elastic-scattering component in correspondence with the measured values. The fitting procedure simultaneously considered the six parameters; real and imaginary strengths, radii and diffusenesses. The resulting OM parameters are given in Table 2. These parameters are relatively conventional. The real strength (measured in integral-per-nucleon J_v/A or Vr^2) is similar to that reported in the literature.^{11,18} The imaginary strength (J_w/A or W_a) tends to be somewhat larger than that frequently reported.

The parameters of Table 2 provide a generally good description of the measured differential-"elastic-scattering" cross sections as illustrated in Fig. 4. There are small systematic differences between the measured and calculated values near the extreme minima of the distributions from ≈ 2 to 3 MeV. This is a region sensitive to compound-elastic-scattering (CE) which here is largely governed by channel competition from the statistically-based levels. The latter are not well defined and may vary from nucleus to nucleus, particularly at these low energies. The measured angle-integrated elastic-scattering cross sections are described well by the calculations as illustrated in Fig. 2. The neutron total cross sections of Ref. 12 are also described well by the calculations. At ≈ 4.0 MeV the elemental total cross sections of Ref. 12 are $\approx 2\%$ to 3% larger than the ^{107}Ag values of Ref. 11. Calculations indicate that the nuclear size effect accounts for $\approx 1\%$ of this difference, leaving a $\approx 1\%$ to 2% difference which is consistent with the respective experimental uncertainties.

Calculated inelastic-scattering cross sections are compared with measured values in Fig. 3. The agreement is good at the lower energies where the excitation of discrete levels governs the processes. However, as the energy increases and the channel competition becomes dominated by statistical level distributions, the calculated values become markedly smaller than the measured quantities; particularly for the excitations of the 328 keV ($3/2^-$) and 419 keV ($5/2^-$) levels. This suggests that the statistical level parameters result in too much channel competition. The discrepancy can be reduced by adjusting the level parameters without a large effect on the elastic-scattering process as the CE component is already small. Such pragmatic parameter adjustment was not pursued. It has been suggested that both silver isotopes are vibrators¹⁹ with the first $3/2^-$ and $5/2^-$ levels coupled to the ground state in a direct-vibrational excitation. The magnitude of such a direct-vibrational excitation was estimated using a coupled-channels calculation assuming $\beta_2 = 0.15$.²⁰ The vibrational coupling results ("V") are compared with the spherical results ("S") in Fig. 3. The enhancement of the inelastic cross sections due to the direct process is significant and will increase with β_2 . However, all the calculated inelastic-scattering cross sections remain smaller than the measured values. The introduction of vibrational coupling will effect the choice of OM parameters to some extent at the present energies. Comprehensive fitting, using the vibrational model was not attempted since the spirit of the interpretation was a simple spherical OM. Comprehensive fitting with a




Fig. 4. Comparison of Measured and Calculated Differential-Elastic-Scattering Cross Sections of Elemental Silver. The measured values are indicated by data symbols and the results of model calculations by curves.

vibrational model is costly in computing time, and the major calculational uncertainty would probably remain the statistical-level parameters.

V. COMPARISONS WITH ENDF/B-V

The ENDF/B-V evaluated data file ²¹ contains isotopic ¹⁰⁷Ag and ¹⁰⁹Ag data sets. These were combined to obtain an elemental file for comparison with the present measurements. Figure 5 compares measured and evaluated neutron total²¹ and elastic-scattering cross sections (small corrections have been made for inelastic contributions to the measured values as necessary). Near 4.0 MeV the agreement is good for both cross sections. However, as the energy decreases the evaluated results become increasingly larger than the measured quantities, the difference amounting to $\approx 8\%$ at several hundred keV. That is a significant discrepancy, particularly with respect to the total cross section which forms the envelope of an evaluated data set. Moreover, the discrepancy is largest at the low energies where the neutron capture cross section is important. The non-elastic-scattering cross section derived from the present measurements and the total cross section of Ref. 12 is in reasonably good agreement with the evaluated total inelastic-scattering cross section. There is a similar reasonable agreement at low energies between the evaluated individual inelastic-excitation cross sections and the measured values of Refs. 10 and 11. The agreement may be fortuitous in view of the above noted discrepancies between measured and evaluated neutron total and elastic-scattering cross sections.

VI. SUMMARY COMMENT

The present measurements give new and detailed definition to neutron elastic scattering from elemental silver and improve the knowledge of inelastic scattering over the few-MeV region. Combined with recently measured neutron total cross sections and previously reported scattering cross sections from this Laboratory¹⁰⁻¹², a reasonably comprehensive experimental understanding of the neutron interaction with elemental silver is obtained from a few-hundred keV to above 4 MeV. The measured "elastic-scattering" results form a data base from which a spherical OM was derived. That OM is consistent with a general OM applicable to the light-mass fission products to be reported elsewhere.²² In addition, details of the experimental interpretation suggest that direct-vibrational processes may play a part in the neutron interaction with silver in the several-MeV region. The present experimental results are reasonably consistent with the evaluated quantities of ENDF/B-V²¹ at energies of 4 to 5 MeV. At lower energies there are discrepancies between measured and evaluated results with the measured neutron total¹² and elastic-scattering cross sections becoming increasingly smaller than the evaluated quantities as the energy decreases.

ACKNOWLEDGEMENTS

The authors are indebted to Drs. W. Poenitz and P. Moldauer for helpful discussions, provision of computational programs and making available pre-publication data.




Fig. 5. Comparison of Measured and Evaluated Neutron Total and Elastic Scattering Cross Sections of Elemental Silver. The light curves were deduced from ENDF/B-V.²¹ The heavy curves are eyeguides constructed through the neutron total cross sections of Ref. 12 and the elastic-scattering results of the present work.

REFERENCES

1. A. Smith, P. Guenther and J. Whalen, Argonne National Laboratory Report, ANL/NDM-69 (1982).
2. Table of Isotopes, 7th Edition, C. M. Lederer and V. S. Shirley Eds., John Wiley and Sons, Inc., New York (1978).
3. M. Goeppert-Mayer and J. Hans D. Jensen, Elementary Theory of Nuclear Shell Structure, John Wiley and Sons, Inc., New York (1955).
4. A. Smith, P. Guenther, R. Larsen, C. Nelson, P. Walker and J. Whalen, Nucl. Instr. and Methods, 50 277 (1967).
5. Proc. of IAEA Consultants Meeting on Neutron Source Properties, INDC (NDS)-114/9T, IAEA Press (1980).
6. A. Smith, P. Guenther and R. Sjoblum, Nucl. Instr. and Methods, 140 397 (1977).
7. A. Smith, P. Guenther, R. Holt and J. Whalen, Argonne National Laboratory Report, ANL/NDM-43 (1978); see also Nucl. Sci. and Eng., 70 281 (1979).
8. A. B. Smith and P. T. Guenther, Argonne National Laboratory Report, ANL/NDM-63 (1982).
9. P. T. Guenther, Elastic and Inelastic Neutron Scattering from the Even Isotopes of Tungsten, Univ. of Illinois Thesis (1977).
10. W. Vonach and A. Smith, Nucl. Phys., 78 389 (1966).
11. A. Smith, P. Guenther, G. Winkler and J. Whalen, Nucl. Phys. A332 297 (1979).
12. W. P. Poenitz, to be published.
13. P. Hodgson, Nuclear Reactions and Nuclear Structure, Clarendon Press, Oxford (1971).
14. W. Hauser and H. Feshbach, Phys. Rev., 87 366 (1952).
15. A. Gilbert and A. Cameron, Can. Jour. of Physics, 43 1446 (1965).
16. ABAREX, a spherical optical-model code, P. A. Moldauer, priv. comm. (1982).
17. P. A. Moldauer, Phys. Rev., C11 426 (1973); also priv. comm. (1982).
18. J. Rapaport, V. Kulkorni and R. Finley, Nucl. Phys., A330 15 (1979).
19. Ch. Lagrange, Proc. of Conf. on Nuclear Data Evaluation Methods and Procedures, Brookhaven National Laboratory Report, BNL-NCS-51363 (1980).

REFERENCES (Contd.)

20. ECIS, a coupled-channels code, J. Raynal, unpublished (1981).
21. Evaluated Nuclear Data File-B, Version-V, Brookhaven National Laboratory Report, ENDF-201 (1979), compiled by R. Kinsey.
22. A. B. Smith, to be published.

TABLE 1. Observed Excitation Energies^a

No.	E _x (exp)	Attributed Components ^b	
		¹⁰⁷ Ag	¹⁰⁹ Ag
1	328 ± 13 ^c	325(3/2-)	311(3/2-)
2	419 ± 50 ^d	423(5/2-)	415(5/2-)
3	748 ± 25	787(3/2-)	702(3/2-) 706(3/2+) 731(3/2+)
4	908 ± 26	922(5/2+) 950(5/2-) 973(5/2-)	863(5/2-) 869(5/2+) 910(7/2+) 912(7/2-)
5	1150 ± 38	1050(?) 1143(5/2-) 1160(?)	1091(9/2-)
6	1286 ± 25	1222(5/2+) 1250(?) 1259(3/2+) 1325(?)	1200(?) 1260(1/2-) 1324(3/2-)
7	1507 ± 20	1465(3/2-) 1482(?)	1430(1/2-) 1490(?) 1524(3/2-)
8	1623 ± 30	1600(?) 1613(1/2-) 1660(1/2+) 1670(?)	1573(?) 1615(1/2-) 1653(1/2-) 1688(?)
9	1835 ± 20	-	-
10	1944 ± 26	-	-

^aAll energies in keV.

^bEnergies and Jπ values taken from Ref. 2.

^cUncertainties are RMS deviations from simple mean. Values of E_x are averages not necessarily corresponding to scattered-neutron resolutions.

^dEnergies uncertain due to presence of the second neutron-group from the source reaction.

Table 2. Optical-Model Parameters^a

Real Potential^b

$$V_0^c = 48.25 \text{ MeV}$$

$$r^d = 1.249 \text{ F}$$

$$a = 0.603 \text{ F}$$

$$J_v/A = 433.7 \text{ MeV-F}^3$$

$$V_0 r^2 = 75.27 \text{ MeV-F}^2$$

Imaginary Potential^e

$$W = 8.50 \text{ MeV}$$

$$r^d = 1.270 \text{ F}$$

$$a = 0.575 \text{ F}$$

$$J_w/A = 85.7 \text{ MeV-F}^3$$

$$W a = 4.89 \text{ MeV-F}$$

^aAlso spin-orbit potential of Thomas form¹³ with 6 MeV depth.

^bSaxon form.¹³

^cEnergy dependence of form $V = V_0 - 0.3 \cdot E$ (MeV).

^dAll radii expressed as $R = r \cdot A^{1/3}$.

^eSaxon-derivative form.¹³.