

(11) (A) No. 1,118,683

(45) ISSUED 820223

(52) CLASS 167-48

(51) INT. CL. A61K 49/02³

(19) (CA) **CANADIAN PATENT** (12)

(54) RADIOACTIVE SCANNING AGENTS
WITH STABILIZER

(72) Fawzi, Mahdi B.,
U.S.A.

(73) Granted to Procter & Gamble Company (The)
U.S.A.

(21) APPLICATION No. 330,719

(22) FILED 790627

(30) PRIORITY DATE U.S.A. (920,082) 780628

No. OF CLAIMS 22 - NO DRAWING

1118683

P&G Case 2583

RADIOACTIVE SCANNING AGENTS WITH STABILIZER

Mahdi Bakir Fawzi

ABSTRACT OF THE DISCLOSURE

Stable compositions, useful as technetium-99m-based scintigraphic agents, comprise gentisyl alcohol or a pharmaceutically-acceptable salt or ester thereof in combination with a pertechnetate reducing agent or dissolved in pertechnetate-99m ($^{99m}\text{TcO}_4^-$) solution. The compositions are especially useful in combination with a phosphate or phosphonate material which carries the radio-nuclide to bone, thus providing a skeletal imaging agent.

This invention relates to compositions useful in the preparation of stable scintigraphic scanning (or, "radio-diagnostic") agents employing technetium-99m. More particularly, gentisyl alcohol (or a soluble gentisyl alcohol compound) is used as a non-interfering stabilizer for such compositions.

Scintigraphy and similar radiographic techniques are finding ever increasing application in biological and medical research and diagnostic procedures. In general, scintigraphy procedures involve the preparation of radioactive scanning agents which, upon introduction into a biological subject, become localized in specific organs,



tissue or skeletal material which are under study. When so localized, traces, plots or scintiphotos of the distribution of the radioactive material can be made by various radiation detectors, i.e., such as traversing scanners, scintillation cameras and the like. The resultant distribution and corresponding relative intensities can then be used to indicate the position occupied by the tissue in which the radionuclide is localized as well as indicate the presence of aberrations, pathological conditions, and the like.

10 In general, and depending on the type of radionuclide used and the organ of interest, a scintigraphic scanning agent as used in the hospital comprises a radionuclide, a carrier agent designed to target the specific organ, various auxiliary agents which affix the radionuclide to the carrier, 15 water or other delivery vehicle suitable for injection into, or aspiration by, the patient, physiologic buffers and salts, and the like.

 Technetium-99m is a radionuclide which is widely known for use in tissue scanning agents. This radionuclide is conveniently available from commercial pertechnetate 20 sources.

 Pertechnetate is in the +7 oxidation state, which is too high to be used in the preparation of scanning agents such as those used for bone mineral and lung. This problem 25 is easily overcome by reducing the pertechnetate to what is believed to be the +3, +4 and/or +5 oxidation state.

 In general, ^{99m}Tc-labeled scanning agents are prepared by admixing a pertechnetate-99m isotonic saline solution with a pertechnetate reducing agent such as the 30 stannous, ferrous or chromous salt of sulfuric or hydrochloric acid, and the desired carrier agent for targeting

the organ of interest. For example, U.S. Patent 3,983,227, Tofe and Francis, discloses the use of reducing salts with radioactive pertechnetate solutions and organophosphonate bone-seeking carriers to prepare bone scanning agents.

5 U.S. Patent 4,002,730, Tofe, Hartman and Kretschmar, describes ^{99m}Tc lung scanning agents prepared by mixing a pertechnetate solution from commercial generators with stannous/starch particles which provide a combined reducing agent/carrier.

While such procedures provide scanning agents which
10 are superior to those previously available, they have shortcomings. Most notably, it has been found that conventional technetium-containing scintigraphic scanning agents are unstable in the presence of oxygen and radiolysis products. Accordingly, previously-described technetium-
15 based scanning agents have been made oxygen-free by saturating the composition with oxygen-free nitrogen gas or by preparing the agent in an oxygen-free atmosphere or in a vacuum. However, even such painstaking precautionary procedures are not entirely satisfactory, since it is extremely difficult
20 to maintain oxygen-free conditions. For instance, pertechnetate solutions may contain dissolved oxygen which, if not detected prior to combination with the pertechnetate reducing agent, forms a product which is unstable and results in the undesirable formation of free pertechnetate- ^{99m}Tc .

25 Others have disclosed means for overcoming the afore-said instability problem using chemical stabilizers. German Offenlegungsschrift 2,618,337, Tofe, published November 11, 1976, discloses the use of ascorbate stabilizers with technetium scanning agents. U.S. Patent 4,075,314, issued
30 February 21, 1978, discloses the use of ascorbate to inhibit

the oxidation of Sn^{+2} and to inhibit the reoxidation of reduced technetium in a pyrophosphate-based bone scanning agent.

It has now been discovered that gentisyl alcohol
5 compounds are safe, effective, and non-interfering stabilizers for scanning agents used in the radiographic diagnosis of tissues (including bone mineral) of humans and lower animals.

U.S. Patent No. 4,233,284 of M. B. Fawzi
10 granted November 11, 1980, said patent being entitled STABILIZED RADIOGRAPHIC SCANNING AGENTS, discloses the use of gentisic acid in radiodiagnostic compositions.

U.S. Patent No. 4,229,427 of
15 Whitehouse, granted October 21, 1980, said Patent being entitled RADIOACTIVE SCANNING AGENTS WITH HYDROQUINONE STABILIZER, teaches the use of hydroquinone in radiodiagnostic compositions.

SUMMARY OF THE INVENTION

Materials which are useful stabilizers for radio-diagnostic agents must exhibit the following properties:

- 5 1) Toxicological acceptability under the conditions of use;
- 2) The ability to stabilize the product for a reasonable period of storage and/or under use conditions; and
- 10 3) No substantial interference with the delivery of the radionuclide to the intended organ.

The present invention is based on the discovery that the gentisyl alcohol compounds, i.e., gentisyl alcohol and its water-soluble phenolate salts and esters, stabilize radiodiagnostic agents without interfering with the ability
15 of such agents to target specific organs. When used in the manner disclosed herein, the gentisyl alcohol compounds meet all of the above three criteria for a stabilizer.

The present invention provides highly stable compositions useful in the preparation of technetium-99m-based
20 scintigraphic scanning agents. The compositions of the present invention comprise a pertechnetate reducing agent or oxidized pertechnetate solution and an effective amount, sufficient to stabilize said compositions in the presence of oxygen and radiolysis products, of gentisyl alcohol or the
25 water-soluble, pharmaceutically-acceptable salts or esters thereof.

In a preferred method aspect, the present invention encompasses an improved method of preparing a technetium-based scanning agent comprising codissolving gentisyl
30 alcohol, or the water-soluble, pharmaceutically-acceptable

salts or esters thereof, and a pertechnetate reducing agent in an oxidized pertechnetate solution.

The stabilizing amount of gentisyl alcohol stabilizer used in the practice of this invention is an amount which
5 inhibits or reduces the oxidation of the pertechnetate reducing agent (e.g., oxidation of Sn^{+2} to Sn^{+4}) during storage and/or inhibits or reduces the reoxidation of reduced technetium and/or formation of technetium-labeled impurities during use.

10 All percentages and ratios herein are by weight, unless otherwise specified.

DETAILED DESCRIPTION OF THE INVENTION

This invention is based on the discovery that gentisyl alcohol and its phenolate salts and esters (all hereinafter "gentisols") can be utilized to prepare highly stable
5 technetium-99m-based scintigraphic scanning agents. As known in the art, ^{99m}Tc -labeled scanning agents are prepared by mixing a pertechnetate solution with a pertechnetate reducing agent. It has been found that small amounts of gentisyl alcohol, its salts or esters, can be combined with
10 either the pertechnetate solution or the pertechnetate reducing agent, thereby providing compositions uniquely suited for the preparation of superior, stable ^{99m}Tc scanning agents.

There are several compositional aspects to the
15 present invention. In one aspect, compositions of the present invention comprise a pertechnetate reducing agent which provides a pertechnetate reducing metal or cation, and the gentisol stabilizer. Such compositions comprise a stabilizing amount of the gentisol stabilizer. In a second
20 aspect, the compositions of the present invention comprise a pertechnetate solution having dissolved therein a stabilizing amount of the gentisol stabilizer.

When practicing the present invention, it is not critical which compositional form is used to prepare the
25 final technetium-based scanning agent. Commercial producers of pertechnetate generators may find it desirable to dissolve low levels of the gentisol stabilizer directly into the pertechnetate solution as it is eluted from the generator, or to incorporate a gentisol stabilizer directly on the
30 generator column. Alternatively, it may be more convenient

to combine the gentisol compound with the pertechnetate
reducing agent. In either case, upon combining the per-
technetate solution with the reducing agent and gentisol
stabilizer, an improved, highly stable scanning agent is
5 provided.

Gentisol stabilizer

Gentisyl alcohol (chemically: 2,5-dihydroxybenzyl
alcohol) is a metabolic product of Penicillium patulum
which can be prepared or isolated by several known methods.
10 See THE MERCK INDEX, 7th Edition, at p. 476.

The pharmaceutically-acceptable esters of gentisyl
alcohol can be prepared by standard well-known neutraliza-
tion and esterification procedures. A thorough discussion
of suitable procedures for preparing carboxylate esters of
15 alcohols can be found in Noller, The Chemistry of Organic
Compounds, Third Ed., 1966. In general, the pharmaceutically-
acceptable phenolate salts of gentisyl alcohol can be
prepared by acid-base neutralization using a base which
is selected to provide a water-soluble reaction product
20 having a pharmaceutically-acceptable counter ion. Similarly,
the pharmaceutically-acceptable esters of gentisyl alcohol
(esterified at the $-CH_2OH$ moiety) can be prepared by reacting
equimolar amounts of the gentisyl alcohol with selected
carboxylic acid or acid halide. Various salts, esters and
25 derivatives of gentisyl alcohol are known in the literature.

In practice, the salts and esters of gentisyl alcohol
suitable for use in the present invention can be selected
according to their solubility in a pertechnetate solution.
It is, of course, preferable that said salts and esters of

gentisyl alcohol be readily soluble in a pertechnetate solution. Accordingly, suitable gentisyl alcohol salts include the soluble alkali metal, alkaline earth metal, heavy metal and ammonium phenolate salts.

5 The pharmaceutically-acceptable esters of gentisyl alcohol (esterified at the $-\text{CH}_2\text{OH}$ group) which are sufficiently soluble in pertechnetate solutions include, for example, the C_1 to C_5 lower acid esters such as the esters of gentisyl alcohol with formic, acetic, propionic, butyric
10 and pentanoic acid.

Gentisyl alcohol, itself, is quite water-soluble, is soluble in TcO_4^- solutions, and is preferred for use herein.

Reducing Agent

15 In embodiments of this invention in which the gentisol stabilizer is combined with the pertechnetate reducing agent, the choice of reducing agent is not critical. As used herein the term "pertechnetate reducing agent" is intended to include compounds, complexes, or the like,
20 comprising a reducing ion capable of reducing heptavalent technetium (TcO_4^-) to trivalent, tetravalent and/or pentavalent technetium. Free metals such as tin are also known for use as pertechnetate reducing agents, although undissolved metal must be removed from the scanning solution
25 prior to injection into the patient. Thus, it is more convenient to use metal compounds which provide the reducing metal cation in soluble form.

Suitable pertechnetate reducing agents can be combined with numerous adjuvants such as fillers and skeletal-
30 or other organ-specific carriers. As disclosed by Tofe and

Francis, and Tofe, et al., above, skeletal scanning agents have been prepared utilizing metallic salts of sulfuric and hydrochloric acid such as stannous chloride, chromous chloride and ferrous sulfate as the pertechnetate reducing agent in combination with various organic phosphonates and/or phosphates as the bone seeking carrier. Other systems capable of reducing pertechnetate-99m include, for example, acid-thiosulfates, acid-hydrogen-sulfates, iron colloids, and acid-borohydrides. U.S. Patents 3,735,001 granted May 22, 1973; 3,863,004 granted January 28, 1975; 3,466,361 granted September 9, 1969; 3,720,761 granted March 13, 1973; 3,723,612 granted March 27, 1973; 3,725,295 granted April 3, 1973; 3,803,299 granted April 9, 1974; and 3,749,556 granted July 31, 1973; disclose

various pertechnetate reducing agents comprising reducing ions capable of reducing heptavalent pertechnetate to appropriate lower valence states.

The amount of the gentisol stabilizer utilized in embodiments of this invention in which the gentisol is combined with a reducing agent will vary depending on the ultimate use of the composition and the amount of inert or filler materials utilized. Too much gentisol stabilizer can cause excess soft tissue uptake and can interfere with organ (especially bone) imaging. In general, the weight ratio of gentisol stabilizer:reducing agent in a gentisol-plus-reducing agent composition is in the range of about 30:1 to about 1:30, preferably about 10:1 to about 1:1, more preferably about 5:1 to 1:1, most preferably about 3:1.

Where it is desirable to incorporate the gentisol stabilizer directly into the pertechnetate solution, the

soluble gentisol compound can be simply dissolved either during or after elution of the pertechnetate source. The elution process is thoroughly described in U.S. Patent 3,369,121.

5 In embodiments of the present invention in which the gentisol stabilizer is dissolved in a pertechnetate solution, the concentration of gentisol will vary somewhat depending upon the degree of aqueous dilution. With current commercial pertechnetate sources, a concentration of
10 gentisol no greater than about 0.1%, preferably no greater than 0.05%, by weight, provides adequate stability and does not interfere with organ distribution of the scanning agent. A concentration within the range of from about 0.0005% to about 0.05% is quite acceptable for many applica-
15 tions.

The present invention also encompasses an improved method of preparing technetium-based scanning agents comprising codissolving a gentisol stabilizer and a pertechnetate reducing agent in an aqueous pertechnetate solution. As
20 described above, the gentisol compound and pertechnetate reducing agent can be either simultaneously dissolved or sequentially dissolved in the pertechnetate solution. Either codissolving procedure results in an improved technetium-based scanning agent.

25 Scanning agents also typically employ carrier agents which direct, or "target", the radionuclide to the desired organ. Broadly speaking, there are two classes of such carrier agents: those which target soft tissue organs such as the heart, marrow, liver, spleen, kidneys and lungs;
30 and those which target calcified tissue, such as bone and


other tissues which may be undergoing pathological calcification. Examples of such carrier, or "targeting", agents for soft tissues include: colloidal sulfur, albumin, and the like. Targeting agents for bone mineral include the water-soluble phosphates and (preferably) phosphonates.

For purposes of the present invention a normal physiologically-acceptable pH of from about 3 to 8, preferably 4 to about 6, is used in the preparation of the scanning agent. At base pH's the gentisol may become colored, but this coloration does not substantially lessen product stability.


In a preferred and convenient embodiment of the present invention, a stable technetium-based skeletal scanning agent can be formed by the direct addition of an aqueous, radioactive pertechnetate solution to a composition comprising: an amount of a pertechnetate reducing agent containing a metallic reducing ion in water-soluble form, such as stannous chloride, sufficient to reduce said pertechnetate; an amount of a water-soluble gentisol compound sufficient to stabilize the skeletal scanning agent; and an amount of a skeletal-specific carrier compound selected from the mono-, di- or polyphosphonates sufficient to carry the radioactive technetium to bone. The ratio of reducing agent:phosphonate:gentisol can be adjusted to provide skeletal uptake with sufficiently low soft tissue uptake that scan quality is not undesirably affected.

A broad range of mono-, di- and polyphosphonates are now known to concentrate on the skeleton upon injection

of solutions thereof into a patient. Operable species for this purpose include mono-, di- and polyphosphonates selected from the group consisting of:




5 wherein each R is hydrogen or CH₂OH and n is an integer of from 3 to 10;




wherein R₁ is hydrogen, alkyl containing from 1 to about 20 carbon atoms, alkenyl containing from 2 to about 20
 10 carbon atoms, aryl (e.g., phenyl, naphthyl), phenyl-ethenyl, benzyl, halogen (e.g., chlorine, bromine and fluorine), hydroxyl, amino, substituted amino (e.g., dimethylamino, diethylamino, N-hydroxy-N-ethylamino, acetylamino), -CH₂COOH, -CH₂PO₃H₂, CH(PO₃H₂)(OH), or
 15 -[CH₂C(PO₃H₂)₂]_n-H where n = 1 to 15, R₂ is hydrogen,


lower alkyl (e.g., methyl, ethyl, propyl and butyl), amino, benzyl, halogen (e.g., chlorine, bromine, and fluorine), hydroxyl, $-\text{CH}_2\text{COOH}$, $-\text{CH}_2\text{PO}_3\text{H}_2$, or $-\text{CH}_2\text{CH}_2\text{PO}_3\text{H}_2$;



5 wherein n is an integer of from 3 to 9;




wherein each R_3 is hydrogen or lower alkyl (e.g., methyl, ethyl, propyl and butyl);




wherein n is an integer of from 2 to 4;


VI.




VII.




VIII.



IX.



X.



wherein X and Y are each hydrogen or hydroxy; and the non-toxic salts of each of the foregoing phosphonates which in an essentially neutral aqueous solution will react with hereinafter enumerated reducing/complexing materials; i.e., stannous, ferrous, or chromous salts to form the corresponding stannous, ferrous or chromous phosphonate salt. Suitable reactive phosphonate salts (hereinafter referred to as pharmaceutically acceptable salts) for use with the present invention include sodium, potassium, ammonium and low molecular weight substituted ammonium (e.g., mono-, di- and triethanol-amine and quaternary ammonium) salts of the above phosphonates and mixtures thereof.

Operable polyphosphonates of the above formula (I) include propane-1,2,3-triphosphonic acid; butane-1,2,3,4-tetraphosphonic acid; hexane-1,2,3,4,5,6-hexaphosphonic acid; hexane-1-hydroxy-2,3,4,5,6-pentaphosphonic acid; hexane-1,6-dihydroxy-2,3,4,5-tetraphosphonic acid; pentane-1,2,3,4,5-pentaphosphonic acid; heptane-1,2,3,4,5,6,7-heptaphosphonic acid; octane-1,2,3,4,5,6,7,8-octaphosphonic acid; nonane-1,2,3,4,5,6,7,8,9-nonaphosphonic acid; decane-1,2,3,4,5,6,7,8,9,10-decaphosphonic acid; and the pharmaceutically acceptable salts of these acids, e.g., sodium, potassium, ammonium, triethanolammonium, diethanolammonium, and monoethanolammonium salts.

Propane-1,2,3-triphosphonic acid and salts thereof can be prepared by a process disclosed in U.S. Patent No. 3,743,688 of D. A. Nicholson and D. Campbell, granted July 8, 1973.

Butane-1,2,3,4-tetraphosphonic acid and salts thereof can be prepared by a process disclosed in U.S. Patent No. 3,755,504 of D. A. Nicholson and D. Campbell, granted August 28, 1973.

The higher aliphatic vicinal polyphosphonates and salts thereof can be prepared by the process disclosed in U.S. Patent 3,584,035 granted June 8, 1971.

Among the operable polyphosphonates encompassed by the above formula (II) are ethane-1-hydroxy-1,1-diphosphonic acid; methanediphosphonic acid; methanhydroxydiphosphonic acid; ethane-1,1,2-triphosphonic acid; propane-1,1,3,3-

tetraphosphonic acid; ethane-2-phenyl-1,1-diphosphonic acid; ethane-2-naphthyl-1,1-diphosphonic acid; methane-phenyldiphosphonic acid; ethane-1-amino-1,1-diphosphonic acid; methanedichlorodiphosphonic acid; nonane-5,5-
 5 diphosphonic acid; n-pentane-1,1-diphosphonic acid; methanedifluorodiphosphonic acid; methanedibromodiphosphonic acid; propane-2,2-diphosphonic acid; ethane-2-carboxy-1,1-diphosphonic acid; propane-1-hydroxy-1,1,3-triphosphonic acid; ethane-2-hydroxy-1,1,2-
 10 triphosphonic acid; ethane-1-hydroxy-1,1,2-triphosphonic acid; propane-1,3-diphenyl-2,2-diphosphonic acid; nonane-1,1-diphosphonic acid; hexadecane-1,1-diphosphonic acid; pent-4-ene-1-hydroxy-1,1-diphosphonic acid; octadec-9-ene-1-hydroxy-1,1-diphosphonic acid; 3-phenyl-1,1-diphos-
 15 phono-prop-2-ene; octane-1,1-diphosphonic acid; dodecane-1,1-diphosphonic acid; phenylaminomethanediphosphonic acid; naphthylaminomethanediphosphonic acid; N,N-dimethylamino-methanediphosphonic acid; N-(2-dihydroxyethyl)-aminomethane-diphosphonic acid; N-acetylamino-methanediphosphonic acid;
 20 aminomethanediphosphonic acid; dihydroxymethanediphosphonic acid; and the pharmaceutically acceptable salts of these acids, e.g., sodium, potassium, ammonium, triethanolammonium, diethanolammonium, and monoethanolammonium salts.

Ethane-1-hydroxy-1,1-diphosphonic acid, an
 25 especially preferred polyphosphonate, has the molecular formula $\text{CH}_3\text{C}(\text{OH})(\text{PO}_3\text{H}_2)_2$. (According to nomenclature by radicals, the acid might also be named 1-hydroxyethylidene diphosphonic acid.)

While any pharmaceutically acceptable salt of ethane-1-hydroxy-1,1-diphosphonic acid can be used in the practice of this invention, mixtures of the disodium and trisodium salts are most preferred. The other sodium, potassium, ammonium, and mono-, di-, and triethanolammonium salts and mixtures thereof are also suitable, provided caution is observed in regulating the total intake of cation species in the salt composition. These compounds can be prepared by any suitable method; however, an especially preferred method is disclosed in U.S. Patent 3,400,149 granted September 3, 1968.

Methanehydroxydiphosphonic acid and related compounds operable herein can be prepared, for example, by reaction of phosgene with an alkali metal dialkylphosphite. A complete description of these compounds and a method for preparing same is found in U.S. Patent 3,422,137 granted January 14, 1969.

Methanedihydroxydiphosphonic acid and salts useful herein and a method for preparing same are disclosed in U.S. Patent 3,497,313 granted February 24, 1970.

Methanediphosphonic acid and related compounds useful herein are described in detail in U.S. Patent 3,213,030, granted October 19, 1965. A preferred method of preparing such compounds is disclosed in U.S. Patent 3,251,907 granted May 17, 1966.

Ethane-1,1,2-triphosphonic acid and related compounds which can be used in the compositions of this invention, as well as a method for their preparation,

are fully described in U.S. Patent 3,551,339 granted December 29, 1970.

Propane-1,1,3,3-tetraphosphonic acid and related compounds useful herein, and a method for preparing same, 5 are fully disclosed in U.S. Patent 3,400,176 granted September 3, 1968. The higher methylene interrupted methylene diphosphonate polymers can be prepared by the polymerization of ethylene-1,1-diphosphonate.

Pentane-2,2-diphosphonic acid and related compounds 10 can be prepared in accordance with the method described by G. M. Kosolopoff in J. Amer. Chem. Soc., 75, 1500 (1953).

Operable phosphonates of formula (III) above include the following:

Methanecyclobutylhydroxydiphosphonic acid
15 Methanecyclopentylhydroxydiphosphonic acid
Methanecyclohexylhydroxydiphosphonic acid
Methanecycloheptylhydroxydiphosphonic acid
Methanecyclooctylhydroxydiphosphonic acid
Methanecyclononylhydroxydiphosphonic acid
20 Methanecyclodecylhydroxydiphosphonic acid.

Each of the sodium, potassium, ammonium, monoethanol- ammonium, diethanolammonium, and triethanolammonium salts of the above-recited methanecycloalkylhydroxydiphosphonic acids as well as any other pharmaceutically acceptable 25 salt of these acids, also selectively seek the skeleton.


The phosphonates of formula (III) can be prepared by methods fully described in U.S. Patent 3,584,125, granted June 8, 1971.

The preferred phosphonates of formula (IV) for the purpose of this invention are tris(phosphonomethyl)amine; tris(1-phosphonoethyl)amine; tris(2-phosphono-2-propyl)amine; and their pharmaceutically acceptable salts.

5 Tris(phosphonomethyl)amine is especially preferred. The following are exemplary of compounds which can also be used:

- (a) bis(phosphonomethyl)-1-phosphonoethyl amine;
- (b) bis(phosphonomethyl)-2-phosphono-2-propyl amine;
- 10 (c) bis(1-phosphonoethyl)phosphonomethyl amine;
- (d) bis(2-phosphono-2-propyl)phosphonomethyl amine;
- (e) tris(1-phosphono-1-pentyl)amine;
- (f) bis(phosphonomethyl)2-phosphono-2-hexyl amine; and
- (g) the pharmaceutically acceptable salts of acids
- 15 (a) through (f), e.g., sodium, potassium, ammonium, triethanolammonium, diethanolammonium, and monoethanolammonium salts.

The tris(phosphonoalkyl)amines can be prepared, for example, by first preparing the corresponding ester in
20 accordance with the general reaction:



wherein R is alkyl and R₁ and R₂ are hydrogen or lower alkyl.

The free acids can be prepared by hydrolysis of the ester using strong mineral acids such as hydrochloric acid. The salts are, of course, prepared by neutralizing the acid with the base of the desired cation. The preparation of tris(phosphonoalkyl)amines is fully disclosed by Irani, et al., in Canadian Patent 753,207, issued February 21, 1967.

The phosphonates of formula (V) include the following: (1) 3,3,4,4,5,5-hexafluoro-1,2-diphosphonocyclopent-1-ene; (2) 3,3,4,4-tetrafluoro-1,2-diphosphonocyclobut-1-ene; and (3) 3,3,4,4,5,5,6,6-octafluoro-1,2-diphosphonocyclohex-1-ene.

The perfluorodiphosphonocycloalkenes can be prepared, for example, by reacting trialkyl phosphites with 1,2-dichloroperfluorocycloalk-1-enes in accordance with the procedures fully described by Frank in J. Org. Chem., 31, #5, p. 1521.

The phosphonate of formula (VI) is referred to herein as cyclic tetraphosphonic acid. This compound

and its pharmaceutically acceptable salts can be prepared by any suitable method; however, an especially preferred method is disclosed by Oscar T. Quimby, U.S. Patent 3,387,024 granted June 4, 1968.

5 Operable phosphonates encompassed by the above formula (VII) are ethene-1,2-dicarboxy-1-phosphonic acid; and the pharmaceutically acceptable salts of these acids, e.g., sodium, potassium, ammonium, triethanolammonium, diethanolammonium, and monoethanolammonium salts. While
10 the above formula (VII) is representative of cis-isomers, the corresponding trans-isomers are also useful herein. Reference hereinafter to ethene-1,2-dicarboxy-1-phosphonic acid or salts thereof, unless otherwise specified, is intended as contemplating the cis- and trans-isomers and
15 mixtures thereof.

Ethene-1,2-dicarboxy-1-phosphonic acid and related compounds useful herein can be prepared by reaction of an ester of acetylenedicarboxylic acid and a dialkyl phosphite followed by hydrolysis and saponification.
20 This method is more fully described in U.S. Patent 3,584,124, granted June 8, 1971.

The sodium salt of formula (VIII) can be made by the rearrangement reaction of a 2-haloethane-1-hydroxy-1,1-diphosphonic acid with about 3 equivalents of sodium
25 hydroxide as disclosed in U.S. Patent 3,641,126.

The phosphonate of formula (IX) can be made by the method of German Offenlegungsschrift 2,026,078.

Operable carboxyphosphonates of the above formula (X) include ethane-1,2-dicarboxy-1,2-diphosphonic acid;

ethane-1,2-dicarboxy-1,2-dihydroxy-1,2-diphosphonic acid;
ethane-1,2-dicarboxy-1-hydroxy-1,2-diphosphonic acid;
and the pharmaceutically acceptable salts of these acids,
e.g., sodium, potassium, ammonium, triethanolammonium,
5 diethanolammonium and monoethanolammonium salts.

Ethane-1,2-dicarboxy-1,2-diphosphonic acid, a preferred carboxyphosphonate herein, has the molecular formula $\text{CH}(\text{COOH})(\text{PO}_3\text{H}_2)\text{CH}(\text{COOH})(\text{PO}_3\text{H}_2)$. The most convenient crystallizable salts of this acid are obtained
10 when three, four or five of the acid hydrogens are replaced by sodium.

While any pharmaceutically acceptable salt of ethane-1,2-dicarboxy-1,2-diphosphonic acid can be used in the practice of this invention, the tetrasodium
15 dihydrogen salt, the trisodium trihydrogen salt, the disodium tetrahydrogen salt, the monosodium pentahydrogen salt, and the mixtures thereof are useful. The other potassium, ammonium, and mono-, di-, and triethanolammonium, etc., salts and mixtures thereof are
20 also suitable, provided caution is observed in regulating the total intake of cation species in the salt composition.

Ethane-1,2-dicarboxy-1,2-diphosphonic acid and suitable salts thereof can be prepared in any convenient manner. For example, the reaction described by Pudovik
25 in "Soviet Research on Organo-Phosphorus Compounds", 1949-1956, Part III, 547-85c. can be used to prepare the ester of ethane-1,2-dicarboxy-1,2-diphosphonic acid which in turn can, by ordinary hydrolysis reactions, be converted

to the free acid form. Neutralization by alkali compounds such as sodium hydroxide, potassium hydroxide, carbonates and the like can be used to prepare a desired salt of the acid. A more detailed description of the preparation of these compounds is described in U.S. Patent 3,562,166 granted February 9, 1971.

Ethane-1,1-dicarboxy-1,2-dihydroxy-1,2-diphosphonic acid and related compounds useful herein can be prepared by reaction of an ester of ethane-1,2-dicarboxy-1,2-diphosphonic acid and an alkali metal hypohalite followed by hydrolysis and saponification. This method is more fully described in U.S. Patent 3,579,570 granted May 18, 1971.

Mixtures of any of the foregoing phosphonic acids and/or salts can be used in the practice of this invention.

In a highly preferred embodiment of this invention, mixtures of disodium- and trisodium-ethane-1-hydroxy-1,1-diphosphonate salts wherein the mole ratio of the disodium salt to trisodium is from about 4:1 to 1:1, more preferably 3:1 to 1:1, are employed in the pertechnetate reducing agent along with a gentisol stabilizer and a reducing metallic ion. These preferred phosphonate/reducing ion/gentisol mixtures provide especially good scintiscans, with excellent stability and skeletal uptake and little soft tissue uptake.

The aminophosphonates, especially the aminopropane diphosphonates disclosed in U.S. Patents 4,054,598 and 3,962,432 are also useful in preparing bone scanning agents stabilized with gentisols.

The following examples are illustrative of compositions made according to this invention. The quantity of each component listed is in milligrams.

EXAMPLE

	<u>COMPONENT</u>	I	II	III	IV	V	VI	VII	VIII	IX	X
	Disodium-ethane-1-hydroxy-1,1-diphosphonate	5.9		5.9	5.9	5.9				4	3
5	Trisodium-ethane-1-hydroxy-1,1-diphosphonate		5.9							1.9	
	Disodium-methane-diphosphonate								5.9		
	Trisodium-methane-diphosphonate							5.9			2.9
	Dichloromethanediphosphonic acid						5.9				
10	Stannous chloride	0.16	0.16			0.16		0.16			0.16
	Ferrous sulfate			0.16			0.16		0.16		
	Chromous chloride				0.16					0.16	
	Sodium chloride	27		27		27		27		27	27
	Glucose		27		27		27				
15	Sodium gentisylate						0.5				0.5
	Gentisyl alcohol	0.2	0.5					0.5		0.1	
	Gentisyl acetate			0.7	0.2				0.7		
	Diethanolammonium gentisylate					0.5					

Each of the compositions of Examples I through X, upon addition of about 5 ml. of a pertechnetate-99m solution from a commercial technetium source, and thorough shaking, yields a skeletal scanning agent suitable for intravenous injection into a human patient. The in-use stability of such skeletal scanning agents is more than three hours, which is ample under ordinary hospital conditions. Preferably, about 1 ml. of the skeletal scanning solution is used in an adult of about 50-100 kilograms body weight and is injected slowly, over a period of about 30 seconds. Administration is preferably done within about three hours of preparation. Kits can, of course, contain multiples or fractions of the above amounts to allow preparation of sufficient agent from a single vial to perform any desired number of scans.

The following example is illustrative of a pertechnetate solution having dissolved therein a gentisol stabilizer.

EXAMPLE XI

A collecting vial containing 0.1 mg. of gentisyl alcohol is placed at the elute orifice of a pertechnetate-99m generator. Saline elute is collected in the vial and completely dissolves the gentisyl alcohol.

Approximately 5 ml. of the pertechnetate solution, with dissolved gentisyl alcohol, is added to a pertechnetate reducing agent comprising 5.9 mg. of the sodium salt of ethane-1-hydroxy-1,1-diphosphonic acid and 0.16 mg. of stannous chloride. After thorough shaking, a stable bone scanning agent suitable for intravenous injection into a human patient is prepared.

In the above example, gentisyl alcohol is replaced by equivalent amounts of gentisyl formate, sodium gentisylate, and gentisyl butyrate, respectively, and stable scanning agents are prepared.

5 The use of methanehydroxydiphosphonate (MHDP) as a highly preferred bone scanning agent which provides exceptionally sharp scintiphotos and excellent lesion detection is described in U.S. Patent No. 4,247,534 of Bevan, granted January 27, 1981, said patent being entitled
10 RADIOGRAPHIC SCANNING AGENT. The following example illustrates the preparation of a freeze-dried product containing MHDP as the bone-seeking agent.

EXAMPLE XII

15 Methanehydroxydiphosphonate, mixture of di-, tri-sodium salts (2.0 mg), stannous chloride (0.16 mg.), and gentisyl alcohol (0.50 mg.) are dissolved in 1 ml. of deoxygenated water at room temperature. The aqueous solution is filtered through a "Millipore"* filter and freeze-dried on a commercial apparatus.

20 The freeze-dried powder prepared in the foregoing manner is admixed with about 5 ml. of a pertechnetate-99m solution from a commercial source. The freeze-dried powder dissolves readily and a stable skeletal scanning agent suitable for intravenous use is secured.

25 The foregoing examples have illustrated the use of gentisols in the preparation of scanning agents using various phosphonate materials to target bone mineral. Inorganic phosphates are also useful in the manufacture

* Trademark

of bone scanning agents and radiodiagnostic products containing inorganic phosphates can also be stabilized with the gentisols herein.

U.S. Patent 4,016,249, of Adler et al., issued April 5, 1977, said patent being entitled "Bone Seeking Technetium 99M Complex", contains a succinct disclosure of the use of inorganic phosphates of various types in the manufacture of bone scanning agents. In particular, certain soluble pyrophosphate species having a molecular weight of less than about 300, said pyrophosphate containing no more than about 25% branched-chain polyphosphate, are quite useful for bone scanning. As with the organophosphonates, the pyrophosphate is conveniently used by admixture with a reducing salt for pertechnetate, thereby providing a kit. In use, the kit is charged with an aqueous solution of pertechnetate, whereupon the heptavalent technetium is reduced to a lower oxidation state, where it combines with the pyrophosphate. When injected into the patient the pyrophosphate targets bone mineral with the technetium radionuclide, in the manner of the organophosphonates. An example of such a product is as follows.

EXAMPLE XIII

<u>Ingredient</u>	<u>Amount (mg.)</u>
Gentisyl alcohol	0.20
Stannous chloride	1.30
Sodium pyrophosphate*	40.0

*As disclosed in U.S. Patent 4,016,249.

The composition of Example XIII is prepared by simple admixture of the listed ingredients and used in the manner disclosed above.

In an alternate mode, the composition of Example XIII is prepared by dissolving the ingredients in water (20 ml.), sterilizing by filtration and freeze-drying using standard equipment.

Highly preferred, gentisol-stabilized bone scanning agents are as follows.

10

EXAMPLE XIV

<u>Ingredient</u>	<u>Milligrams/vial</u>
EHDP*	5.9
SnCl ₂	0.16
Gentisyl alcohol	0.50
15 NaCl	26.5

*Mixture of di- and tri-sodium salts of ethane-1-hydroxy-1,1-diphosphonic acid.

The ingredients are dry mixed. Five mls. of eluate from a commercial pertéchnetate source added to one vial of the composition of Example XIV provides sufficient solution for 5 bone scans.

The composition of Example XIV is modified by replacing the EHDP with an equal amount of 3-amino-1-hydroxypropane-1,1-diphosphonic acid and a stabilized bone imaging agent is secured.

EXAMPLE XV

<u>Ingredient</u>	<u>Milligrams/vial</u>
MHDP*	2.0
SnCl ₂	0.16
5 Gentisyl alcohol	0.50.
NaCl	26.5

*Mixture of sodium salts of methanehydroxydiphosphonic acid.

The ingredients are dry mixed. Five mls. of eluate from a commercial pertechnetate source added to one vial of the composition of Example XV provides sufficient solution for 10 5 bone scans.

EXAMPLE XVI

Ethane-1-hydroxy-1,1-diphosphonate, mixture of di-, and trisodium salts (5.9 mg.), stannous chloride (0.16 mg.), 15 and gentisyl alcohol (0.50 mg.) are dissolved in 1 ml. of deoxygenated water at room temperature. The aqueous solution is filtered through a millipore filter and freeze-dried on a commercial apparatus.

The freeze-dried powder prepared in the foregoing 20 manner is admixed with about 5 ml. of a pertechnetate-99m solution from a commercial source. The freeze-dried powder dissolves readily and a stable skeletal scanning agent suitable for intravenous use is secured.

As can be seen from the foregoing, gentisol stabilizers 25 provide excellent storage stability and in-use stability for ^{99m}Tc-based scanning agents. When an organ-specific carrier such as an organophosphonate or inorganic phosphate is used in such compositions in the manner disclosed, it

is preferred that the weight ratio of gentisol:carrier be in the range of from about 10:1 to about 1:30, preferably 1:1 to about 1:20, most preferably from about 1:5 to about 1:12.

5 As disclosed in the art, the weight ratio of reducing agent:carrier in such compositions is generally in the range 1:50 to about 1:20, although this ratio can vary with the type of organ-specific carrier being used. The gentisol stabilizer does not appreciably alter these ratios.

10 Stannous gentisylate, i.e., the water-soluble product prepared by allowing stannous chloride to react with gentisyl alcohol in oxygen-free water for ca. 1 hour and freeze-drying the reaction mixture, can be used as a combined source of reducing agent and stabilizer in the
15 herein-disclosed compositions and processes.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A stable composition, useful in the preparation of technetium-99m-based radiographic scanning agents, comprising: a pertechnetate reducing agent; and a stabilizing amount of a gentisol stabilizer selected from gentisyl alcohol and the water-soluble, pharmaceutically-acceptable salts and esters thereof.
2. A composition according to Claim 1, comprising a weight ratio of gentisol stabilizer to reducing agent in the range of about 30:1 to about 1:30.
3. A composition according to Claim 2, wherein the ratio of stabilizer to reducing agent is in the range of from about 10:1 to about 1:1.
4. A composition according to Claim 3, wherein the gentisol stabilizer is gentisyl alcohol.
5. A composition according to Claim 1, wherein the pertechnetate reducing agent is selected from the group consisting of the soluble stannous, chromous, titanous and ferrous salts.
6. A composition according to Claim 5 wherein the pertechnetate reducing agent is selected from the group consisting of stannous chloride, chromous chloride and ferrous sulfate.

7. A composition according to Claim 1 wherein the reducing agent is stannous chloride, the gentisol is gentisyl alcohol, and the weight ratio of gentisol:SnCl₂ is about 5:1 to about 1:1.

8. A composition according to Claim 1, additionally comprising an organ-specific carrier.

9. A composition according to Claim 8 wherein the organ-specific carrier is an organophosphonate.

10. A composition according to Claim 9 wherein the organophosphonate is selected from the group consisting of ethane-1-hydroxy-1,1-diphosphonic acid, methanediphosphonic acid, dichloromethanediphosphonic acid, methanehydroxy-
5 diphosphonic acid, and the soluble salts and esters thereof.

11. A composition according to Claim 10 wherein the pertechnetate reducing agent is stannous chloride, the organ-specific carrier is a sodium salt form of ethane-1-hydroxy-1,1-diphosphonic acid, and the gentisol stabilizer
5 is gentisyl alcohol.

12. A composition according to Claim 10 wherein the pertechnetate reducing agent is stannous chloride, the organ-specific carrier is a sodium salt form of methanediphosphonic acid, and the gentisol stabilizer
5 is gentisyl alcohol.

13. A composition according to Claim 10 wherein the pertechnetate reducing agent is stannous chloride, the organ-specific carrier is a sodium salt form of dichloromethanediphosphonic acid, and the gentisol stabilizer is gentisyl alcohol.

14. A composition according to Claim 10 wherein the pertechnetate reducing agent is stannous chloride, the organ-specific carrier is a sodium salt form of methane-hydroxydiphosphonic acid, and the gentisol stabilizer is gentisyl alcohol.

15. A composition according to Claim 9 wherein the organophosphonate is a 3-aminopropane-1-hydroxy-1,1-diphosphonate.

16. A composition according to Claim 8 wherein the organ-specific carrier is a water-soluble inorganic phosphate.

17. A composition according to Claim 16 wherein the inorganic phosphate is a sodium pyrophosphate.

18. A composition useful in the preparation of technetium-99m-based radiographic scanning agents, comprising: an oxidized pertechnetate solution having dissolved therein a stabilizing amount of a gentisol stabilizer selected from gentisyl alcohol or a pharmaceutically-acceptable salt or ester thereof.

19. A composition according to Claim 18, comprising an oxidized pertechnetate solution having dissolved therein no more than about 0.1% by weight of a gentisol stabilizer.

20. A composition according to Claim 19, comprising an oxidized pertechnetate solution having dissolved therein no more than about 0.1%, by weight, of gentisyl alcohol.

21. A method of preparing a stabilized technetium-99m-based scanning agent, comprising codissolving a stabilizing amount of a gentisol stabilizer selected from gentisyl alcohol or a pharmaceutically-acceptable salt or ester thereof with an aqueous solution of radioactive technetium in the +3, +4 or +5 valence state.

22. A method according to Claim 21 wherein an organ-seeking carrier is included in the solution.



SUBSTITUTE

REPLACEMENT

SECTION is not Present

Cette Section est Absente