

REFERENCE

2163/80

INTERNATIONAL CENTRE FOR
THEORETICAL PHYSICS

ARE LEPTON AND QUARK FAMILIES QUANTIZED DYNAMICAL SYSTEMS?

W. Królikowski




INTERNATIONAL
ATOMIC ENERGY
AGENCY



UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION

1980 MIRAMARE-TRIESTE



International Atomic Energy Agency
and
United Nations Educational Scientific and Cultural Organization

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

ARE LEPTON AND QUARK FAMILIES QUANTIZED DYNAMICAL SYSTEMS? *

W. Królikowski **

International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

Lepton and quark families $\{v_N\}$, $\{e_N\}$ and $\{u_N\}$, $\{d_N\}$ are conjectured to be quantum-dynamical systems in the space of generations $N = 0, 1, 2, \dots$. Such a conjecture, called the "zereth quantization", implies the form of lepton and quark mass spectra, if lepton and quark families are supposed to be almost in thermal equilibrium with the rest of the Universe. Toponium is predicted at about 38 GeV, while the next charged lepton at 28.5 GeV.

MIRAMARE - TRIESTE

April 1980

* To be submitted for publication.

** On leave of absence from Institute of Theoretical Physics, Warsaw University, Warsaw, Poland.

In order to introduce the idea which is proposed in this paper, let us begin with the following pedagogical example. Consider a bound particle in an external potential (e.g. an electron in a hydrogen atom, where the proton is treated as infinitely heavy). Assume further that this system is in thermal equilibrium with a heat reservoir (e.g. our hydrogen atom may be put into a cavity with black-body radiation). Then the particle is in a mixed state given by the density matrix ¹⁾

$$\rho = \sum_n |n\rangle \rho_n \langle n|, \quad \sum_n \rho_n = 1, \quad (1)$$

where $H|n\rangle = E_n|n\rangle$ and

$$\rho_n = \frac{1}{Z} \exp(-\beta E_n), \quad Z = \sum_n \exp(-\beta E_n), \quad \beta = 1/kT. \quad (2)$$

In the mixed state ρ , the mass m of the particle is statistically distributed over the component states n with the weights ρ_n

$$\mu_n = m \rho_n, \quad \sum_n \mu_n = m. \quad (3)$$

Of course, if the state ρ were a pure state n_0 , i.e. if $\rho = |n_0\rangle \langle n_0|$, then the total amount of mass m would be concentrated in the particular component state n_0 , $\mu_n = m \delta_{nn_0}$. Obviously, the numbers μ_n are not masses of the particle in the component states n , but masses of the fractions of the particle in these component states. Note that the same as above concerning the mass m can be also said about the electric charge Q of the particle.

The above example concerns the familiar level of quantum theory which is the first quantization. It may suggest, however, some extensions in a new direction. Namely, one may ask the exciting question, whether there is possibly a more primordial level of quantum theory, where particles are fractions of some new quantum-dynamical systems which might be called the "particle families". If it is so, the latter are to be identified with the

experimentally observed (directly or indirectly) lepton or quark families

$f = \nu, e, u, d$, each consisting of several generations $N = 0, 1, 2, \dots$, viz. ²⁾

$$f = \{f_N\}, \quad f_N = \begin{cases} \nu_N = \nu_e, \nu_\mu, \nu_\tau, \dots \\ e_N = e, \mu, \tau, \dots \\ u_N = u, c, (t), \dots \\ d_N = d, s, b, \dots \end{cases}, \quad (4)$$

where electric charges of f_N particles are independent of N , $Q_{f_N} = Q_f$ with $Q_f = 0, -1, \frac{2}{3}, -\frac{1}{3}$ for $f = \nu, e, u, d$, respectively. Such a level of quantum theory, where particle families are quantum-dynamical systems, may be called the "zereth quantization" ³⁾ because it is analogical to but logically precedes the first quantization, where particles are quantum-dynamical systems. On the other hand, in the second quantization logically following the first, particle fields are quantum-dynamical systems.

If the conjecture of the zeroth quantization is true, each lepton or quark family f can be described by the density matrix

$$\rho_f = \sum_N |N\rangle \rho_{f_N} \langle N|, \quad \sum_N \rho_{f_N} = 1, \quad (5)$$

where $H_f |N\rangle = E_{f_N} |N\rangle$, the operator H_f being an unknown Hamiltonian of f family in the zeroth quantization. The density matrix (5) gives the statistical distribution of some mass M_f of f family over the generations $N = 0, 1, 2, \dots$ with the weights ρ_{f_N}

$$m_{f_N} = M_f \rho_{f_N}, \quad \sum_N m_{f_N} = M_f < +\infty. \quad (6)$$

Here, the numbers m_{f_N} are masses of f_N particles, as the latter are the fractions of f family in the component states N . We can see from Eq.(6) that

the masses m_{f_N} of f_N particles depend on the state ρ_f of f family. The observed masses m_{f_N} correspond to the actual state ρ_f which ought to be theoretically determined. Note that, in contrast to the mass $M_f = \sum_N m_{f_N}$,

no electric charge $\sum_N Q_{f_N}$ can be ascribed to f family, because all f_N particles possess equal electric charges $Q_{f_N} = Q_f$, so that

$Q_{f_N} \neq \left(\sum_N Q_{f_N} \right) \rho_{f_N}$. Thus, electric charge as well as all space-time quantities (e.g. position, momentum and spin) seem to be particle attributes.

The first quantization is needed to localize in space-time the fractions of particle families which become particles thereby. Then the second quantization allows them to interact via local field interactions connected with the existence of particle charges.

To proceed further with the idea of the zeroth quantization and then pass to its applications we must be more specific about the Hamiltonian H_f and the distribution ρ_{f_N} .

First of all let us note that the particle family f appears as a one-dimensional quantum-dynamical system since its states can be apparently numerated by one quantum number $N = 0, 1, 2, \dots$. So, as a first guess, it is very natural to try for f family a one-dimensional oscillator with the canonical variables $q = (a + a^\dagger)/\sqrt{2}$ and $p = (a - a^\dagger)/i\sqrt{2}$, where $[a, a^\dagger] = 1$. Then the Hamiltonian of f family is

$$H_f = \omega_f a^\dagger a + \theta(N_f - a^\dagger a) V_f, \quad (7)$$

where $\theta(x) = 0$ or 1 for $x \geq 0$ or $x < 0$ and $V_f = +$ in order to provide the possible cut $N \leq N_f$ for the number of generations of f family. The eigenvalues of H_f are

$$E_{f_N} = \omega_f N \text{ for } N \leq N_f. \quad (8)$$

Assumption (7) may be considered as the zero-order approximation in a weak coupling regime for the zeroth quantization. In this case, introducing the phase ϕ of $a^\dagger = \sqrt{a^\dagger} \exp(i\phi)$ which is canonically conjugate to $\pm a^\dagger a = -id/d\phi$, we get, respectively, for the dynamical systems considered

on the level of the zeroth, first and second quantization the following co-ordinates:

$$\phi, \quad \vec{r}(\phi), \quad \psi(\vec{r}), \quad (9)$$

where $\vec{r}(\phi)$, or relativistically $x^\mu(\phi)$, represents a string of the dual model! However, in contrast to this model where the usual first quantization follows by averaging ⁴⁾

$$\vec{r} = \frac{1}{2\pi} \int_{-\pi}^{\pi} d\phi \vec{r}(\phi), \quad (10)$$

in our case the usual first quantization can be obtained for f_N particles ($N = 0, 1, 2, \dots$) by the Fourier transformation

$$\vec{r}_N = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} d\phi \vec{r}(\phi) \cos(N\phi). \quad (11)$$

So the effect of dual excitations around the usual first-quantization motion is absent in our case.

The Hamiltonian (7) of f family is perturbed by the interaction with the rest of the Universe which is responsible for setting up the distribution ρ_{f_N} of f family. Let us suppose that, if the electric charges Q_f of f_N particles can be neglected, f family is in thermal equilibrium with the rest of the Universe considered as a heat and N -number reservoir. Then we get the grand canonical distribution

$$\rho_{f_N} |_{Q_f=0} = \frac{1}{Z_f} \exp[\beta(\mu_f N - E_{f_N})], \quad (12)$$

where

$$Z_f = \sum_N \exp[\beta(\mu_f N - E_{f_N})], \quad \beta = 1/kT, \quad (13)$$

the constant μ_f being a chemical potential. The restriction to the approximation of $Q_f = 0$ is in Eq.(12) necessary, since electric charge contributes to particle mass and so it is a kind of "source" of the "field" ρ_{f_N} in the space of generations $N = 0, 1, 2, \dots$ (other virtual "sources" of ρ_{f_N} are neglected here).

If $\mu_f \neq 0$, the grand canonical ensemble (12) implies the conservation of N number for the Universe as a whole. This conservation, appearing here on the level of the zeroth quantization, corresponds to the (sometimes approximate) flavour conservations on the level of the second quantization (where N number for an antiparticle comes with minus sign into the balance). For example, the N -number conservation forbids such unwanted processes as $\mu^- + e^- e^+ e^-$, $\mu^- + e^- \gamma$ and $\tau^- + e^- e^+ e^-$ or $\mu^- e^+ e^-$ or $\mu^- \mu^+ \mu^-$, $\mu^- + e^- \gamma$ or $\mu^- \gamma$, but it would allow for $\tau^- + \mu^- \bar{\nu}_e \nu_\mu$ if there were charged and/or neutral intermediate bosons of the generation $N = 1$ (which, however, would get presumably much larger masses than W^\pm and Z belonging to the generation $N = 0$). The proton decay $p \equiv uud + e^+ \bar{\nu}_e \bar{\nu}_e$ is also allowed from the viewpoint of N -number conservation, depending on the existence of appropriate intermediate bosons (of the generation $N = 0$, but necessarily with a big mass). Of course, the flavour conservations in weak processes involving quarks are only approximate in the cases where Cabibbo-like mixing appears.

Now, let us switch on the electric charges Q_f of f_N particles and suppose that the thermal equilibrium (12) is then perturbed linearly by a term proportional to $Q_{f_N}^2 = Q_f^2$

$$\rho_{f_N} = A_f \rho_{f_N} |_{Q_f=0} + B_f Q_f^2, \quad (14)$$

where

$$A_f + B_f Q_f^2 N_f = 1 \quad (15)$$

due to the normalization condition $\sum_N \rho_{f_N} = 1$. Here, because of the constant term in the distribution (14), it must ^N be assumed that the number of generations of f family is finite, $N \leq N_f < +\infty$.

In consequence of Eqs.(8) and (12)-(15) one can eventually write

$$\rho_{f_N} = (1 - B_f Q_f^2 N_f) \frac{1}{Z_f} \lambda_f^{2N} + B_f Q_f^2, \quad (16)$$

where

$$Z_f = \frac{\lambda_f^{2N_f} - 1}{\lambda_f - 1}, \quad \lambda_f^2 = \exp[\beta(\mu_f - \omega_f)]. \quad (17)$$

We can see that in the case of Eqs.(8), (12) and (14) each f_N particle of N generation contains N identical Bose objects characterized thermodynamically — on the level of the zeroth quantization — by the energy ω_f and chemical potential μ_f . These objects, playing the role of excitation quanta, may be called the "generatons" as they are responsible for the higher generations $N > 0$ of leptons and quarks.

In order to compare the distribution (16) with experimental data, we multiply it by M_f , obtaining the mass spectrum of f family. Introducing into it the initial mass values m_{f_0} and m_{f_1} we obtain the following spectral formula:

$$m_{f_N} = (m_{f_0} + \kappa_f Q_f^2) \lambda_f^{2N} - \kappa_f Q_f^2, \quad (18)$$

where

$$\kappa_f Q_f^2 = -M_f B_f Q_f^2 = \frac{m_{f_1} - \lambda_f^2 m_{f_0}}{\lambda_f^2 - 1}. \quad (19)$$

As will shortly be seen, experimentally $\lambda_f > 1$ and $\kappa_f > 0$. The mass spectrum (18) satisfies the simple recurrence formula

$$m_{f_{N+1}} = \lambda_f^2 m_{f_N} + (\lambda_f^2 - 1) \kappa_f Q_f^2 \quad (20)$$

with an inhomogeneous term proportional to Q_f^2 which shows explicitly that electric charge is here a "source" of particle mass. The spectrum (18) and the recurrence formula (20) were previously proposed on a phenomenological ground ⁵⁾.

In the case of leptons, Eq.(20), if fitted to the masses m_e, m_μ and $m_\tau = 1782_{-7}^{+2}$ MeV ⁶⁾, gives

$$\lambda_e = 3.992_{-0.009}^{+0.004}, \quad (\lambda_e^2 - 1) \kappa_e = 97.51_{-0.04}^{+0.01} \text{ MeV} \quad (21)$$

and then predicts ⁵⁾

$$m_{e_3} = 28.5_{-0.5}^{+0.2} \text{ GeV}, \quad m_{e_4} = 455_{-10}^{+3} \text{ GeV} \quad (22)$$

and also

$$m_{\nu_N} = 0 \quad (N > 0) \quad (23)$$

if $m_{\nu_e} = 0$.

In the case of quarks, if taking the small current masses $m_u \approx 0 \approx m_d$ (of a few MeV) and assuming that

$$m_u \ll \frac{4}{9} \kappa_u, \quad m_d \ll \frac{1}{9} \kappa_d, \quad (24)$$

one gets from Eq.(18)

$$m_{u_N} \approx \frac{4}{9} \kappa_u (\lambda_u^{2N} - 1), \quad m_{d_N} \approx \frac{1}{9} \kappa_d (\lambda_d^{2N} - 1) \quad (N > 0). \quad (25)$$

Thus, if

$$\lambda_u \approx \lambda_d, \quad \kappa_u \approx \kappa_d, \quad (26)$$

one concludes that $m_{u_N} \approx 4m_{d_N}$ and in particular

$$m_c \approx 4m_s \approx \frac{4}{9} (\lambda_u^2 - 1) \kappa_u, \quad m_t \approx 4m_b \approx \frac{4}{9} (\lambda_u^4 - 1) \kappa_u \quad (27)$$

(cf. the mass relations discussed on a different ground in Ref.7). Hence, if $m_c \approx 1.5$ GeV and $m_b \approx 5$ GeV, one obtains

$$m_g \approx 0.38 \text{ GeV}, \quad m_t \approx 20 \text{ GeV}, \quad (28)$$

predicting toponium $t\bar{t}$ at about 38 GeV ⁵⁾ as $m_{t\bar{t}} \approx 4m_{b\bar{b}} = 38$ GeV for $m_{b\bar{b}} = 9.5$ GeV. One also obtains

$$\lambda_u \approx 3.5, \quad (\lambda_u^2 - 1) \kappa_u \approx 3.4 \text{ GeV}. \quad (29)$$

Then from Eqs.(25) and (26) one predicts

$$m_{u_3} \approx 250 \text{ GeV}, \quad m_{d_3} \approx 62 \text{ GeV}. \quad (30)$$

One can verify a posteriori that the assumption (24) is satisfied for m_u very well, while for m_d with the factor 5 or so (for m_u and m_d equal to a few MeV).

Note that the mass spectrum (18) and its normalization condition

$\sum_N m_{fN} = M_f < +\infty$ confirm the necessity of a finite number of lepton and quark generations, $N \leq N_f < +\infty$, since $\lambda_f > 1$ (see Eqs.(21), (26) and (29)) and in the spectrum (18) there is a constant term. Explicitly, the normalization condition gives

$$(m_{f0} + \kappa_f Q_f^2) Z_f - \kappa_f Q_f^2 N_f = M_f. \quad (31)$$

It would be very natural to speculate ⁵⁾ that the upper bound for N should be caused by the gravitational cut for the magnitude of particle masses

$$m_{fN} \lesssim \sqrt{\frac{\hbar c}{2G}} = \frac{m_{PL}}{\sqrt{2}}, \quad (32)$$

where $m_{PL} = 1.2211027 \times 10^{19}$ GeV/c² is the Planck mass. In this case (excitingly enough!) our spectral formula (18) and the values (21) and (29) obtained for λ_f and κ_f by their fitting to experimental data give for leptons and quarks very close upper bounds

$$N_e = 18, \quad N_u = 19 \quad (33)$$

which become even identical,

$$N_e = N_u = 18, \quad (34)$$

if $\lambda_u \simeq 3.6$ instead of $\lambda_u \simeq 3.5$ (then $m_b \simeq 5.2$ GeV and $m_t \simeq 21$ GeV instead of $m_b \simeq 5$ GeV and $m_t \simeq 20$ GeV). However, it is well known that there exist various arguments for a much lower number of lepton and quark generations, among them the important perturbative argument ⁸⁾ based on one-loop corrections to the relation $m_W^2/m_Z^2 \cos^2 \theta_W = 1$ in the standard Glashow-Salam-Weinberg theory.

Let us remark that because of the rise of experimental mass spectrum m_{fN} with N (for $f \neq \nu$), the actual distribution $\rho_{fN} = m_{fN}/M_f$ also increases with N , showing that the probability of finding the component state N in the actual state ρ_f of f family rises with N . It does not imply, however, that heavy fermions should be more abundant in our present Universe than light

ones, as the former decay into the latter via the electroweak interactions (including Cabibbo-like mixing). These decays cannot be described on the level of the zeroth quantization (similarly as they cannot be on the level of the first quantization) and require the second quantization to be introduced (with particle masses consistent with the zeroth quantization considered as a zero-order approximation or an effective approach).

In conclusion, we would like to stress that the zeroth quantization provides a new mechanism of particle excitations which may be responsible for the generation of mass differences in the experimental known lepton and quark families. This quantum-statistical mechanism has nothing to do with the radial or orbital excitations connected with a hypohetic composite structure of leptons and quarks ⁹⁾.

ACKNOWLEDGMENTS

The author would like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste.

REFERENCES

- 1) Cf. e.g. P.A.M. Dirac, The Principles of Quantum Mechanics, 4th Edition (revised) (Clarendon Press, Oxford 1967), p.132.
- 2) Cf. e.g. H. Harari, Phys. Rep. C42, 235 (1978).
- 3) W. Królikowski, ICTP, Trieste, Internal Report IC/80/14.
- 4) P. Ramond, Phys. Rev. D3, 2415 (1971);
Cf. also W. Królikowski, Bull. Acad. Pol. Sci., Ser. Sci. Math. Astron. Phys. 20, 495 (1972).
- 5) W. Królikowski, Acta Phys. Polon. B10, 767 (1979); B10, 1061 (1979);
in Proc. of the International Symposium on Lepton and Hadron Interactions, Visegrád, Hungary, 1979 (to appear).
- 6) W. Bacino et al., Phys. Rev. Letters 41, 13 (1978).
- 7) G. Preparata, Phys. Letters 82B, 398 (1979).
- 8) M. Veltman, Nucl. Phys. B123, 89 (1977); Acta Phys. Polon. B8, 475 (1978).
- 9) Cf. e.g. W. Królikowski, Phys. Letters 90B, 241 (1980); ICTP, Trieste, preprint IC/80/1 (to appear in Acta Phys. Polon. B.).

CURRENT ICTP PREPRINTS AND INTERNAL REPORTS

- IC/79/98 INT.REP.* V.E. GODWIN and E. TOGATTI: Local field corrections to the binding energies of core excitons and shallow impurities in semiconductors.
- IC/79/99 INT.REP.* M.F. MOSTAFA, M.A. SEMARY and M.A. AHMED: Magnetic susceptibility investigation of some antiferromagnetic Fe^{2+} complexes.
- IC/79/100 INT.REP.* I.A. AMIN: The exchange property of modules.
- IC/79/101 E. MAHDAVI-HEZAVEH: Remarks concerning the running coupling constants and the unifying mass scale of grand unified gauge theories.
- IC/79/102 INT.REP.* ISMAIL A. AMIN: On a conjecture of Erdős.
- IC/79/103 M. YUSSOUFF and A. MOOKERJEE: Phonon frequency spectrum in random binary alloys.
- IC/79/104 INT.REP.* S. GOETTIG: Anisotropic plasmon-phonon modes in degenerate semiconductors.
- IC/79/105 T.N. SHERRY: Supersymmetric extension of the SU(5) model.
- IC/79/106 N.S. BAAKLINI: An SU(3) theory of electroweak interactions.
- IC/79/107 INT.REP.* M.Y.M. HASSAN and H.M.M. MANSOUR: Relativistic calculation of polarized nuclear matter.
- IC/79/108 INT.REP.* M.Y.M. HASSAN and S.S. MONTASSER: On the thermal properties of nuclear matter with neutron excess.
- IC/79/109 INT.REP.* J.S. NKOMA: Theory of absorption by exciton polaritons in a spatially dispersive media.
- IC/79/110 A. OSMAN: Nucleon-nucleon interaction in the three-nucleon system.
- IC/79/111 INT.REP.* S. FERRARA: Superspace aspects of supersymmetry and supergravity.
- IC/79/112 INT.REP.* M. STESLICKA and K. KEMPA: Variational calculation of surface states for a three-dimensional array of δ -function potentials.
- IC/79/113 INT.REP.* K.S. SINGWI and M.P. TOSI: Relation between bulk compressibility and surface energy of electron-hole liquids.
- IC/79/114 INT.REP.* A.R. HASSAN: Two-photon transitions to exciton polaritons.
- IC/79/115 INT.REP.* G. AKDENIZ and A.O. BARUT: Gauge-invariant formulation of dyonium Hamiltonian on the sphere S^3 .
- IC/79/117 INT.REP.* J.S. NKOMA: Linear photon and two photon absorption by surface polaritons.
- IC/79/118 INT.REP.* A. VISINESCU and A. CORCIOVEI: Dechanneling in the WKB approximation.
- IC/79/119 INT.REP.* M. APOSTOL: Finite size effects on the plasma frequency in layered electron gas.
- IC/79/120 INT.REP.* F. DESTEFANO and K. TAHIR SHAH: Quasi-catastrophes as a non-standard model and changes of topology.
- IC/79/121 A. OSMAN: Effect of Coulomb forces in the three-body problem with application to direct nuclear reactions.

- IC/79/122 INT.REP.* K. TAHIR SHAH: A note on the violation of angular-momentum conservation.
- IC/79/123 INT.REP.* A.A. EL SHAZLY, F.A. GANI and M.K. EL MOUSLY: Determination of optical absorption edge in amorphous thin films of selenium and selenium doped with sulphur.
- IC/79/124 INT.REP.* M.M. PANT: Variationally optimized muffin-tin potentials for band calculations.
- IC/79/125 INT.REP.* W. WADIA and L. BALLOOMAL: On the transformation of positive definite hermitian form to unit form.
- IC/79/126 INT.REP.* G.S. DUBEY and D.K. CHATURVEDI: Dynamical study of liquid aluminium.
- IC/79/127 A. NDUKA: Spherically symmetric cosmological solutions of the Lyttleton-Bondi Universe.
- IC/79/128 INT.REP.* M.M. BAKRI and H.M.M. MANSOUR: The relativistic two-fermion equations (I).
- IC/79/129 INT.REP.* N.S. TOHCHEV and J.G. BRANKOV: On the s-d model for coexistence of ferromagnetism and superconductivity.
- IC/79/130 INT.REP.* B.D. KANDILAROV, M.T. PRIMATAROVA and V. DETCHEVA: Interface states in a class of heterojunctions between diatomic semiconductors.
- IC/79/131 W. MECKLENBURG: Geometrical unification of gauge and Higgs fields.
- IC/79/132 LUNG CHI-WEI, SUN SHIANH-KAI and SHYURNG LJARNG-YUEH: The surface energy, thermal vibrations of dislocation lines and the critical crack.
- IC/79/133 INT.REP.* M.Y.M. HASSAN, S.S. MONTASSER and S. RAMADAN: On the thermal properties of polarized nuclear matter.
- IC/79/134 INT.REP.* A. OSMAN: Two-, three- and four-body correlations in nuclear matter.
- IC/79/135 T. BARNES and G.I. GHANDOUR: On quantizing gauge theories without constraints.
- IC/79/136 J. NIEDERLE: Supergravity.
- IC/79/137 INT.REP.* T.N. SHERRY: Higgs potential in the SU(5) model.
- IC/79/138 INT.REP.* M.M. BAKRI and H.M.M. MANSOUR: The relativistic two-fermion equations (II).
- IC/79/139 INT.REP.* R.R. BASILY: The effect of the gate electrode on the C-V characteristics of the structure M-TmF₃-SiO₂-Si.
- IC/79/140 J. NIEDERLE: Quantization as mapping and as deformation.
- IC/79/142 ABDUS SALAM: A gauge appreciation of developments in particle physics - 1979.
- IC/79/143 INT.REP.* B. BUTI: Ion-acoustic holes in a two-electron temperature plasma.
- IC/79/144 INT.REP.* W. KRÓLIKOWSKI: Recurrence formulae for lepton and quark generations.
- IC/79/145 INT.REP.* G. SENATORE, M. ROVERE, M. PARRINELLO and M.P. TOSI: Structure and thermodynamics of two-component classical plasmas in the mean spherical approximation.
- IC/79/146 INT.REP.* C. TOMÉ: Change of elastic constants induced by point defects in hcp crystals.
- IC/79/147 P. ŠTOVČEK and J. TOLAR: Quantum mechanics in a discrete space-time.
- IC/79/148 INT.REP.* M.T. TELI: Quaternionic form of unified Lorentz transformations.
- IC/79/149 INT.REP.* W. KRÓLIKOWSKI: Primordial quantum chromodynamics: QCD of a Fermi-Bose couple of coloured preons.
- IC/79/154 INT.REP.* RIAZUDDIN and FAYYAZUDDIN: A model for electroweak interactions based on the left-right symmetric gauge group U_L(2) ⊗ U_R(2).
- IC/79/158 INT.REP.* B. JULIA and J.F. LUCIANI: Non-linear realizations of compact and non-compact gauge groups.
- IC/79/159 INT.REP.* J. BOHÁČIK, P. LICHARD, A. NOGOVÁ and J. PIŠŮT: Monte Carlo quark-parton model and deep inelastic electroproduction.
- IC/79/162 P. BUDINI: Reflections and internal symmetry.
- IC/79/164 P. BUDINI: On conformal covariance of spinor field equations.
- IC/79/167 INT.REP.* T.D. PALEV: Para-Bose and para-Fermi operators as generators of orthosymplectic Lie superalgebras.

IC/80/5 RIAZUDDIN: Two-body D-meson decays in non-relativistic quark model.
INT.REP.*

IC/80/6 G. ALBERI, M. BLESZYNSKI, T. JAROSZEWICZ and S. SANTOS: Deuteron
D-wave and the non-eikonal effects in tensor asymmetries in elastic
proton-deuteron scattering.
INT.REP.*

IC/80/7 A.M. Kurbatov and D.P. Sankovich: On the one variational principle in
quantum statistical mechanics.
INT.REP.*

IC/80/8 G. Stratan: On the alpha decay branching ratios of nuclei around $A = 110$.
INT.REP.*

IC/80/12 M.V. MIHAILOVIĆ and M.A. NAGARAJAN: A proposal for calculating the
importance of exchange effects in rearrangement collisions.
INT.REP.*

IC/80/14 W. KROLIKOWSKI: Lepton and quark families as quantum-dynamical systems.
INT.REP.*

IC/80/17 NAMIK K. PAK: Introduction to instantons in Yang-Mills theory. (Part I).
INT.REP.*