



RESEARCH REPORT

IC/80/86  
INTERNAL REPORT  
(Limited distribution)

International Atomic Energy Agency  
and  
United Nations Educational Scientific and Cultural Organization

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

DENSITY OF STATES, POISSON'S FORMULA OF SUMMATION AND WALFISZ'S FORMULA \*

Pu Fucho \*\*

International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

Using Poisson's formula for summation, we obtain an expression for density of states of d-dimensional scalar Helmholtz's equation under various boundary conditions. Likewise, we also obtain formulas of Walfisz's type. It becomes evident that the formulas obtained by Pathria et al. in connection with ideal bosons in a finite system are exactly the same as those obtained by utilizing the formulas for density of states.

MIRAMARE - TRIESTE  
June 1980

\* To be submitted for publication.  
\*\* Permanent address: Institute of Physics, Academia Sinica, Beijing, People's Republic of China.

In order to study the statistical properties of a system of finite volume, it is necessary to know the density of states of the system. For a rectangular body satisfying the periodic boundary condition (PBC), Neumann boundary condition (NBC) and Dirichlet boundary condition (DBC) for dimensionality  $d = 1, 2, 3$ , the expressions for density of states are given in Refs.1 to 3. They are derived indirectly from a temporal coherent function of a finite black body. It had been pointed out <sup>2)</sup> that this expression can follow from Walfisz's formula of lattice summation which was known more than fifty years ago. It was clarified further in Ref.3 that the formulas for a finite system of ideal bosons obtained by Pathria et al. <sup>4)-6)</sup> with the help of Poisson's formula of summation are exactly the same as those derived from the formula for density of states. Also, the Poisson formula itself is nothing but the one-dimensional Walfisz formula. In this short communication we have obtained not only the general formula for density of states for any dimension  $d$  by using Poisson's formula of summation but also showed that Walfisz's type formulas can be readily derived from Poisson's formula. The method of Pathria et al. <sup>4)-6)</sup> is to apply Poisson's formula to the function to be summed, while in obtaining these formulas from that of density of states we must sum the delta function at the beginning. The two methods are naturally equivalent.

Starting from the corresponding one-dimensional Poisson formula, it is not difficult to show by induction that the d-dimensional Poisson equations are

$$\sum_{l_1, \dots, l_d = -\infty}^{\infty} f(l_1, \dots, l_d) = \sum_{n_1, \dots, n_d = -\infty}^{\infty} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} f(x_1, \dots, x_d) e^{2\pi i (n_1 x_1 + \dots + n_d x_d)} dx_1 \dots dx_d \quad (1)$$

$$\sum_{l_1, \dots, l_d = \frac{1}{2}(1-\eta)}^{\infty} f(l_1, \dots, l_d) = \left(\frac{\eta}{2}\right)^d f(0, \dots, 0) + \sum_{s=1}^d \left(\frac{\eta}{2}\right)^{d-s} \sum_{1 \leq j_1 < \dots < j_s \leq d} \sum_{n_{j_1}, \dots, n_{j_s} = -\infty}^{\infty} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} f(0, \dots, x_{j_1}, \dots, 0, \dots, x_{j_s}, \dots, 0) \exp \left\{ 2\pi i \sum_{p=1}^s n_{j_p} x_{j_p} \right\} \prod_p dx_{j_p} \quad (2)$$

The summation in the above equations are restricted to integers. In formula (2)  $n = \pm 1$  and when  $n = 1$  the sum at the left-hand side begins from zero, for  $n = -1$ , it begins from one.  $f(x_1, \dots, x_d)$  is a function of  $d$  variables. If

$$f(x_{p1}, \dots, x_{pd}) = f(x_1, \dots, x_d) \quad (3)$$

for all permutations  $P$  applying on  $(1, \dots, d)$  and

$$f(x_1, \dots, -x_j, \dots, x_d) = f(x_1, \dots, x_j, \dots, x_d) \quad (4)$$

then (2) becomes

$$\sum_{l_i, l_d = \frac{1}{2}(1-\eta)}^{\infty} f(l_1, \dots, l_d) = \left(\frac{\eta}{2}\right)^d f(0, \dots, 0) + \left(\frac{\eta}{2}\right)^d \sum_{s=1}^d \eta^s \binom{d}{s} \sum_{n_1, \dots, n_s = -\infty}^{\infty} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} f(x_1, \dots, x_s, 0, \dots, 0) e^{2\pi i(n_1 x_1 + \dots + n_s x_s)} dx_1 \dots dx_s \quad (5)$$

The scalar Helmholtz equation in the rectangular region  $L_1 \times \dots \times L_d$  is

$$\left\{ (\lambda_1^2 + \dots + \lambda_d^2) + k^2 \right\} \psi(x_1, \dots, x_d) = 0 \quad (6)$$

For PBC, the wave vector  $k$  is equal to

$$k = 2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}} \quad (7)$$

where  $l_1, \dots, l_d$  are integers. For

$$k = \pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}} \quad (8)$$

where  $l_1, \dots, l_d$  are integers from 0 to  $\infty$ . For DBC the same formula applies but  $l_1, \dots, l_d$  are integers from 1 to  $\infty$ .

For PBC, the density of states  $D_d(k)$  is, by definition, equal to

$$D_d(k) = \sum_{l_1, \dots, l_d}^{\infty} \delta(k - 2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}) \quad (9)$$

From (1) it follows that

$$D_d(k) = \sum_{n_1, \dots, n_d = -\infty}^{\infty} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \delta(k - 2\pi \sqrt{\frac{x_1^2}{L_1^2} + \dots + \frac{x_d^2}{L_d^2}}) \exp 2\pi i(n_1 x_1 + \dots + n_d x_d) dx_1 \dots dx_d \quad (10)$$

Introducing the  $d$ -dimensional spherical co-ordinate system  $(y, \theta_1, \dots, \theta_{d-1})$ ,  $0 \leq y < \infty$ ,  $0 \leq \theta_1 \leq \pi$ ,  $i = 1, \dots, d-2$ ,  $0 \leq \theta_{d-1} \leq 2\pi$  and taking  $\theta_1$  as the angle between vectors  $(x_1/L_1, \dots, x_d/L_d)$  and  $(L_1 n_1, \dots, L_d n_d)$  we have

$$D_d(k) = \frac{L_1 \dots L_d}{(2\pi)^d} k^{d-1} \Omega_{d-1} \sum_n \int_{-1}^1 e^{in y} (1-y^2)^{\frac{d-3}{2}} dy \quad (11)$$

where  $n = \sqrt{n_1^2 + \dots + n_d^2}$ .  $\sum_n$  represents that  $n_1, \dots, n_d$  are taken over all integers from  $-\infty$  to  $\infty$  and  $\Omega_{d-1}$  is the solid angle subtended by a  $(d-1)$  dimensional sphere. From the integral representation of the Bessel function, it gives

$$D_d(k) = \frac{L_1 \dots L_d}{(2\pi)^d} k^{d-1} \Omega_d \sum_n \Gamma\left(\frac{d}{2}\right) \frac{J_{\frac{d}{2}-1}(nk)}{(nk/2)^{d/2-1}} \quad (12)$$

which can be written as

$$= \frac{L_1 \dots L_d}{(2\pi)^d} k^{d-1} \Omega_d + \frac{L_1 \dots L_d}{(2\pi)^d} k^{d-1} \Omega_d \sum_n' \Gamma\left(\frac{d}{2}\right) \frac{J_{\frac{d}{2}-1}(nk)}{(nk/2)^{d/2-1}} \quad (13)$$

the first term is the Weyl term which forms the essential part and the second term is the oscillatory part. The meaning attached to each of them is evident.

For NBC and DBC, the density of states  $D^{\eta}(k)$  is, by definition, equal to

$$D_d^{\eta}(k) = \sum_{l_1, \dots, l_d = \frac{1}{2}(1-\eta)}^{\infty} \delta(k - \pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}), \quad (14)$$

while for NBC,  $\eta = 1$  and for DBC,  $\eta = -1$ .

Because conditions (3) and (4) are satisfied, we may apply Eq.(5) to obtain

$$D_d^{\eta}(k) = \left(\frac{1}{2}\right)^d \left\{ \delta(k) + \sum_{s=1}^d \eta^s \binom{d}{s} D_s(k; 2L_1, \dots, 2L_s) \right\}. \quad (15)$$

Thus the density of states for NBC and DBC can be expressed in terms of that for PBC. It should be noted that each side of the rectangular is doubled in the corresponding expression for PBC. This is denoted explicitly in Eq.(15). Substituting Eq.(12) in (15), we have

$$D_d^{\eta}(k) = \left(\frac{1}{2}\right)^d \left\{ \delta(k) + \sum_{s=1}^d \eta^s \binom{d}{s} \frac{L_1 \dots L_s}{\pi^s} k^{s-1} \Omega_s \sum_n \Gamma\left(\frac{s}{2}\right) \frac{J_{\frac{s}{2}-1}(2nk)}{(nk)^{s/2-1}} \right\}. \quad (16)$$

When  $d = 1, 2, 3$  the results in Refs.1 to 3 are obtained from Eqs.(16) and (12).

Now, we calculate the total number of states with the magnitude of a vector less than  $k$  in wave vector space. For PBC, it is given by definition that

$$N_d(k) = \sum_{l_1, \dots, l_d = -\infty}^{\infty} \theta(k - 2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}), \quad (17)$$

where  $\theta$  is the usual step function defined by

$$\theta(k) = \begin{cases} 1 & k > 0 \\ 0 & k \leq 0 \end{cases}.$$

At  $k = 0$ ,  $\theta(k)$  can be either defined as  $\frac{1}{2}$ , 1 or 0 with the understanding that the states situated at the boundary are counted  $\frac{1}{2}$ , counted or not counted, respectively.

Repeating the calculations, we have

$$N_d(k) = \frac{L_1 \dots L_d}{(2\pi)^d} \Omega_{d-1} \sum_n \int_0^k \frac{\Gamma(1/2) \Gamma((d-1)/2)}{(nw)^{d/2-1}} 2^{d/2-1} J_{d/2-1}(nw) w^{d-1} dw. \quad (18)$$

Utilizing the formula for Bessel functions

$$\frac{d}{dy} y^{\nu} J_{\nu}(y) = y^{\nu} J_{\nu-1}(y)$$

and completing the integral with respect to  $w$ , we obtain

$$N_d(k) = \frac{L_1 \dots L_d}{(2\pi)^d} V_d \sum_n \Gamma\left(\frac{d}{2} + 1\right) \frac{J_{d/2}(nk)}{(nk/2)^{d/2}}, \quad (19)$$

where  $V_d$  is the volume of a  $d$ -dimensional sphere of radius  $k$ . This is just the Walfisz formula. For NBC ( $\eta = 1$ ) and DBC ( $\eta = -1$ ), the number of states  $N_d^{\eta}(k)$  with magnitude less than  $k$  is given by

$$N_d^{\eta}(k) = \sum_{l_1, \dots, l_d = \frac{1}{2}(1-\eta)}^{\infty} \theta(k - \pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}). \quad (20)$$

Similarly we have

$$\begin{aligned} N_d^{\eta}(k) &= \left(\frac{1}{2}\right)^d \left\{ \theta(k) + \sum_{s=1}^d \eta^s \binom{d}{s} N_d(k; 2L_1, \dots, 2L_s) \right\} \\ &= \left(\frac{1}{2}\right)^d \left\{ \theta(k) + \sum_{s=1}^d \eta^s \binom{d}{s} \frac{L_1 \dots L_s}{\pi^s} V_d \sum_n \Gamma\left(\frac{s}{2} + 1\right) \frac{J_{s/2}(2nk)}{(nk)^{s/2}} \right\}. \quad (21) \end{aligned}$$

These are formulas of the Walfisz type. From the point of view of the lattice summation, it can be interpreted as the number of lattice points in the first quadrant in a sphere with radius  $k$  in wave number space. For  $n = 1$ , the points on the co-ordinate plane is taken into account in the first quadrant. For  $n = -1$ , it is not taken into account. The term  $\theta(k)$  comes from the contribution of the vertex, terms with  $s = 1, 2, \dots$  come from the boundary line, boundary plane. From definitions, it follows that

$$\frac{d}{dk} N_d(k) = D_d(k)$$

and

$$\frac{d}{dk} N_d^n(k) = D_d^n(k)$$

which may be verified readily from Eqs.(12), (19), (16) and (21).

The technique <sup>4)-6)</sup> used by Pathria et al. is to apply Poisson's summation formula directly to the function

$$f(k) = f\left(2\pi \sqrt{\frac{k_1^2}{L_1^2} + \dots + \frac{k_d^2}{L_d^2}}\right)$$

i. e.

$$\sum_{l_1, \dots, l_d} f\left(2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}\right) = \sum_{n_1, \dots, n_d} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} f\left(2\pi \sqrt{\frac{x_1^2}{L_1^2} + \dots + \frac{x_d^2}{L_d^2}}\right) \exp\left\{2\pi i (n_1 x_1 + \dots + n_d x_d)\right\} dx_1 \dots dx_d$$

Repeating the calculations performed above, we have

$$\sum_{l_1, \dots, l_d} f\left(2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}\right) = \int_0^{\infty} f(k) \frac{L_1 \dots L_d}{(2\pi)^d} k^{d-1} \Omega_d \sum_n \Gamma\left(\frac{d}{2}\right) \frac{J_{d/2-1}(nk)}{(nk/2)^{d/2-1}} dk \quad (22)$$

This result is certainly the same as that obtained from

$$\sum_{l_1, \dots, l_d} f\left(2\pi \sqrt{\frac{l_1^2}{L_1^2} + \dots + \frac{l_d^2}{L_d^2}}\right) = \int_0^{\infty} f(k) D_d(k) dk$$

by substituting into it the  $D_d(k)$  given by Eq.(12). While in obtaining Eq.(12) for  $D_d(k)$ , the Poisson formula is applied to the delta function.

#### ACKNOWLEDGMENTS

The author would like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste. He would also like to thank Professor M.P. Tosi for reading the manuscript.

#### REFERENCES

- 1) B. Steinhilber, H.P. Baltes and M. Pabst, Phys. Rev. A12, 1519 (1975).
- 2) H.P. Baltes and B. Steinhilber, J. Math. Phys. 18, 1275 (1977).
- 3) A.N. Chaba, Phys. Rev. A30, 1292 (1979).
- 4) S. Greenspoon and R.K. Pathria, Phys. Rev. A9, 2103 (1974).
- 5) C.S. Zasada and R.K. Pathria, Phys. Rev. A15, 2439 (1977).
- 6) A.N. Chaba and R.K. Pathria, Phys. Rev. A18, 1277 (1978).
- 7) A. Walfisz, Math. Z. 19, 1275 (1977).

- IC/79/163 G. GELMINI: The s-wave  $Q\bar{q}q$  states.
- IC/79/165 S. MARCULESCU and L. MEZINCESCU: Phase factors and point-splitting in supersymmetry.
- IC/79/166 A. FASOLINO, G. SANTORO and E. TOSATTI: Instability, distortion and dynamics of the  $W(100)$  surface.
- IC/79/168 T.D. PALEEV: The trace formulae for realizations of Lie algebras with A-operators. Analysis of the Fock representations.
- IC/80/1 W. KRÓLIKOWSKI: Towards a dynamical preon model.
- IC/80/5 RIAZUDDIN: Two-body D-meson decays in non-relativistic quark model.  
INT.REP.\*
- IC/80/6 G. ALBERI, M. BLESZYNSKI, T. JAROSZEWICZ and S. SANTOS: Deuteron D-wave and the non-eikonal effects in tensor asymmetries in elastic proton-deuteron scattering.  
INT.REP.\*
- IC/80/7 A.M. KURBATOV and D.P. SANKOVICH: On the one variational principle in quantum statistical mechanics.  
INT.REP.\*
- IC/80/8 G. STRATAN: On the alpha decay branching ratios of nuclei around  $A = 110$ .  
INT.REP.\*
- IC/80/9 N.Y. AYOUB: Effect of high lying states on the ground and few low lying excited  $0^+$  energy levels of some closed-shell nuclei.
- IC/80/10 L. FONDA, N. MANKOĆ-BORSTNIK and M. ROSINA: The decay of coherent rotational states subject to random quantum measurements.
- IC/80/11 G.C. GIRARDI, V. GORINI and G. PARRAVICINI: Spatial localization of quantum states and physical meaning of the matrix elements of the resolvent operator.
- IC/80/12 M.V. MIHAILOVIĆ and M.A. NAGARAJAN: A proposal for calculating the importance of exchange effects in rearrangement collisions.  
INT.REP.\*
- IC/80/13 A. BULGAC, F. CARSTOIU and O. DUMITRESCU: Double folded Yukawa interaction potential between two heavy ions.
- IC/80/14 W. KRÓLIKOWSKI: Lepton and quark families as quantum-dynamical systems.  
INT.REP.\*
- IC/80/16 AHMED OSMAN: Four-body problem for four bound alpha particles in  $^{16}\text{O}$ .
- IC/80/17 NAMIK K. PAK: Introduction to instantons in Yang-Mills theory (Part I).  
INT.REP.\*
- IC/80/18 RIAZUDDIN: Neutral current weak interaction without electro-weak unification.  
INT.REP.\*
- IC/80/19 M.S. CRAIGIE, S. NARISON and RIAZUDDIN: An apparent inconsistency between the Dyson and renormalization group equations in QCD.
- IC/80/20 W.I. FURMAN and G. STRATAN: On alpha decay of some isomeric states in Po-Bi region.  
INT.REP.\*
- IC/80/23 P. ROZMEJ, J. DUDEK and W. NAZAREWICZ: Possible interplay between non-axial and hexadecapole degrees of freedom - An explanation for "enormously" large  $Q_4$ ?
- IC/80/24 REVISED.
- IC/80/25 G. MAIELLA: Path-integral measure and the Fermion-Bose equivalence in the Schwinger model.  
INT.REP.\*
- IC/80/26 M.S. CRAIGIE, S. NARISON and RIAZUDDIN: A critical analysis of the electromagnetic mass shift problem in QCD.
- IC/80/27 D.K. CHATURVEDI, U. MARINI BETTOLO MARCONI and M.P. TOBI: Mode-coupling theory of charge fluctuation spectrum in a binary ionic liquid.  
INT.REP.\*
- IC/80/28 ABDUS SALAM: Gauge unification of fundamental forces (Nobel lecture).
- IC/80/30 R. NOJAROV, E. NADJAKOV and V. ANTONOVA: High spin structure in a coupled bands model.
- IC/80/31 AHMED OSMAN: Rearrangement collisions between four identical particles as a four-body problem.
- IC/80/34 I.Zh. PETKOV and M.V. STOITSOV: On a generalization of the Thomas-Fermi method to finite Fermi systems.  
INT.REP.\*

- IC/80/37 A.N. ANTONOV, V.A. NIKOLAEV and I.Zh. PETKOV: Nucleon momentum and density distributions of nuclei.
- IC/80/40 W. KRÓLIKOWSKI: An integral transform of the Salpeter equation.  
INT.REP.\*
- IC/80/41 Z.A. KHAN: Elastic scattering of intermediate energy protons on  ${}^4\text{He}$  and  ${}^{12}\text{C}$ .  
INT.REP.\*
- IC/80/44 A. OSMAN: Two-nucleon transfer reactions with form factor models.
- IC/80/45 E.W. MIELKE: Towards exact solutions of the non-linear Heisenberg-Pauli-Weyl spinor equation.  
INT.REP.\*
- IC/80/46 H.S. CRAIGIE: Catastrophe theory and disorders of the family system.  
INT.REP.\*
- IC/80/53 R. BECK, M.V. MIHAILOVIĆ and M. POLJSAK: Calculation of nuclear reaction parameters with the generator co-ordinate method and their interpretation.
- IC/80/58 S. YOKSAM: Spatial variations of order parameter around Kondo impurity for  $T \leq T_c$ .  
INT.REP.\*
- IC/80/59 J.K.A. AMUZU: Sliding friction of some metallic glasses.  
INT.REP.\*
- IC/80/60 Zi-Zhao GAN and Guo-Zhen YANG: A theory of coherent propagation of light wave in semiconductors.
- IC/80/61 K.G. AKDENIZ and O. OGUZ: A new class of meronic solutions.  
INT.REP.\*
- IC/80/63 H.B. SINGH and H. HAUG: Optical absorption spectrum of highly excited direct gap semiconductors.  
INT.REP.\*
- IC/80/64 W. KLONOWSKI: Living protein macromolecule as a non-equilibrium metastable system.  
INT.REP.\*
- IC/80/65 H.G. RNIK: An approximation to the ground state of  $E \otimes e$  and  $T_2 \otimes T_2$  Jahn-Teller systems based on Judd's isolated exact solutions.  
INT.REP.\*
- IC/80/66 A.N. ERMILOV, A.N. KYREEV and A.M. KURBATOV: Spin glasses with non-Gaussian distributions. Frustration model.  
INT.REP.\*