


2083052.80



СООБЩЕНИЯ
ОБЪЕДИНЕННОГО
ИНСТИТУТА
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

E5-82-669

G.S. Popov

ASYMPTOTIC BEHAVIOUR
OF THE SCATTERING PHASE
FOR NON-TRAPPING METRICS

1982

1. INTRODUCTION

The aim of this article is to study the asymptotic behaviour of the scattering phase $s(\lambda)$ related to an elliptic second order formally self-adjoint operator H , defined either in \mathbb{R}^n or in an unbounded domain Ω with Dirichlet or Neumann boundary conditions. Recently, this problem was investigated by many authors. In ref.^{/4/} Buslaev announced a result about the asymptotic of $s(\lambda)$ as $\lambda \rightarrow \infty$ for differential operators in \mathbb{R}^n , as well as in the obstacle case with Dirichlet boundary conditions. The perturbed operator H , considered in ref.^{/4/}, has a principal symbol with constant coefficients and 0 is not an eigenvalue of H . Later, the same problem was studied by A.Majda and J.Ralston^{/14/}. They proved the existence of an asymptotic expansion and computed the first three coefficients when $s(\lambda)$ is the scattering phase of the Laplacian with Dirichlet boundary conditions on a convex obstacle, and when $s(\lambda)$ is the scattering phase of the Laplace-Beltrami operator for a non-trapping metric on \mathbb{R}^n , which is Euclidean in a neighbourhood of ∞ . The authors conjectured that the same asymptotic expansion holds for any non-trapping obstacle. This conjecture was proved in ref.^{/18/} by V.Petcov and the author for the Laplacian with Dirichlet or Neumann boundary conditions.

For the Schrödinger operator $H = -\Delta + V$, $V \in C_0^\infty(\mathbb{R}^n)$ and $n = 3$, the asymptotic behaviour of $s(\lambda)$ as $\lambda \rightarrow \infty$ was investigated by Colin de Verdiere^{/5/}, and for any n -odd by Guillopé^{/7/}. Recently, an asymptotic expansion of $s(\lambda)$ related to a first or second order elliptic operator on a Hermitian bundle over an odd dimensional Riemannian manifold was announced by V.Ivriij and M.Shubin^{/9/}.

In this paper, both cases, n -even and n -odd, are considered, as well as the case when $\lambda = 0$ is an eigenvalue of the perturbed operator H . The asymptotic behaviour of the scattering phase $s(\lambda)$ as $\lambda \rightarrow \infty$ is investigated for arbitrary second order elliptic, formally self-adjoint differential operators H in a domain $\Omega \subset \mathbb{R}^n$, satisfying a non-trapping condition and such that $H = -\Delta$ in a neighbourhood of ∞ . The self-adjoint extension of H in $L^2(\Omega)$ with Dirichlet or Neumann boundary conditions on $\partial\Omega$ when $\Omega \neq \mathbb{R}^n$ is bounded from below but it allows to have a finite number of non-positive eigenvalues in contrast to refs.^{/14,18/}. Therefore there is not always a good rate of local decay for $H_{ac}^{-1/2} \sin(tH_{ac}^{1/2})$, $t \rightarrow \infty$ essentially used in ref.^{/18/}

where H_{ac} is the absolutely continuous part of the operator H . In order to overcome this difficulty we study the asymptotic behaviour of the S -matrix at infinity.

Suppose K is a bounded domain in R^n with smooth boundary ∂K and $\Omega = R^n \setminus K$ or $\Omega = R^n$. Consider an elliptic, formally self-adjoint second order differential operator P in Ω with Dirichlet or Neumann boundary conditions on ∂K , when $\Omega \neq R^n$ and $P = -\Delta$ outside the ball $B_R = \{x; |x| \leq R\}$. Without loss of generality assume that P has the form $P = -\Delta_g + hD + V$,

where $V \in C_0^\infty(R^n)$, $hD = \sum_{j=1}^n h_j(x) D_j$, $D_j = -i\partial/\partial x_j$ and Δ_g is the Laplace-Beltrami operator for a Riemannian metric g .

$$\Delta_g = \sum_{i,j=1}^n g^{-1/2} \partial/\partial x_i (g^{ij} g^{1/2}) \partial/\partial x_j$$

$g^{ij} \in C^\infty(\bar{\Omega})$, $g = \det(g_{ij})$, $g_{ij} = (g^{ij})^{-1}$ and $g_{ij} = \delta_{ij}$ for $|x| > R$. The projections of the (generalized) bicharacteristics of P on $\bar{\Omega}$ are called (generalized) geodesics of g .

Definition. The metric g is said to be non-trapping if there is $T > 0$ such that every (generalized) geodesic, beginning in B_R , leaves the ball B_R by the time T_R .

Let H_0 and H be the self-adjoint extension of the free Laplacian $-\Delta$ in $L^2(R^n)$ and of P in $L^2(\Omega)$ with Dirichlet or Neumann boundary conditions on ∂K when $\Omega \neq R^n$. These operators generate groups of unitary operators $\exp(itH_0)$ and $\exp(itH) \otimes 1$ in $L^2(R^n) = L^2(\Omega) \oplus L^2(K)$. The wave operators W_\pm are defined as follows

$$W_\pm = s\text{-}\lim_{t \rightarrow \mp \infty} (e^{itH} \otimes 1) e^{-itH_0}$$

It is well known^{3/} that W_\pm are isometrics on $L^2(R^n)$ and $\text{Rang}(W_+) = \text{Rang}(W_-)$, so the scattering operator $S = W_+^* W_-$ exists as a unitary operator on $L^2(R^n)$. In the spectral representation of H_0 on $L^2(R^+, L^2(S^{n-1}))$ the scattering operator S can be considered as a function of unitary operators $S(\lambda)$ on $L^2(S^{n-1})$ which is called a scattering matrix. Moreover, $S(\lambda) = I + K(\lambda)$, where $K(\lambda)$ is a trace class operator for $\lambda > 0$. This enables us to define the function $\det S(\lambda) : R^+ \rightarrow S^1 = \{z \in C; |z| = 1\}$ as a product of the eigenvalues of $S(\lambda)$. It was proved in refs.^{7,10,11/} that there exists a continuous (even analytic) in R^+ function $s(\lambda)$, satisfying the equality

$$\det S(\lambda) = \exp(2\pi i s(\lambda)), \quad \lambda > 0.$$

Such a function $s(\lambda)$ is called a scattering phase.

We shall prove the following results.

Theorem 1. Suppose the metric g is non-trapping in R^n . Then

$$s(\lambda) \sim \sum_{j=0}^{\infty} a_j \lambda^{n/2-j} \quad \text{as } \lambda \rightarrow \infty. \quad (1.1)$$

Moreover,

$$a_0 = (4\pi)^{-n/2} (\Gamma(n/2+1))^{-1} \{ \text{vol}_g(B_R) - \text{vol}_e(B_R) \},$$

$$a_1 = (4\pi)^{-n/2} (\Gamma(n/2))^{-1} \int \left(\frac{K\sqrt{g}}{3} - \frac{|h|_g}{4} + V(x) \right) dx,$$

where $\text{vol}_g(B_R)$ and $\text{vol}_e(B_R)$ are the Riemannian and Euclidean volume of the ball B_R , $K(x)$ is the scalar curvature and $|h|_g = \left(\sum_{i=1}^n g_{ij} h^i h^j \right)^{1/2}$ is the Riemannian length of the vector h .

In the case of the Schrödinger operator we prove

Theorem 2. Let $H = -\Delta + V$, $V \in C^\infty(R^n)$, $n \geq 3$. Then $s(\lambda)$ has the form (1.1) when $\lambda \rightarrow \infty$, where

$$a_j = \int P_j^n(V, DV, \dots, D^\alpha V) dx, \quad D^\alpha = D_1^{\alpha_1} \dots D_n^{\alpha_n}$$

and P_j^n are some universal polynomials. Moreover $P_0^n = 0$, $P_1^n(V) = (4\pi)^{-n/2} (\Gamma(n/2))^{-1} V$ and

$$P_j^n(\lambda V, \lambda^{3/2} DV, \dots, \lambda^{1+|\alpha|/2} D^\alpha V) = \lambda^j P_j^n(V, DV, \dots, D^\alpha V), \quad \lambda > 0.$$

In the obstacle case we prove

Theorem 3. Let $H = -\Delta_g + hD + V$ in $L^2(\Omega)$ with Dirichlet or Neumann boundary conditions on $\partial\Omega$ and suppose the metric g is non-trapping in $\Omega \subset R^n$, $n \geq 3$. Then

$$s(\lambda) \sim \sum_{j=0}^{\infty} a_j \lambda^{(n-j)/2} \quad \text{as } \lambda \rightarrow \infty$$

and

$$a_0 = (4\pi)^{-n/2} (\Gamma(n/2+1))^{-1} \{ \text{vol}_g(\Omega \cap B_R) - \text{vol}_e(B_R) \},$$

$$a_1 = \pm \frac{1}{4} (4\pi)^{-(n-1)/2} \left(\Gamma\left(\frac{n+1}{2}\right) \right)^{-1} \text{vol}_g(\partial\Omega),$$

where $+(-)$ sign is used in the case of Dirichlet (Neumann)

boundary conditions and $\text{vol}_g(\partial\Omega)$ is the Riemannian volume of $\partial\Omega$.

The plan of the paper is as follows. In section 1 we prove that the point spectrum of the operator H is finite and investigate some properties of the scattering phase. In section 2 we study the behaviour of the scattering matrix at ∞ in order to find functions $s_1(\lambda)$ and $s_2(\lambda)$ such that $s(\lambda^2) = s_1(\lambda) + s_2(\lambda)$ and $s_1(\lambda) \in C^\infty(\mathbb{R}^1)$, $\hat{s}_2(\lambda) = O(\lambda^{-N})$, $N \in \mathbb{Z}$, $\lambda \rightarrow \infty$ for any $N \in \mathbb{Z}$. In section 3 we investigate the distribution $\hat{s}_1(\lambda)$ using suitable trace formulas and prove a similar to theorem 1 result in the case of matrices of first order differential operators.

2. THE SCATTERING PHASE AND THE SPECTRUM OF H

We begin to study the spectrum of H in $L^2(\mathbb{R}^n)$. First we prove that the point spectrum of H is finite. Since $H = -\Delta$ outside the ball B_R the Rellich's theorem and the unique continuation property of second order elliptic operators yield the absence of the positive point spectrum of H . Moreover, $H \geq -\epsilon \Delta + V_1$ for some $\epsilon > 0$, $V_1 \in C_0^\infty(\mathbb{R}^n)$ when $\Omega = \mathbb{R}^n$ and since the negative point spectrum of $-\epsilon \Delta + V_1$ is finite, so is those of H . In the case $\Omega \neq \mathbb{R}^n$ we use the inequality $H \geq H_1 \oplus H_2$ in $L^2(\Omega \cap B_R) \oplus L^2(\mathbb{R}^n \setminus B_R)$, where $H_1 = H$ in $\Omega \cap B_R$, $H_2 = -\Delta$ in $\mathbb{R}^n \setminus B_R$ with Dirichlet boundary conditions on $\partial(\Omega \cap B_R)$ and ∂B_R respectively. Notice that both operators H_1 and H_2 have finite negative point spectrum. Moreover, the eigenvalue 0 has a finite multiplicity. Indeed, assume there exist infinitely many $\phi_j \in L^2(\mathbb{R}^n)$, $(\phi_j, \phi_k) = \delta_{jk}$ such that $H\phi_j = 0$. Then $\Delta\phi_j = (\Delta - H)\phi_j$ and $\phi_j(x) = \int |x-y|^{-n+2} (\Delta - H)\phi_j(y) dy$. Let $n > 4$, $\chi \in C_0^\infty(\mathbb{R}^n)$, $\chi = 0$ on B_R and $\chi(x) = 1$ for $|x| > R+1$. Then $|\chi\phi_j(x)| \leq C|x|^{-n+2}$ and $\|(\chi + |x|^2)^\epsilon \phi_j\|_{H^2(\mathbb{R}^n)} \leq C$ for some $\epsilon > 0$, $C > 0$. Now it is not hard to choose a Cauchy subsequence of ϕ_j in $L^2(\mathbb{R}^n)$, which contradicts our assumption. When $n=3$ or $n=4$,

we have $\int (\Delta - H)\phi_j(y) dy = 0$ since $|\xi|^{-2} (\Delta - H)\phi_j(\xi) = \hat{\phi}_j(\xi) \in L^2(\mathbb{R}^n)$. Then $\phi_j(x) = \int (\Delta - H)\phi_j(y) [|x-y|^{-n+2} |x|^{-n+2}] dy$ and the arguments given in the case $n > 4$ can be repeated. Therefore the point spectrum $\sigma_p(H)$ of H is finite and non-positive. Moreover, the continuous spectrum of H is absolutely continuous and coincides with \mathbb{R}^+ .

In the rest of this section we study the scattering phase $s(\lambda)$ related to the pair H, H_0 . First consider $K_0 = (a + H_0)^{-1}$, $K = (a + H)^{-1}$ which are bounded, self-adjoint operators for $a > \inf \sigma_p(H), 0 = \lambda_1$. Moreover, the operator $K^p - K_0^p$ is a trace class one for $p > n$ (see ref. [2]). Then the scattering phase $s(\lambda; K^p, K_0^p)$ related to the pair K^p, K_0^p is defined as follows

$$s(\lambda; K^p, K_0^p) = \pi^{-1} \lim_{\epsilon \rightarrow +0} \arg \det [1 + (K^p - K_0^p)(K_0^p - \lambda - i\epsilon)^{-1}], \lambda \in \mathbb{R}^1 \quad (2.1)$$

and has the properties (see refs. ^{/3,7,10/})

- (i) $s(\lambda; K^P, K_0^P) \in L^1(\mathbb{R}^n)$ and $\text{supp } s \subset [0, (\lambda_1 + a)^{-1}]$,
 (ii) For any $\Phi \in C^\infty(\mathbb{R}^n)$ the operator $\Phi(K^P) - \Phi(K_0^P)$ is a trace class one and

$$\text{Tr} \{ \Phi(K^P) - \Phi(K_0^P) \} = \int \Phi'(\lambda) s(\lambda; K^P, K_0^P) d\lambda,$$

$$(iii) \det S(\lambda; K^P, K_0^P) = \exp(-2\pi i s(\lambda; K^P, K_0^P)), \lambda > 0,$$

where $S(\lambda; K^P, K_0^P)$ is the scattering matrix for the pair H, H_0 . The function $s(\lambda) = s((a+\lambda)^{-1}; K^P, K_0^P)$ will be called a scattering phase for the pair H, H_0 . This notion is motivated by the property (iii) $\det S(\lambda) = \exp(-2\pi i s(\lambda))$ derived from (iii)' by the invariance principle. Using (i)' and (ii)' it is not hard to see, that (i) $(1 + \lambda^2)^{-p} s(\lambda) \in L^2(\mathbb{R}^1)$ for $p > n$ and $\text{supp } s \subset [a_1, \infty)$. (ii) For any $\Phi \in \mathcal{S}(\mathbb{R}^1)$ the operator $\Phi(H) - \Phi(H_0)$ is a trace class one and

$$\text{Tr} \{ \Phi(H) - \Phi(H_0) \} = \int \Phi'(\lambda) s(\lambda) d\lambda.$$

Moreover, the function $s(\lambda)$ is analytic in \mathbb{R}^+ since the operator H has no positive point spectrum (see refs. ^{/10,18/}).

Two special choices of the function Φ in (ii) are very useful for studying the asymptotics of $s(\lambda)$ at infinity. Let $\Phi(\lambda) = e^{-\lambda} \phi(\lambda)$, $\phi \in C^\infty(\mathbb{R}^1)$, $\phi(\lambda) = 1$ for $\lambda \in [-a, \infty)$ and $\phi(\lambda) = 0$ for $\lambda \in (-\infty, -a-1)$. Then $\Phi \in \mathcal{S}(\mathbb{R}^1)$ and

$$\text{Tr} \{ e^{-iH} \otimes 0 - e^{-iH_0} \} = -t \int_{-\infty}^{\infty} e^{-\lambda} s(\lambda) d\lambda, \quad t > 0, \quad (2.2)$$

where $e^{-iH} \otimes 0$ acts as e^{-iH} in $L^2(\Omega)$ and as 0 in $L^2(\mathbb{R}^n \setminus \Omega)$. Denote $\Phi(\lambda) = \phi(\lambda) \int \cos(\sqrt{\lambda} t) \rho(t) dt$, $\rho \in C_0^\infty(\mathbb{R}^1)$ and $B_0 = \sqrt{H_0}$, $B_1 = \sqrt{H_{ac}} \otimes i\sqrt{-H_p}$, where H_{ac} and H_p are respectively the absolutely continuous and discrete part of H . Then $\Phi \in \mathcal{S}(\mathbb{R}^1)$ and it is not hard to see from (ii), that

$$\text{Tr} \int_{-\infty}^{\infty} \rho(t) \{ \cos B_1 t \otimes 0 - \cos B_0 t \} dt = 1/2 \int_{\Gamma} \frac{d}{d\mu} \hat{\rho}(\mu) \tilde{s}(\mu) d\mu, \quad (2.3)$$

where $\Gamma = (-\infty, \infty) \cup (-ia, ia)$ and $\tilde{s}(\mu) = s(\mu^2)$ for $\mu \in (0, \infty)$ ($ia, 0$), $\tilde{s}(\mu) = -s(\mu^2)$ for $\mu \in (-\infty, 0) \cup (0, ia)$.

Remark. It turns out that the function $\tilde{s}(\mu)$, $\mu \in \mathbb{R}^1$ is the scattering phase for the wave equation in Lax-Phillips scattering theory (see ref. ^{/18/}). Moreover, using (2.1) one can obtain the equality (see refs. ^{/5,7/})

$$1/2 \int_{-ia}^{ia} \frac{d}{d\mu} \hat{\rho}(\mu) \tilde{s}(\mu) d\mu = \sum_{\lambda_j \in \sigma_p(H) \setminus \{0\}} \frac{\hat{\rho}(\sqrt{\lambda_j}) + \hat{\rho}(-\sqrt{\lambda_j})}{2} s(\lambda_j).$$

3. DECOMPOSITION OF $s(\lambda)$

In this section we construct functions $s_j(\lambda)$, $j=1,2$ with the properties

- (i) $s(\lambda^2) = s_1(\lambda) + s_2(\lambda)$,
- (ii) $|s_2(\lambda)| \leq C_N(1+\lambda)^{-N}$ when $\lambda \rightarrow \infty$, $N \in \mathbb{Z}$,
- (iii) The Fourier transform of $s_1(\lambda)$ is a compactly supported distribution.

To do this we use the equality

$$-\frac{d}{d\lambda} s(\lambda) = \text{Tr} \{ S(\lambda) \frac{d}{d\lambda} S^*(\lambda) \}, \quad \lambda > 0,$$

as well as an explicit form of the scattering matrix. We are going to obtain a representation formula for the S -matrix. By the invariance principle we have $S(\lambda) = S((a+\lambda)^{-1}; K^p, K_0^p)$, $\lambda > 0$. Moreover, the stationary approach^{/1,11,12/} can be applied to derive a representation formula for the S -matrix of the pair K, K_0 . Denote by A the operator of multiplication by $(1+|x|^2)^{-\beta/2}$, $\beta > n$ and let C be the operator given by $K = K_0 + ACA$. Since H coincides with H outside the ball B_R , the operator C is a compact one from H^{0,m_1} to H^{0,m_2} for every $m_1, m_2 \in \mathbb{R}$. Hereafter $H^{s,m}$ will be the weighted Sobolev space with norm

$$\|f\|_{s,m}^2 = \int (1+|\xi|^2)^s |\mathcal{F}[(1+|x|^2)^{m/2} f](\xi)|^2 d\xi$$

and \mathcal{F} stands for the Fourier transform $\mathcal{F}(f)(\xi) = \int e^{ix\xi} f(x) dx$. The operator $Q_0(\zeta) = A(K_0 - \zeta)^{-1}A$ has the norm-continuous boundary values $Q_0^\pm(\mu)$ for $\mu \in I = (0, a^{-1})$ as $\zeta \rightarrow \mu \pm i0$. Moreover, the compact operator $CQ_0^\pm(\mu)$ has no eigenvalue 1 in $L^2(\mathbb{R}^n)$ since H has no positive point spectrum (see ref.^{/11/} §7). Following Agmon, Kato, Kuroda^{/1,11,12/} one can prove that $Q^\pm(\mu) = \lim_{\zeta \rightarrow \mu \pm i0} A(K - \zeta)^{-1}A$ exists as a continuous function of operators bounded in $L^2(\mathbb{R}^n)$ for $\mu \in I$. Moreover $1 - CQ^\pm(\mu) = (1 + CQ_0^\pm(\mu))^{-1}$ for $\mu \in I$. The S -matrix for the pair K, K_0 can be written in the form

$$\begin{aligned} S(\mu; K, K_0) &= 1 - 2\pi i F_0(\mu) [1 + CQ_0^+(\mu)]^{-1} F_0(\mu)^* \\ &= 1 - 2\pi i F_0(\mu) [1 - CQ^+(\mu)] F_0(\mu)^*. \end{aligned} \tag{3.2}$$

The operator $F_0(\mu) : L^2(\mathbb{R}^n) \rightarrow L^2(S^{n-1})$ is determined by the equality $F_0(\mu) F_0^*(\mu) = -(2\pi i)^{-1} [Q_0^+(\mu) - Q_0^-(\mu)]$. Denote by $\gamma(\lambda)$ the trace operator on the sphere with a radius λ , $(\gamma(\lambda)u)(\omega) = u(\lambda\omega)$, $\omega \in S^{n-1}$ for $u \in C^\infty(\mathbb{R}^n)$, where polar coordinates $\xi = \rho\omega$ are used. The operator $\gamma(\lambda)$ extends to a Hölder continuity with respect to λ function of bounded operators from $H^{s,m}(\mathbb{R}^n)$ to $L^2(S^{n-1})$ for any $s > 1/2$, $m \in \mathbb{R}$. Using the equality $(K_0 - \zeta)^{-1} =$

$= -(1+z) - (1+z)^2 (H_0 - z)^{-1}$, $\zeta = (a+z)^{-1}$ and the Hölder continuity of $\gamma(\lambda)$ we obtain $E_0(\mu) = 2^{-1/2} (1+\lambda) \lambda^{(n-1)/4} \sigma(\lambda^{1/2}) \mathcal{F}A$, $\mu = (a+\lambda)^{-1}$. Then (3.2) and the invariance principle yield

$$S(\lambda) = 1 - \pi i (1+\lambda)^2 \lambda^{(n-1)/2} G(\lambda) [1 + (1+\lambda)V + (1+\lambda)^2 VR(\lambda^2 + i0)] VG^*(\lambda) \quad (3.3)$$

for $\lambda > 0$, where $G(\lambda) = \gamma(\lambda^{1/2}) \mathcal{F}$, $V = K - K_0$ and $R(z) = (H - z)^{-1}$.

Remark 1. In the case $\Omega = \mathbb{R}^n$ a more simple formula than (3.3) is known

$$S(\lambda) = 1 - \pi i \lambda^{(n-2)/2} G(\lambda) [V - VR(\lambda + i0)V] G^*(\lambda), \quad (3.4)$$

where $V = H - H_0$ (see ref. /18/). This formula is also valid when H and H_0 are matrices of differential operators and $G(\lambda)$ is suitably chosen.

Lemma 1. The S -matrix has the form $S(\lambda^2) = S_1(\lambda) + S_2(\lambda)$, where

(1) $(\frac{d}{d\lambda})^j S_2(\lambda)$ is a trace class operator with norm

$$\|(\frac{d}{d\lambda})^j S_2(\lambda)\|_{Tr} \leq C_N (1+\lambda)^{-N}, \quad \lambda > 0, \quad N \in \mathbb{Z}, \quad j < n-1.$$

(2) $\hat{S}_1(t) = \int_0^\infty e^{it\lambda} S_1(\lambda) d\lambda$ has a compact support with respect to t .

Obviously Lemma 1 and (3.1) give together the desired decomposition of the scattering phase. In order to prove Lemma 1 we need the following assertion.

Lemma 2. The operator $V = K - K_0$ has the form $V = V_1 + V_2$ where the distribution kernel of V_1 is compactly supported and $V_2: H^{s, m_1} \rightarrow H^{s+N, m_2}$ is a bounded operator for each $s, m_1, m_2, N \in \mathbb{R}^1$. Moreover $\text{supp } V_2 \subset \mathbb{R}^n \setminus B_R$ for any $u \in H^{s, m_1}$.

Proof. Let $\phi \in C^\infty(\mathbb{R}^{n+1})$, $\phi(t, x) = 1$ for $|x| < t + R$, $\phi(t, x) = 0$ for $|x| > t + R + 1$. Choose $\chi \in C_0^\infty(\mathbb{R}^{n+1})$, $\chi = 1$ on B_R and $\psi \in C_0^\infty(\mathbb{R}^1)$, $\psi(t) = 1$ for $|t| < 1$, $\psi(t) = 0$ for $|t| > 2$. Using the finite propagation speed of $B_j^{-1} \text{ sint } B_j$ we obtain

$$\begin{aligned} V &= \int_0^\infty e^{-t} \psi(t) \phi \{ B_1^{-1} \text{ sint } B_1 \otimes 0 - B_0^{-1} \text{ sint } B_0 \} \phi dt + \\ &+ \int_0^\infty e^{-t} (1-\psi) \chi \{ B_1^{-1} \text{ sint } B_1 \otimes 0 - B_0^{-1} \text{ sint } B_0 \} \phi dt + \\ &+ \int_0^\infty e^{-t} (1-\psi)(1-\chi) \phi \{ B_1^{-1} \text{ sint } B_1 \otimes 0 - B_0^{-1} \text{ sint } B_0 \} \phi dt. \end{aligned}$$

Denote the third integral by V_2 and the sum of the first and

second one by V_1 . Obviously the distribution kernel of V_1 is compactly supported. Moreover $\text{supp } V_2 u \subset \text{supp}(1-\gamma) \subset \mathbb{R}^n \setminus B_R$. Integrating by parts in the third integral and taking into account the inequality $|x| \leq t + R$ on $\text{supp } \phi$ we claim that $V_2 : H^{s, m_1} \rightarrow H^{s+N, m_2}$ is a bounded operator.

Lemma 3. Suppose that the operators $W_j \in \mathcal{L}(L^2)$ have compactly supported distribution kernels. Let the metric g be non-trapping in $\bar{\Omega}$. Then the operator $Q(\lambda) = W_1 R(\lambda^2 + i0) W_2$ has the form $Q(\lambda) = Q_1(\lambda) + Q_2(\lambda)$, where

$$(i) \quad \|(d/d\lambda)^k Q_2(\lambda)\|_{\mathcal{L}(L^2(\Omega))} \leq C_N (1+\lambda)^{-N}, \quad \lambda \in \mathbb{R}^+, \quad N \in \mathbb{Z}, \quad k < n-1$$

$$(ii) \quad \hat{Q}_1(t) = \int_0^\infty e^{i\lambda t} Q_1(\lambda) d\lambda \quad \text{is compactly supported.}$$

Proof. Consider the operators $P_j(t) = B^{-1} \text{sint} B_j$, $j = 0, 1$. Obviously $P_0(t)$ and $P_1(t)$ solve the problems

$$\begin{aligned} (D_t^2 - D_0)P_0(t) &= 0 & (D_t^2 - H)P_1(t) &= 0 \\ P_0(0) &= 0, \quad P_{0t}(0) = I & P_1(0) &= 0, \quad P_{1t}(0) = I \\ & & BP_1(t) &= 0, \end{aligned}$$

where $Bu = u/\partial\Omega$ or $Bu = \frac{\partial u}{\partial n}$ and n is the outward normal to

$\partial\Omega$. Let $\chi \in C_0^\infty(\mathbb{R}^n)$, $\chi \equiv 1$ on $\text{supp}_{x,y} W_j(x,y)$, $j = 1, 2$ and $\chi(x) = 0$ for $x \notin B_{R_1}$ where $W_j(x,y)$ are the distribution kernels of W_j . Due to the non-trapping condition, there exist $T > 0$, such that every generalized null bicharacteristic of $D_t^2 - H$ passing over $\text{supp } \chi \cap \bar{\Omega}$ at $t=0$ lies for $|t| > T$ completely over the set $\mathbb{R}^n \setminus B_R$. Moreover the bicharacteristics of $D_t^2 - H$ are straight lines outside the ball B_R . The propagation of singularities for the distribution kernel $P_1(t, x, y)$ of $P_1(t)$ yield

$$\text{sign } \text{supp } P(t, x, y) \chi(y) \subset \{(t, x, y) : ||x| - t| < T\}, \quad T > R_1. \quad (3.5)$$

Choose a cut-off function $\xi \in C^\infty(\mathbb{R}^{n+1})$ such that $\xi \equiv 1$ on a neighbourhood of $\{(t, x) : ||x| - t| < T\}$, $\xi(t, x) = 0$ if $(t, x) \notin \{(t, x) : ||x| - t| < T + 1\}$ and suppose $\int_x \xi(t, x) = 0$ for $t \in \mathbb{R}^1$.

Consider the operators $P_{0\chi} = \chi P_0(t) \chi$, $P_{1\chi} = \chi P_1(t) \chi$, $E_0 = \xi P_{1\chi}(t) \chi$, $R_\chi(\lambda) = \chi R(\lambda) \chi$. Then we have

$$W_1 R(\lambda^2 + i0) W_2 = W_1 \{ \chi \hat{E}(\lambda) + [R_\chi(\lambda^2 + i0) - \chi \hat{E}_0(\lambda)] \} W_2.$$

It is easy to see that the operator $\chi E_0(t)$ has a compact support with respect to t .

So we need the following estimate

$$\|D_\lambda^j [R(\lambda^2 + i0) - \chi \hat{E}(\lambda)]\|_{\mathcal{L}(L(\Omega))} \leq C_N (1 + \lambda)^{-N}, \quad N \in \mathbb{Z}, j < n-1, \quad (3.6)$$

where $\mathcal{L}(L^2(\Omega))$ is the space of bounded operators from $L^2(\Omega)$ to $L^2(\Omega)$. A similar to (3.6) estimate was obtained by Vainberg^{/21/} and Rauch^{/20/}. Our proof of (3.6) is close to that given in ref.^{/20/} and we only shall sketch it.

Consider the operator $F(t) = [D_t^2 - H, \xi] P(t)\chi$, $F(t) \in \mathcal{L}(L^2(\Omega))$. It follows from (3.5) that the kernel $\tilde{F}(t, x, y)$ of $F(t)$ is a smooth function, $\text{supp } F \subset \{(t, x, y); T < |x| - t < T+1\}$ and $F^{(k)}(0) = 0$ for any $k \in \mathbb{Z}^+$, since $\xi \equiv 1$ on $\text{supp } \chi$. Moreover

$$\begin{aligned} (D_t^2 - H)E_0(t) &= F(t), \\ E_0(0) &= 0, \quad E_{0t}(0) = \chi, \quad BE_0(t) = 0, \end{aligned} \quad (3.7)$$

where $Bu = u/\partial\Omega$ or $Bu = \frac{\partial u}{\partial n}/\partial\Omega$. Let $\tilde{F}(t, x, y)$ be a smooth function in $R^1 \times R^n \times \Omega$ such that $\tilde{F} = F$ for $x \in \Omega$ and $F(t, x, y) = 0$ for $x \in K$, $t > 2T$. Consider the problem

$$\begin{aligned} (D^2 - H_0)W(t) &= \tilde{F}(t) \\ W(0) &= 0, \quad W_t(0) = 0. \end{aligned} \quad (3.8)$$

Choose $\psi \in C^\infty(R^n)$, $\psi \equiv 1$ on $\text{supp } \chi$, $\psi(x) = 1$ for $|x| > 2T$. From (3.7), (3.8) and Duhamel's formula we have

$$W(t) = E_0(t) \otimes 0 - P_0(t)\chi + \int_0^t P_0(t-s)(H_0 - H \otimes 0)(E_0(s) \otimes 0) ds \quad (3.9)$$

in $L^2(R^n)$,

$$E_0(t) = \psi W(t) + P_1(t)\chi + \int_0^t P_1(t-s)Q(s) ds \quad (3.10)$$

in $L^2(\Omega)$, where $Q(s) = (1 - \psi)F(s) + [H, \psi]W(s)$. Since χE_0 has a compact support with respect to t , we can choose $T > 0$ so that

$$\chi W(t) = P_{0\chi}(t) + \int_0^T P_{0\chi}(t-s)(H_0 - H \otimes 0)(E_0(s) \otimes 0) ds.$$

The local energy decay of the operator $P_0(t)$, i.e.,

$$\|D_t^j P_{0\chi}(t)\|_{\mathcal{L}(H^{-s}, s^s)} \leq C_{s,j} t^{-n}, \quad t > C$$

and the smoothness of the kernel of $W(t)$ yield the estimate

$$\|D_t^j \chi W(t)\|_{\mathcal{L}(H^{-s}, H^s)} \leq C_{s,j} (1+t)^{-n} \quad \text{for } t \in \bar{R}^+.$$

Using the equalities $\chi W^{(\ell)}(0) = 0$ for $\ell \in Z^+$, we obtain

$$\|D_{\lambda}^j \chi \hat{W}(\lambda)\|_{\mathcal{L}(H^{-s}, H^s)} \leq C_N (1+\lambda)^{-N} \quad \text{for } N \in Z^+, j < n-1.$$

Therefore

$$\|D_{\lambda}^j \hat{Q}(\lambda)\|_{\mathcal{L}(L^2(\Omega))} \leq C_N (1+\lambda)^{-N} \quad \text{for } N \in Z^+, j < n-1. \quad (3.11)$$

Moreover, the function $\hat{Q}(\lambda) = \int_0^{\infty} e^{ikt} Q(t) dt$ is analytic on the half-plane $\text{Im} k > 0$ with values in $\mathcal{L}(L^2(\Omega))$ and it has a C^{n-2} continuation on \mathbb{R} . Multiplying (3.10) by χ and taking a Fourier-Laplace transform with respect to t we get

$$\chi \hat{E}_0(k) - R_{\chi}(k^2) = R_{\chi}(k^2) \hat{Q}(k)$$

for $\text{Im} k \geq C$, C -sufficiently large. We can extend this equality in $\{k; \text{Im} k > 0, \text{Re} k > 0\}$ since the functions $R_{\chi}(k^2)$ and $\hat{Q}(k)$ are analytical in this region with values in $\mathcal{L}(L^2(\Omega))$. Using (3.11) we obtain

$$\|D_{\lambda}^j R_{\chi}(\lambda^2 + i0)\|_{\mathcal{L}(L^2(\Omega))} \leq C \lambda^p, \quad \lambda \geq \lambda_0, j < n-1, \quad (3.12)$$

for some p , and prove the estimate (3.6). So we complete the proof of Lemma 3.

We are ready to prove Lemma 1. Using Lemma 2 and Lemma 3 with $W_j = V_j$ we can write $S(\lambda^2)$ in the form $S(\lambda^2) = S_1(\lambda) + S_2(\lambda)$, where

$$S_1(\lambda) = 1 + \pi i (1 + \lambda^2) G(\lambda^2) \chi [V_1 + Q_1(\lambda)] \chi G^*(\lambda^2) \lambda^{n-2}.$$

$$\begin{aligned} S_2(\lambda) = & \pi i (1 + \lambda^2) \lambda^{n-2} \{ (1 + \lambda^2)^2 G(\lambda^2) Q_2(\lambda) G^*(\lambda^2) + \\ & + G(\lambda^2) [1 + (1 + \lambda^2) V_1 + (1 + \lambda^2)^2 V_1 R(\lambda + i0) \bullet 0] V_2 G^*(\lambda^2) + \\ & + G(\lambda^2) V_2 [1 + \lambda^2 + (1 + \lambda^2)^2 R(\lambda + i0) \bullet 0] V G^*(\lambda^2) \}. \end{aligned}$$

The operator $S_1(\lambda)$ satisfies the second condition of Lemma 1. Indeed, the operator $\hat{Q}_1(t)$ has a compact support with respect to t in view of Lemma 3 and so does $G(\lambda^2) \chi(t)$ with distribution kernel $\delta(t-x\omega) \chi(x)$, $\chi \in C_n^{\infty}(\mathbb{R}^D)$, $\omega \in S^{n-1}$. In what follows we shall prove that $S_2(\lambda)$ satisfies the first condition of Lemma 1.

1. First consider the operator $I_1(\lambda) = G(\lambda^2) Q_1(\lambda) G^*(\lambda^2)$. The kernel of $G(\lambda^2)$ is equal to $e^{i\lambda\omega x}$, therefore $I_1(\lambda)$ is an operator with smooth kernel $I_1(\lambda, \omega, \theta)$ and

$$\begin{aligned} |I_1(\lambda, \omega, \theta)| &= \left| \int e^{i\lambda\omega x} \chi(x) Q_2(\lambda) (e^{-i\lambda\theta y} \chi(y)) dx \right| \leq \\ &\leq C \|Q_2(\lambda)\|_{\mathcal{L}(L^2(\Omega))} \leq C_N (1+\lambda)^{-N}. \end{aligned}$$

Therefore $I_1(\lambda)$ is a trace class operator and

$$\|I_1(\lambda)\|_{\text{Tr}} \leq C_N(1+\lambda)^{-N}.$$

2. In order to estimate the other terms of $S_2(\lambda)$ we use the inequality

$$\|R(\lambda^2 + i0)\|_{\mathcal{L}(H^{0,n}, H^{0,-n})} \leq C(1+\lambda)^p \quad \text{for some } p. \quad (3.13)$$

This estimate was proved for $\chi R(\lambda^2 + i0) \chi$ (see (3.12)). To derive it for $R(\lambda^2 + i0) \chi$ consider the resolvent equation $R(\lambda^2 + i0) \chi = R_0(\lambda^2 + i0) \chi - R_0(\lambda^2 + i0)(H - H_0) \chi R(\lambda^2 + i0) \chi$. Using the inequality $\|D_x^\alpha R_0(\lambda^2 + i0)\|_{\mathcal{L}(H^{0,n}, H^{0,-n})} \leq C\lambda$ for $|\alpha| \leq 2$ we obtain (3.13) for $R(\lambda^2 + i0) \chi$ and repeating this argument we prove (3.13).

Consider the operator $I_2(\lambda) = G(\lambda) V_1 R(\lambda^2 + i0) V_2 G^*(\lambda)$. This operator has a smooth kernel

$$I_2(\lambda, \omega, \theta) = (\lambda)^{-2N} \int e^{i\lambda x \omega} V_1 R(\lambda^2 + i0) (V_2 \Delta^N (e^{-i\lambda y \theta})) dx$$

and Lemma 2 yields $|I_2(\lambda, \omega, \theta)| \leq C\lambda^{-2N}$. The other terms of $S_2(\lambda)$ can be estimate in a similar way.

4. PROOF OF THE THEOREMS

In this section we show that the scattering phase has an asymptotic development at infinity and compute the coefficients. Denote by σ the distribution

$$\langle \sigma, \rho \rangle = \text{Tr} \int \rho(t) \{ \cos B_1 t \otimes 0 - \cos B_0 t \} dt, \quad \rho \in C_0^\infty(\mathbb{R}^1).$$

Using the trace formula (2.2) we have

$$\hat{\rho} \sigma(\lambda) = \frac{1}{2} \int_{-\infty}^{\infty} \frac{d}{d\mu} \hat{\rho}(\lambda - \mu) \tilde{s}(\mu) d\mu + \frac{1}{2} \int_{-i\alpha}^{i\alpha} \frac{d}{d\mu} \hat{\rho}(\lambda - \mu) \tilde{s}(\mu) d\mu$$

and the second integral is $O(\lambda^{-N})$ for any N as $\lambda \rightarrow \infty$. The decomposition of $s(\lambda^2)$ obtained in Sect.2 yields

$$\hat{\rho} * \frac{d}{d\lambda} s_3(\lambda) = -2\hat{\rho} \sigma(\lambda) + O(\lambda^{-N})$$

with $s_3(\lambda) = s_1(\lambda)$ for $\lambda > 0$ and $s_3(\lambda) = -s_1(\lambda)$ for $\lambda < 0$. For $\rho \in C_0^\infty(\mathbb{R}^1)$ and $\rho \equiv 1$ on $\text{supp } \hat{s}_1(t)$ the last equality leads to

$$\frac{d}{d\lambda} s(\lambda^2) = -2\hat{\rho} \sigma(\lambda) + O(\lambda^{-N}), \quad N \in \mathbb{Z}^+, \quad \lambda > 0. \quad (4.1)$$

First consider the case $\Omega = \mathbb{R}^n$. To study the right-hand side of (4.1), introduce the distribution kernels $v_0(t, x, y)$ and $v_1(t, x, y)$ of the operators $\cos B_0 t$ and $\cos B_1 t$. Obviously v_0 and v_1 are solutions of the problems

$$\begin{aligned} (D_t^2 - H_0)v_0 &= 0 & (D_t^2 - H)v_1 &= 0 \\ v_0|_{t=0} &= \delta(x-y), \quad v_0|_{t=0} = 0 & v_1|_{t=0} &= \delta(x-y), \quad v_1|_{t=0} = 0 \end{aligned}$$

and $\sigma(t)$ is equal to the distribution $\int [v_1(t, x, x) - v_0(t, x, x)] dx$. Repeating the arguments in the proof of Corollary 1.2 in ref. 6, one can prove that $\text{sing supp } \sigma \subset \{T_j, T_j\}$ is a period of a periodic geodesic of g . Since the non-trapping condition $\text{sing supp } \sigma = \{0\}$. Then (4.1) holds for any $\rho \in C_0^\infty(\mathbb{R}^1)$, $\rho \equiv 1$ on a neighbourhood of $t=0$. Using the finite speed of propagation and applying a finite partition of unity, one can reduce the problem to the investigation of the functions

$$I_j(\lambda) = \iint e^{-i\lambda t} \rho(t) \phi(x) v_j(t, x, x) dx dt, \quad j=0,1,$$

with $\phi \in C_0^\infty(\mathbb{R}^n)$. It turns out that for $|t| < \delta$ and δ sufficiently small, the distributions v_1 and v_0 are sums of oscillating integrals

$$v_\pm(t, x, y) = \int e^{i\Phi_\pm(t, x, y, \theta)} a^\pm(t, x, y, \theta) d\theta, \quad (4.2)$$

where a^\pm are classical amplitudes, $a^\pm \sim \sum_{j=0}^\infty c_j^\pm$, c_j^\pm - homogeneous of order $-j$ with respect to θ . The phase functions Φ_\pm have the form $\Phi_\pm = \psi(x, y, \theta) \pm \text{tg}(y, \theta)$ (see 6), q^2 is the principle symbol of H and ψ is a local solution of $q(x, y, \theta) = 0$ when $\langle x-y, \theta \rangle = 0$ and $d_x \psi(x, y, \theta) = 0$ for $x=y$. Then the integral $I_1(\lambda)$ became

$$I_1(\lambda) = \lambda^n \int e^{i\lambda t(1-q(y, \theta))} \phi(x) \rho(t) a^+(t, x, x, \theta) d\theta dx + O(\lambda^{-N}).$$

Substituting

$$\theta = r\omega \quad q(x, \omega) = 1, \quad r > 0, \quad S = \{\omega; q(x, \omega) = 1\}$$

we have

$$I(\lambda) = \lambda^n \int_0^\infty \int_S e^{i\lambda t(1-r)} \rho(t) \phi(x) a^+(t, x, x, \lambda r \omega) |Vq|^{-1} dS dr dx dt,$$

and applying the method of stationary phase we obtain

$$I_1(\lambda) \sim (2\pi)^{-n} \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \frac{\lambda^{n-j-k-1}}{k!} \iint_S (i^{-1} \frac{\partial^2}{\partial t \partial r}) [r^{n-1-j} c_j^+(t, x, x, \omega)]_{r=0} \phi(x) \frac{dS}{Vq} dx.$$

This formula leads to (1.1) with

$$a_j = (2\pi)^{-n} \iint [c_j^+(0, x, x, \omega) + \sum_{k=1}^j (n-j-1)\dots(n-j-k)(k!)^{-1} (i^{-1} \partial/\partial t)^k c_{j-k}^+(0, x, x, \omega) \frac{dS dx}{|\nabla q|}]. \quad (4.3)$$

In order to compute the coefficients in the case of the Schrödinger operator $H = -\Delta + V$, observe that $\Phi_{\pm} = \langle x-y, \theta \rangle \pm t|\theta| \nabla c_1^+ = 1/2$, and c_ℓ , $\ell > 0$ solves the transport equation

$$\partial c_\ell^+ - \langle \theta, \nabla_x \rangle c_\ell^+ - i/2(\partial_t^2 - \Delta + V)C_{\ell-1}^+ = 0.$$

$$c_\ell^+ |_{t=0} = 0.$$

Using (4.3) we prove inductively that a_j has the form prescribed in theorem 2.

The investigation of the asymptotic behaviour of $\hat{\rho}\sigma(\lambda)$ as $\lambda \rightarrow \infty$ in the obstacle case $\Omega = \mathbb{R}^n$ for the Laplace operator with Dirichlet or Neumann boundary conditions was done in refs.^{8,18/}. It was proved, that $\hat{\rho}\sigma(\lambda) \sim \sum_{j=0}^{\infty} a_j \lambda^{(n-j-2)/2}$ and the first three coefficients a_j , $j=0,1,2$ were obtained in the case $a_j = \delta_{ij}$. The method used in refs.^{8,18/} by Ivrii, can be applied to the investigation of $\rho\sigma(\lambda)$ for arbitrary second order differential operators in Ω with Dirichlet or Neumann boundary conditions. In order to compute the first two coefficients of $s(\lambda)$ one can use the trace formula (2.2) as well as the asymptotics of the right-hand side of (2.2) as $t \rightarrow +0$ given by Mc Keen and Singer (see^{15/} §4 and §5, formula (2)). Comparing the coefficients of the two sides of (2.2) as $t \rightarrow +0$ we get a_0 and a_1 .

The method used in the previous sections can be applied without to study the asymptotics of the scattering phase related to systems of first order differential operators. Let $H_0 = \sum_{j=1}^n A_j^0 D_j$, $H_1 = \sum_{j=1}^n A_j^1(x) D_j + B(x)$ be self-adjoint operators in $L^2(\mathbb{R}^n; \mathbb{C}^{2m})$, A_j^0 be constant $2m \times 2m$ matrices, $A_j^1 \in C^\infty(\mathbb{R}^n; \mathbb{R}^{4m^2})$, $A^k = \sum_{j=1}^n A_j^k \xi_j$, $k=0,1$; $H_1 = H_0$ outside the ball B_R . Assume, that the eigenvalues $\lambda_j(x, \xi)$ of $A^1(x, \xi)$ are simple and

$$\lambda_1(x, \xi) < \dots < \lambda_m(x, \xi) < 0 < \lambda_{m+1}(x, \xi) < \dots < \lambda_{2m}(x, \xi). \quad (4.4)$$

Then the spectrum of H_0 is absolutely continuous and $\sigma(H_0) = \mathbb{R}^1$. The eigenfunctions of H_1 in L^2 corresponding to a non-zero eigenvalue are smooth and supported in B_R and so they are finitely many. Moreover, the eigenvalue $\lambda=0$ has a finite multiplicity. Thus $\sigma(H_1) = \sigma_p(H_1) \cup \sigma_{ac}(H_1)$ and $\sigma_p(H_1)$ is finite, $\sigma_{ac}(H_1) = \mathbb{R}^1$.

Consider the scattering phase $s(\lambda)$ related to the pair H_1, H_0 . The function $s(\lambda)$ has the properties (i)-(iii) described in Sect.2. Choosing $\Phi(\lambda) = \hat{\rho}(\lambda)$, $\rho \in C_0^\infty(\mathbb{R}^1)$ in (iii) we obtain the following trace formula

$$\text{Tr} \int \rho(t) \{ e^{itH} - e^{itH_0} \} dt = \int \frac{d}{d\lambda} \hat{\rho}(\lambda) s(\lambda) d\lambda. \quad (4.5)$$

Denote by $\rho_j(t)$, $1 \leq j \leq 2m$ the projections of the bicharacteristics of $\lambda_j(x, \xi)$ on the x -space. We shall use the following non-trapping condition. There exists $T > 0$, such that

$$\rho_j(t) \notin B_R \quad \text{for } t > T \quad \text{if} \quad \rho_j(0) \in B_R. \quad (4.6)$$

Theorem 4. Suppose that (4.4) and (4.6) are valid. Then

$$s(\lambda) \sim \sum_{j=0}^{\infty} a_j^\pm \lambda^{n-j} \quad \text{as} \quad \lambda \rightarrow \pm \infty$$

and

$$a_0^\pm = (4\pi)^{-n/2} (\Gamma(n/2+1))^{-1} \int \text{Tr}(\pi_j^\pm A^1(x, \xi) - \pi_0^\pm A^0(\xi)) dx d\xi,$$

where π_j^+ (π_j^-) is the projection on the positive (negative) eigenspace of A . The proof of theorem 4 is similar to that of theorem 1 and we shall only sketch it. In order to decompose $s(\lambda)$ as a sum of functions $s_j(\lambda)$, $j=1,2$ with the properties (ii) and (iii) described in Sect.3, we use the formula

$$S(\lambda) = 1 - 2\pi i G(\lambda) \{ V + VR(\lambda + i0)V \} G^*(\lambda),$$

where $\lambda \in \mathbb{R}^1$ ($\sigma_p(H_1) = 0$), $V = H_1 - H_0$. Here $G(\lambda)$ are bounded operators from $H^{\ell, s}$, $s > 1/2$, $\ell \in \mathbb{R}^1$ to an auxiliary space \mathcal{H} . Denote by $\pi_j(\xi)$ the orthogonal projection onto the eigenspace of $A^0(\xi)$ corresponding to $\lambda_j(\xi)$. Then $\pi_j(\xi)$ is a smooth, homogeneous function of order one in \mathbb{R}^n . Let $S_{j, \lambda} = \{ \xi \in \mathbb{R}^n; \lambda_j(\xi) = \lambda \}$ and $d\mu_j(\omega) = |\lambda_j(\xi)|^{-1} dS_j$, where dS_j is the usual Lebesgue measure on $S_{j, \lambda}$. Consider the trace operators $\gamma_j(\lambda)$ on $S_{j, \lambda}$ defined by $(\gamma_j(\lambda) u)(\omega) = u(\lambda\omega)$, $u \in C^\infty(\mathbb{R}^n)$, where polar coordinates $\xi = \lambda\omega$, $\omega \in S_{j, 1}$ are used. Denote $\gamma_\lambda = \sum_{j=1}^{2m} \gamma_j(\lambda) \pi_j(\xi)$ and $\mathcal{H} = \sum_{j=1}^{2m} \pi_j(\lambda\omega) L^2(S_{j, \lambda}; d\mu_j; \mathbb{C}^{2m})$. It turns out that $G_\lambda = \gamma_\lambda \mathcal{F}$. Moreover, the operator $\int e^{i\lambda t} G_\lambda V d\lambda$ has a compactly supported distribution kernel and using an analogy of Lemma 2 we find the functions $s_j(\lambda)$ $j=1,2$. From (4.5) we obtain $\frac{d}{d\lambda} s(\lambda) = -\hat{\rho} \sigma(\lambda) + O(\lambda^{-N})$, where $\sigma(t) = \int [u_1(t, x, x) - u_0(t, x, x)] dx$ and u_j are the fundamental solutions of the Cauchy problem for $D_t - H_1$ and $D_t - H_0$ respectively. Using a microlocal

parametrix for the Cauchy problem and the method of the stationary phase we complete the proof of theorem 4.

REFERENCES

1. Agmon S. Spectral Properties of Schrödinger Operators and Scattering Theory. Ann. Scuola Norm. Sup. Pisa, 1975, 11, p.151-218.
2. Birman M.S. Vestnik Leningrad Univ., 1962, 1, p.22-25.
3. Birman M.S., Krein M.G. Dokl.Akad.Nauk SSSR, 1962, 144, p.475-478.
4. Buslaev V.S. Dokl.Akad.Nauk SSSR, 1981, 197, p.999-1002.
5. Colin de Verdiere Y. Une formule de trace pour l'operateur de Schrödinger dans R^3 . Ann.Scient.Es.Norm.Sup., 1981, 14, p.27-49.
6. Duistermaat J., Guillemin V. Invent Math., 1975, 29, p.39-79.
7. Guillipé L. Une formule de trace pour l'operateur de Schrödinger dans R^3 , thèse. Universite Scient. et Medicale Grenoble, 1981.
8. Ivrij V. Dunc.Anal. i Pril., 1980, 14, p.23-34.
9. Ivrij V., Schubín M.S. Dokl.Akad.Nauk SSSR, 1982, p.282-284.
10. Jensen A., Kato T. Comm. in P.D.E., 1978, 3, p.1165-1195.
11. Kato T. Hadronic Journ., 1978, 1, p.134-154.
12. Krein M. Dokl.Akad.Nauk SSSR, 1962, 144, p.268-271.
13. Kuroda An Introduction to Scattering Theory Lecture Notes. Springer Verlag, 1980, p.51.
14. Majda A., Ralston J. Duke Math.J., 1978, 45, p.183,196, p.517-536; 1979, 46, p.725-731.
15. McKean H., Singer I. J.Diff.Geometry, 1967, 1, p.43-69.
16. Melrose R., Sjöstrand J. Comm.Pure Appl.Math. 1978, 31, p.593-617.
17. Petkov V., Popov G. C.R.Acad.Sc.Paris, 1981, 292, p.275-277.
18. Petkov V., Popov G. Ann.Inst. Fourier, 1982, 32, p.123-162.
19. Popov G. Dokl.BAN, 1982, 51, p.168-172.
20. Rauch J. Comm. Pure Appl.Math., 1978, 31, p.431-480.
21. Vainberg B. Russian Math.Surveys, 1975, 30, p.1-38.


Received by Publishing Department
on September 14 1982.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices - in US \$,
including the packing and registered postage

D13-11807	Proceedings of the III International Meeting on Proportional and Drift Chambers. Dubna, 1978.	14.00
	Proceedings of the VI All-Union Conference on Charged Particle Accelerators. Dubna, 1978. 2 volumes.	25.00
D1,2-12450	Proceedings of the XII International School on High Energy Physics for Young Scientists. Bulgaria, Primorsko, 1978.	18.00
D-12965	The Proceedings of the International School on the Problems of Charged Particle Accelerators for Young Scientists. Minsk, 1979.	8.00
D11-80-13	The Proceedings of the International Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics. Dubna, 1979.	8.00
D4-80-271	The Proceedings of the International Symposium on Few Particle Problems in Nuclear Physics. Dubna, 1979.	8.50
D4-80-385	The Proceedings of the International School on Nuclear Structure. Alushta, 1980.	10.00
	Proceedings of the VII All-Union Conference on Charged Particle Accelerators. Dubna, 1980. 2 volumes.	25.00
D4-80-572	N.N.Kolesnikov et al. "The Energies and Half-Lives for the α - and β -Decays of Transfermium Elements"	10.00
D2-81-543	Proceedings of the VI International Conference on the Problems of Quantum Field Theory. Alushta, 1981	9.50
D10,11-81-622	Proceedings of the International Meeting on Problems of Mathematical Simulation in Nuclear Physics Researches. Dubna, 1980	9.00
D1,2-81-728	Proceedings of the VI International Seminar on High Energy Physics Problems. Dubna, 1981.	9.50
D17-81-758	Proceedings of the II International Symposium on Selected Problems in Statistical Mechanics. Dubna, 1981.	15.50
D1,2-82-27	Proceedings of the International Symposium on Polarization Phenomena in High Energy Physics. Dubna, 1981.	9.00

Orders for the above-mentioned books can be sent at the address:
Publishing Department, JINR
Head Post Office, P.O.Box 79 101000 Moscow, USSR



Попов Г.С.

E5-82-669

Асимптотическое поведение фазы рассеяния в незахватывающих метриках

Рассмотрено асимптотическое поведение фазы рассеяния на бесконечности для эллиптического самосопряженного дифференциального оператора H либо в \mathbb{R}^n , либо в области $\Omega \subset \mathbb{R}^n$ с краевыми условиями Дирихле или Неймана. Оператор H имеет вид $H = -\Delta_g + hD + V(x)$, где Δ_g - оператор Лапласа-Бельтрами римановой метрики g в Ω , $hD = \sum_{j=1}^n h_j D_j$, $D_j = -\frac{1}{i} \frac{\partial}{\partial x_j}$. Предполагается, что H равен оператору Лапласа Δ в окрестности бесконечности и что метрика g не является ловушкой для лучей, т.е. все геодезические метрики уходят с любого компакта в $\bar{\Omega}$ через некоторое время, зависящее только от компакта. При этих ограничениях получено полное асимптотическое разложение фазы рассеяния $s(\lambda)$ для $\lambda \rightarrow \infty$. Найдены первые члены этого разложения.

Работа выполнена в Лаборатории вычислительной техники и автоматизации ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1982

Popov G.S.

E5-82-669

Asymptotic Behaviour of the Scattering Phase for Non-Trapping Metrics

The asymptotic behaviour of the scattering phase is considered at infinity for an elliptic, self-adjoint, second order differential operator H , defined either in \mathbb{R}^n or in an unbounded domain $\Omega \subset \mathbb{R}^n$ with Dirichlet or Neumann boundary conditions. The operator H has the form $H = -\Delta_g + hD + V$, where Δ_g is the Laplace-Beltrami operator related to a Riemannian metric g

in $\bar{\Omega}$, $hD = \sum_{j=1}^n h_j D_j$, $D_j = -\frac{1}{i} \frac{\partial}{\partial x_j}$. Provided a non-trapping hypothesis

is fulfilled and H coincides with the Laplace operator Δ in a neighbourhood of infinity, an asymptotic development of the scattering phase $s(\lambda)$ is obtained as $\lambda \rightarrow \infty$. The first coefficients in this development are found.

The investigation has been performed at the Laboratory of Computing Techniques and Automation, JINR.

Communication of the Joint Institute for Nuclear Research. Dubna 1982

Редактор Э.В.Ивашкевич. Макет Р.Д.Фоминой.
Набор В.С.Румянцевой, Н.И.Коротковой.

Подписано в печать 4.11.82.

Формат 60x90/16. Офсетная печать. Уч.-изд.листов 1,37.

Тираж 430. Заказ 32007.

Издательский отдел Объединенного института ядерных исследований.
Дубна Московской области.