


2

SU8505885



**ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА**

**E11-84-495**

**A.B.Shvachka**

**SYMMETRY PROPERTIES  
OF SOME NONLINEAR FIELD  
THEORY MODELS**

Submitted to XXII International Conference  
on High Energy Physics (Leipzig, July 1984)

**1984**

1. In this paper various approaches towards the study of symmetry properties of nonlinear equations as well as possible ways of their computer implementation using the language of systems of analytical computations are discussed. A detailed investigation on the problems met in this paper may be found in <sup>/1-3/</sup>. In each section we deal with the possibilities of automation of routine calculations, connected with the study of symmetry properties of nonlinear field theory equations. Special attention is paid to the method of pseudopotential investigation of formal integrability and isovector method for the equations of balance.

2. The method of quasipotentials has been proposed by Wahlquist and Estabrook <sup>/4/</sup>. It allows one to write down for a nonlinear evolution equation a finite or infinite set of conservation laws as well as to find the Bäcklund transformation and the equations of associated linear scattering problem, by using the tools of outer differential forms. In order to illustrate the approach, consider the following <sup>/5/</sup> nonlinear 2nd order equation

$$H(\xi, \eta, u_{\xi\xi}, u_{\eta\eta}, u_{\xi\eta}, u_{\eta}, u_{\xi}, u) = 0. \quad (2.1)$$

A column vector is said to be a quasipotential if it satisfies the following overdetermined system of the 1st order differential equations

$$\begin{aligned} y_{\xi}^{\mu}(\xi, \eta) &= F^{\mu}(\xi, \eta, u, u_{\xi}, u_{\eta}, \dots, u^*, u_{\xi}^*, u_{\eta}^*, \dots, y^{\mu}, y^{\mu*}), \\ y_{\eta}^{\mu}(\xi, \eta) &= G^{\mu}(\xi, \eta, u, u_{\xi}, u_{\eta}, \dots, u^*, u_{\xi}^*, u_{\eta}^*, \dots, y^{\mu}, y^{\mu*}), \quad \mu = 1, 2, \dots, \eta. \end{aligned} \quad (2.2)$$

In this case the consistency conditions for the system (2.2), i.e.,

$$F_{\eta}^{\mu} = G_{\xi}^{\mu}, \quad (2.3)$$

are equivalent with the original equation (2.1).

Hence, the problem is reduced to that one of writing down equations (2.2) for a given equation (2.1) (i.e., to find  $F^{\mu}$  and  $G^{\mu}$ ) and, subsequently, of solving them relative to  $y^{\mu}(\xi, \eta)$ . The essence of the method consists of replacement of (2.1) by a closed Pfaffian system of differential forms which together with prolongation forms  $\theta^{\mu}$  form a differential ideal.

The condition

$$\Theta^\mu = \alpha dy^\mu - F^\mu d\xi - G^\mu d\eta \quad (2.4)$$

leads to a family of commutation relations between vector-fields. On the next step, it is necessary to find the vector-fields  $y^\mu$  in the explicit form of their commutation relations for components of vector-fields, and thereby also the functions  $F^\mu$  and  $G^\mu$ . The algorithm of finding quasipotentials requires cumbersome calculations. In order to automatize such calculations one can employ a system of analytical computations (say, REDUCE <sup>/7/</sup>). In ref. <sup>/7/</sup> the algorithm is described and the text of program is given on the language of REDUCE for performing the basic operations with outer differential forms. Our experience shown that the program works rather slowly. Therefore the author jointly with I.G.Resnikov and V.L.Topunov created a program on the language of a system of analytical calculations, which carries over the following steps on a computer: 1) replaces the original equation by a system of outer differential forms, 2) tests the Frobenius condition for a given equation, and 3) writes down the system of equations for the purpose of finding isovectors <sup>/7/</sup> and simplifies that system when possible. Consequently, the most part of routine work is carried out by a computer. In ref. <sup>/8/</sup> a description is given to the programs in REDUCE which construct prolongation structures for nonlinear equations. Unfortunately, we have no possibility of becoming familiar with that paper. In <sup>/6,9/</sup> REDUCE-programs are described for the study of Lie symmetries of differential equations.

3. A somewhat different, nevertheless conceptually close to WE <sup>/4/</sup> approach towards the study of symmetry properties for nonlinear equations is given in <sup>/10,11/</sup> In particular, in P.Kepsten's paper <sup>/11/</sup> the formalism of local jets <sup>/12/</sup> is employed. Similarly to the WE approach, the original equation is replaced by a family of outer differential forms. For the nonlinear Dirac equation

$$\sum_{k=1}^3 h \frac{\partial}{\partial x_k} (y_k \psi) - i h \frac{\partial}{\partial x_4} (y_4 \psi) + m_0 c \psi + n_0 \psi (\bar{\psi} \psi) = 0 \quad (3.1)$$

the class of generators of the local symmetry group is constructed in <sup>/11/</sup>. Using contact forms the ideal  $I$  of differential 1-forms, connected with (3.1), is constructed. The infinitesimal symmetries of the closed ideal  $I$  in the  $n$ -dimensional space formed by the class of differential forms  $a(I), \dots, a(m)$  are determined by vector-fields

$$V = V^i \frac{\partial}{\partial x_i} \quad (3.2)$$

such that  $\mathcal{L}_V |C|$ , where  $\mathcal{L}_V$  stands for the Lie derivative relative to the vector-field  $V$ . As is seen from (3.2), the vector-field  $V$  necessarily satisfies the conditions

$$\mathcal{L}_V a(i) + \gamma(i, j) \wedge a(j) = 0, \quad (i = 1, \dots, m), \quad (3.3)$$

where  $\gamma(i, j)$  are the corresponding differential forms. Beside the definition of the generators for the local symmetry group of the nonlinear Dirac equation, using the program package on the language of REDUCE in<sup>/2/</sup> there are determined also the preserved flows. The author jointly with I.G. Resnikov and V.L. Topunov carried over calculations and determined the generators of local symmetry groups for a number of the nonlinear equations. This required development of the corresponding program packet on the language of a system of analytical computations. In this way all routine calculations connected with the determination of generators and the construction of preserved flows are carried over with the aid of a computer<sup>/30/</sup>.

4. When investigating formal integrability of nonlinear evolution equations (NEE) one may successfully employ Computer Algebra Systems (CAS). In<sup>/13/</sup> there are described algorithms and programs on the language REDUCE designed for the study of formal integrability of NEE having the form

$$u_t = F(u, u_1, \dots, u_n), \quad n \geq 2, \quad u = u(x, t), \quad u_i = \frac{d^i u}{dx^i}. \quad (4.1)$$

A systematic account of questions related to formal integrability, existence of nontrivial Lie-Bäcklund algebra, and infinite series of conservation laws may be found in<sup>/2, 12/</sup>. Here we restrict ourselves to merely presenting basic concepts and necessary formulae<sup>/13/</sup>.

An equation of the form (4.1) is said to be formally integrable if there exists a formal series

$$L = \sum_{i=-\infty}^{\infty} a_i D^i, \quad a_i = a_i(u, u_1, \dots, u_{k_i}), \quad D \stackrel{\text{def}}{=} \frac{d}{dx},$$

satisfying on solutions to (4.1) the following operator relation

$$L_t - [F^*, L] = 0, \quad (4.2)$$

where  $F^* \stackrel{\text{def}}{=} \sum_{i=0}^n \frac{\partial F}{\partial u_i} D^i$ ,  $L_t \stackrel{\text{def}}{=} \left[ \frac{d}{dt}, L \right]$ .

The set of all functions  $H(u, u_1, \dots, u_m)$  such that

$$H_*(F) - F_*(H) = 0 \quad (4.3)$$

is said to be the Lie-Bäcklund algebra,  $A(F)$ , of the equation (4.1). A Lie-Bäcklund algebra is called nontrivial, if it contains elements different from  $u_1$  and  $F$ . A conservation law for (4.1) is the relation

$$\frac{d}{dt} p(u, u_1, \dots, u_k) = \frac{d}{dx} q(u, u_1, \dots, u_{k+n}) \quad (4.4)$$

valid on the solutions to (4.1). The order of conservation laws

is, by definition, the number  $\frac{1}{2} \text{ord} \frac{\delta p}{\delta u}$ , where  $\text{ord} f = n$  is defined

as such an  $n$  for which  $\frac{df}{du_n} \neq 0$ ,  $\frac{df}{du_m} = 0$ ,  $m > n$ ; and  $\frac{\delta}{\delta u} \stackrel{\text{def}}{=} \sum_{i=0}^k (-1)^i D^i \frac{\partial}{\partial u_i}$

is the variational derivative. If  $p \in \text{Im}(D)$  (i.e.,  $p$  coincides with the total derivative with respect to  $x$ ), then  $\delta p / \delta u = 0$  and the conservation law becomes trivial. Let us summarize basic results on formally integrable equations <sup>/14-16/</sup>.

Theorem 1. If the equation (4.1) admits for an infinite Lie-Bäcklund algebra, then it is formally integrable <sup>/14/</sup>.

Theorem 2. If the equation (4.1) admits for an infinite series of conservation laws of form (4.4), then it is formally integrable <sup>/15/</sup>.

Theorem 3. The equation (4.1) is formally integrable if and only if there exists an infinite series of conservation laws <sup>/16/</sup> of the following type:

$$(\text{Res } L^{-1})_i \in \text{Im } D, \quad (4.5a)$$

$$(\text{Res } L^{-1} L)_i \in \text{Im } D, \quad (4.5b)$$

$$(\text{Res } L^m)_i \in \text{Im } D, \quad m = 1, 2, \dots, \quad (4.5c)$$

where  $\text{Res } \sum b_i D^i \stackrel{\text{def}}{=} b_{-1}$ .

Conditions (4.5a) and (4.5b) may be rewritten in terms of the right-hand side  $F$  of (4.1)

$$\left[ \left( \frac{\partial F}{\partial u_n} \right)^{-1/n} \right]_i \in \text{Im } D, \quad (4.6a)$$

$$\left( \frac{\partial F / \partial u_{n-1}}{\partial F / \partial u_n} \right) \in \text{Im } D. \quad (4.6b)$$

In order to find explicit formulae for the coefficients of the series  $L$  and of densities  $\text{Res}L^m$  the following algorithm was proposed in <sup>/13/</sup>. It follows from (4.2) that the coefficients of the series  $L^m = \sum_{i=-\infty}^m n_i D^i$  should satisfy the relations

$$b_i = \frac{F_n^{i/m}}{n} \int dx \left\{ F_n^{-\frac{(i+n)}{n}} [c_{n+i-1} |_{b_i=0} + \partial_t b_{n+i-1}] \right\}, \quad (4.7)$$

$$i = m-1, m-2, \dots; m = 1, 2, \dots, F_n = \frac{\partial F}{\partial u_n},$$

where  $c_j$  are coefficients of  $D^j$  in the commutator  $[L^m, F_*]$ ,  $b_j = 0$  for  $j > m$ . The integration constants in (4.2) are taken to be zeroes.

Since the expressions  $c_{n+i-1}$  depend on  $b_j$ ,  $j \leq i$ , the relation (4.7) makes it possible to define an arbitrary number of coefficients  $b_j$  by recurrence. In particular, when  $i = -1$ , we may determine the explicit form of  $\text{Res} L^m$ .

The algorithm described so far enables us to solve the following problems <sup>/13/</sup>.

a) Testing the conditions for formal integrability. For any concrete equation of the form (4.1) one first has to check the conditions (4.6a) and (4.6b). If the result is positive, then the recurrent procedure (4.7) is employed and the quantities  $\text{Res}L^m$ ,  $m = 1, 2, 3, \dots$ , are calculated. If the original equation is formally integrable, then the integrands in (4.7) must be total derivatives with respect to  $x$ . Hence, an implicit checking of conditions (4.5c) is contained in the recurrent procedure: if for  $m = 1$  and on the  $i$ -th step a "nonintegrable" expression appears under the integral sign in (4.7), then the condition (4.5c) for  $m = 1 - n - i$  is not satisfied.

b) Inclusion of the requirement of formal integrability into the equations. When the proposed algorithm is applied to equations containing arbitrary constants, then these constants in general will appear in the left-hand sides of (4.6a), (4.6b), as well as in the integrand of (4.7), i.e., in expressions which for formally integrable equations should coincide with total derivatives. Hence, by taking the variational derivatives of these expressions equal to zero, we can resolve the resulting equations and in this way adjust the constants so that the original equation becomes formally integrable. The same results may be also obtained in a more simple manner, namely, by singling out the "nonintegrable" part of (4.7) and taking this equation to zero.

c) Finding the densities of conservation laws. Here it suffices to observe that for concrete equations of the form (4.1) the algorithm yields explicit formulae for  $\text{Res}L^m$ ,  $m = 1, 2, \dots$ , these being the densities we seek for.

d) Finding the elements of Lie-Bäcklund algebra. Calculations of  $b_m, b_{m-1}, \dots, b_0$  according to (4.7) give us the preliminary information for the determination of the  $m$ -th order element of the Lie-Bäcklund algebra for any given equation of the form (4.1). The corresponding algorithm is described in<sup>/16/</sup>. It allows either to find the element of a prescribed order or, to show that such an element does not exist.

A detailed description of programs developed by the author jointly with A.Yu.Zharkov is given in<sup>/13/</sup>. The programs are applied to S.I.Svinolupov-V.V.Sokolov equations

$$u_t = u_5 + 5(u_2 - u_1^2 + \alpha e^{2u} + \beta e^{-4u})u_3 - 5u_1 u_2^2 + 15(\alpha e^{2u} - 4\beta e^{-4u})^2 u_1 u_2 + u_1^5 + 90\beta e^{-4u} u_1^3 + 5(\alpha e^{2u} + \beta e^{-4u})u_1, \quad \alpha, \beta \in \mathbb{C}, \quad (4.8)$$

and result in a nontrivial element of the Lie-Bäcklund algebra in the form<sup>/13/</sup>

$$H = u_7 + 7(u_2 - u_1^2)u_5 + 14(u_3 - 2u_1 u_2)u_4 - 21u_1 u_2^2 - 14(2u_2^2 + u_1^2 u_2 - u_1^4)u_3 - \frac{28}{3}u_1 u_2^3 + 28u_1^3 u_2^2 - \frac{4}{3}u_1^7 + 7(\alpha e^{2u} + \beta e^{-4u})u_5 + 7(5\alpha e^{2u} - 16\beta e^{-4u})u_1 u_4 + 14(5\alpha e^{2u} - 13\beta e^{-4u})u_2 u_3 + 28(2\alpha e^{2u} + 29\beta e^{-4u})u_1^2 u_3 + 42(\alpha e^{2u} - 76\beta e^{-4u})u_1^3 u_2 + 14(7\alpha^2 e^{4u} - 13\alpha\beta e^{-2u} - 20\beta^2 e^{-8u})u_1 u_2 + 1260\beta e^{-4u} u_1^5 + 70(\alpha^2 e^{4u} + 2\alpha\beta e^{-2u} + 10\beta^2 e^{-8u})u_1^3 + \frac{28}{3}(\alpha e^{2u} + \beta e^{-4u})^3 u_1. \quad (4.9)$$

5. In<sup>/17,18/</sup> various concepts of group symmetry for differential equations are discussed. Since information concerning the group symmetry in many cases is equivalent to information concerning the solutions of a given equation, it appears reasonable to employ an as general concept of group symmetry as possible. We present the definition of the symmetry group proposed in<sup>/19,20/</sup>. Suppose the functions  $\phi(x_0, \dots, x_{n-1})$  on the  $n$ -dimensional space  $R^n(x)$  satisfy

$$A\phi = 0, \quad (5.1)$$

where  $A$  is a differential operator.

Definition 1. Let a family of operators  $g_L$  form the Lie algebra for some group  $G$ . The group  $G$  is said to be the symmetry group of the equation (5.1) if  $g_L$  and  $A$  commute on the set of solutions to (5.1):

$$[A, g_\alpha]\phi(x) = 0. \quad (5.2)$$

In other words, the generators of  $G$  map one solution,  $\phi(x)$ , into another one,  $\psi_\alpha(x) = g_\alpha\phi(x)$ .

A more general definition of symmetry groups for equations is given in /17/.

**Definition 2.** Let a family of operators  $g_a$  ( $1 \leq a < \infty$ ) form an algebra closed under multiplication ( $\cdot$ ) and generate some group G. The group G is said to be a symmetry group for  $A\phi=0$ , if

$$A \cdot A \cdot \dots \cdot A \cdot g_a \phi(x) = A^p g_a \cdot \phi(x) = 0 \quad (5.3)$$

(i.e., the multiplication  $p$  - yields an operator transforming a non-zero solution  $\phi(x)$  into the zero solution) or, if

$$\psi_a(x) = A^{p-1} \cdot g_a \phi(x) \quad (5.4)$$

(i.e., the operator  $g_a$  considered on the set of all solutions to  $A\phi=0$ , yields an operator transforming one non-zero solution into another one (in general, non-zero as well)).

The algebra of generators  $g_a$  ( $1 \leq a < \infty$ ) of the symmetry group G is said to be the invariance algebra of the equation  $A\phi=0$  /17/. It is clear that the second definition contains the first one as a particular case, namely, one for  $p=1$  and the multiplication ( $\cdot$ ) corresponds to commutation, i.e., when G is a Lie group.

6. The symmetry properties of multidimensional nonlinear differential equations make it possible, in some cases, to construct the solutions explicitly. In /21,22// classes of exact solutions are found using the symmetry properties for the following multidimensional nonlinear equations:

$$\square u + \lambda u^k = 0, \quad u = u(x), \quad x = (x_0, x_1, \dots, x_{n-1}), \quad (6.1)$$

$$\square u + \lambda \exp(u) = 0, \quad \square = -p_\mu p^\mu, \quad (6.2)$$

$$p^\mu = i g_{\mu\nu} \frac{\partial}{\partial x_\nu}; \quad (6.3)$$

$$\frac{\partial u}{\partial x_\mu} \cdot \frac{\partial u}{\partial x^\mu} = m^2, \quad [\gamma_\mu p^\mu - m - \lambda(\bar{\psi}\psi)^k] \psi = 0, \quad (6.4)$$

where  $\gamma_\mu$  is the Dirac matrix,  $\psi$  is a spinor and  $m, k, \lambda$  are arbitrary constants.

In order to find explicit solutions to equations (6.1) through (6.4) the following approach is used /21/:

$$u(x) = \phi(\omega) f(x) + g(x), \quad (6.5)$$

where  $\phi(\omega)$  is a certain unknown function (or a vector-function in case of a system of differential equations) depending on new

invariant variables  $\omega = \omega(\mathbf{x}) = \{\omega_1, \omega_2, \dots, \omega_{n-1}\}$ , the number of these being  $n-1$ , where  $n$  is the number of variables in the original equation. New variables  $\omega(\mathbf{x})$  and explicit expressions for  $f(\mathbf{x})$  and  $g(\mathbf{x})$  are determined from the system of Euler-Lagrange equations

$$\frac{dx_0}{\xi^0} = \frac{dx_1}{\xi^1} = \dots = \frac{dx_{n-1}}{\xi^{n-1}} = \frac{du}{\eta}, \quad (6.6)$$

where  $\xi^\mu$  and  $\eta$  are functions which determine the infinitesimal invariance group for (6.1)-(6.3), that is,

$$x'_\mu = x_\mu + \epsilon \xi^\mu(\mathbf{x}, u) + O(\epsilon^2), \quad u'(x') = u(\mathbf{x}) + \epsilon \eta(\mathbf{x}, u) + O(\epsilon^2). \quad (6.7)$$

Of course, the explicit form of  $\xi^\mu$  and  $\eta$  depends upon the concrete symmetry.

In order to solve the nonlinear Dirac equation (6.4) one uses /21/

$$\psi(\mathbf{x}) = A(\mathbf{x}) \phi(\omega), \quad (6.8)$$

where  $A(\mathbf{x})$  is a  $4 \times 4$  nonsingular matrix,  $\phi(\omega)$  is an unknown 4-component spinor depending merely on invariant variables. The explicit form of the matrix  $A$  is determined from the equation /21/

$$QA(\mathbf{x}) = (\xi^\mu(\mathbf{x}) \partial_\mu + \eta_x) A(\mathbf{x}) = 0, \quad (6.9)$$

(i.e.,  $Q$  is infinitesimal generator of the invariance group for (6.4)). Explicit forms of solutions to (6.4) are given in /23/. Besides the equations (6.1)-(6.4) the above explained method was employed by Fuschich and coauthors in order to find classes of exact solutions for multidimensional nonlinear Schrödinger equation /24/ Hamilton-Jacobi equation /21/ and Born-Infeld equation /25/, respectively. It was shown that:

1. If a nonlinear equation is in possession of nontrivial symmetry, there is a hope to determine multi-parameter families of its exact solutions.
2. The exact solutions may serve the purpose of "etalons" when developing constructive approximative methods for solution of nonlinear equations.

The method proposed in /21-25/ can be implemented on the language of systems of analytic computations. At present, the author jointly with I.G. Resnikov and V.L. Topunov are working on its computer realization.

7. When studying systems of differential equations in different areas of mathematics the necessity arises in checking their consistency <sup>/26/</sup>. This is usually connected with cumbersome symbolic calculations. One of the methods designed for this purpose is Cartan's algorithm <sup>/27/</sup>.

The possibility of using a computer for the above-mentioned problem is pointed out in <sup>/28/</sup>. In <sup>/29-37/</sup> the algorithm is described for the investigation on consistency properties of various systems of equations.

Suppose we are given a system of  $m$ -th order differential equations

$$(s) \Phi_{\mu}(x, u, p) = 0, \quad \mu = 1, 2, \dots, s, \quad (7.1)$$

where  $x = (x_1, \dots, x_n)$ ,  $u = (u_1, \dots, u_r)$ ,  $p$  are the derivatives up to some specified order  $m$  of the functions  $u$  relative to the variables  $x$ , i.e.,  $p_{\mu}^a = \frac{\partial^{(a)} u_{\mu}}{\partial x_a}$ ,  $a = (a_1, \dots, a_n)$ . In order to solve

the consistency problem for (7.1) it is necessary to clarify whether the given system is in involution, otherwise, it is necessary to construct its prolongation with the aid of adding

differential consequences  $\frac{d\Phi_{\mu}}{dx_i} = 0$ , where  $d/dx_i$  is the total

derivative with respect to  $x_i$ . The system (7.1) is in involution, if the following conditions are satisfied <sup>/35/</sup>:

1)  $c_{x_0}(s)$  is an involutory space;

2) there exist a neighbourhood  $\mathcal{G} \subset J^m$  of the point  $x_0 \in S$  such that the triple  $(r(s) \cap (p_m^{m+1})^{-1}(y), s, \mathcal{G}p_m^{m+1})$  is a fibred manifold,

where  $s$  is the manifold determined by  $\Phi_{\mu} = 0$ ,  $\frac{d\Phi_{\mu}}{dx_i} = 0$ ,  $J^m$  is the set of all  $m$ -jets  $j_x^m$ , and  $p_m^{m+1}(j_x^{m+1}(f)) = j_x^m(f)$  respectively.

In <sup>/27/</sup> the application of analytical computations is illustrated on the Navier-Stokes equation and the existence of a partially invariant solution is exhibited. Furthermore, the corresponding system of equations determining that solution is reduced to a system in involution, and it is pointed out that there is a certain freedom in doing so.

When partial solutions are determined via the method of differential connections <sup>/36/</sup> then the original system of differential equations is completed using different differential connections.

## REFERENCES

1. Ovsyanikov L.V. Grupovoy analiz differentsialnykh uravneniy. Nauka, M., 1978 (in Russian).
2. Ibragimov N.Ch. Grupy preobrazovaniy v matematicheskoy fizike (in Russian), Nauka, M., 1983.
3. Blumen G.W., Cole J.D. Similarity Methods for differential Equations. Springer-Verlag, N.Y., 1974.
4. Wahlquist H.D., Estabrook F.B. Phys.Rev.Lett., 1973, 31, p. 1386; J. of Math.Phys., 1975, 16, p.1; ibid, 1976, 17, p. 1293.
5. Witernitz P. Preprint CRMA-1150, Montreal, 1983.
6. Schwarz F. Comp.Phys.Commun., 1982, 27, p. 179.
7. Edelen D.G.B. Comp. & Math. With Appl., 1980, 6, p. 415.
8. Gragert P.K.H. Symbolic Computations in Prolongation Theory. Ph.D. thesis, Twente University of Technology, The Netherlands, 1981.
9. Eliseev V.P., Korniyak V.V., Fedorova R.N. JINR, 11-84-238, Dubna, 1984 (in Russian).
10. Steeb M.-H., Erig W., Strampp W. J.Math.Phys., 1981, 22, p. 970; ibid, 1982, 23, p. 145.
11. Kersten P.H.M. J.Math.Phys., 1983, 24, p. 2374.
12. Pirani E.A.E., Robinson D.C., Shadwick W.F. Local Jet Bundle Formulation of Bäcklund Transformations, In: Math.Phys.Studies 1, Reidel, Boston, 1979.
13. Zharkov A.Yu., Shvachka A.B. JINR, P11-83-914, Dubna, 1983 (in Russian).
14. Ibragimov N.Ch., Shabat A.B. Functional. analys, 1980, 14, No. 4, p. 79 (in Russian).
15. Svinolupov S.I., Sokolov V.V. In: Integrable systems, Bashkirian branch of the USSR Acad. of Sciences, 1982, Ufa, p. 71 (in Russian).
16. Sokolov V.V., Shabat A.B. Bashkirian branch of the USSR Acad. of Sciences. Preprint, Ufa, 1982 (in Russian).
17. Kotelnikov G.A. In: Group-theoretical methods in physics. Proceedings of the International Seminar, Zvenigorod, 1982. Nauka, M., 1983 (in Russian).
18. Malkin I.A., Manko V.I. Dynamical symmetries and coherent states of quantum systems. Nauka, M., 1979 (in Russian).
19. Malkin I.A., Manko V.I. Pysma v ZhETPh, 1965, 2, p. 230 (in Russian).
20. Fushchich V.I. Theor. and Math.Phys., 1971, 7, p.3.
21. Fushchich V.I. In: Theoretically-algebraical investigations in Mathematical physics. Kiev, Inst. of Mathem.. Acad. of Sciences of UkSSR, 1981 pp. 6-28 (in Russian).
22. Fushchich V.I. In: Group-theoretical methods in physics. Proceedings of the Int. Seminar, Zvenigorod, 1982, M., Nauka, 1982, pp. 407-413 (in Russian).

23. Fushchich V.I., Moskalink S.S. Lett.Nuovo Cim., 1981, 31, p. 571.
24. Fushchich V.I., Senov N.I. Doklady Acad. Nauk USSR, 1982, 263, p. 582 (in Russian).
25. Ganja V.G. et al. In: International Conference on Systems and Techniques of Analytical Computing and Their Applications in Theoretical Physics, Dubna, 1982, JINR, D11-83-511, Dubna, 1983, p. 125-131 (in Russian).
26. Cartan E. External differential systems and their geometrical applications. M., 1962, State Univ., (in Russian).  
Finikov S.P. Metod vneshnich form Cartana. M.-L., GITTL, 1948 (in Russian).
27. Arais E.A., Shapeev V.P., Yanenko N.N. DAN SSSR, 1974, 214, No. 4, p. 296 (in Russian).
28. Palamodov V.P. Linear differential operators with constant coefficients. M., 1964 (in Russian).
29. Samborsky S.N. Differential equations, 1980, 16, No. 3, p. 516 (in Russian).
30. Chebotarevsky B.D. Differential geometry of figures many-folds. Kaliningrad, 1979, No.10, p. 148 (in Russian).
31. Vinogradov A.M., Krasylshyk I.S., Lychagyn V.V. Geometry of jet spaces and nonlinear differential equations. Nauka, M., 1977 (in Russian).
32. Zhishyn A.E. DAN SSSR, 1978, 238, No. 2, p. 15 (in Russian).
33. Meleshko S.V., Shapeev V.P., Yanenko N.N. DAN SSSR, 1980, 254, No. 4 (in Russian).
34. Kuranishi M. Lectures on involutive systems of partial differential equations, Publ. Soc.Math., Sao Paulo, 1967.
35. Yanenko N.N. Trudy IV Vsesouznogo s'ezda, Nauka, L., 1964, 2, p. 247 (in Russian).
36. Reznikov I.G., Steletsky V.I., Topunov V.L. Tezisy Vsesouznoy konferentsyy. Novosibirsk, 1981, p. 177 (in Russian).
37. Melnikov V.K. JINR, P2-81-249, P2-81-262, Dubna, 1981, (in Russian).

**Received by Publishing Department  
on July 11, 1984.**

В Объединенном институте ядерных исследований начал выходить сборник "*Краткие сообщения ОИЯИ*". В нем будут помещаться статьи, содержащие оригинальные научные, научно-технические, методические и прикладные результаты, требующие срочной публикации. Будучи частью "*Сообщений ОИЯИ*", статьи, вошедшие в сборник, имеют, как и другие издания ОИЯИ, статус официальных публикаций.

Сборник "*Краткие сообщения ОИЯИ*" будет выходить регулярно.

The Joint Institute for Nuclear Research begins publishing a collection of papers entitled *JINR Rapid Communications* which is a section of the *JINR Communications* and is intended for the accelerated publication of important results on the following subjects:

- Physics of elementary particles and atomic nuclei.
- Theoretical physics.
- Experimental techniques and methods.
- Accelerators.
- Cryogenics.
- Computing mathematics and methods.
- Solid state physics. Liquids.
- Theory of condensed matter.
- Applied researches.

Being a part of the *JINR Communications*, the articles of new collection like all other publications of the Joint Institute for Nuclear Research have the status of official publications.

*JINR Rapid Communications* will be issued regularly.



SI-64-495

... ..  
... ..  
... ..  
... ..  
... ..

... ..  
... ..

*[Handwritten signature]*

... ..

SI-64-495

... ..  
... ..  
... ..  
... ..  
... ..

... ..

