

REFERENCE

IC/86/95

**INTERNATIONAL CENTRE FOR
THEORETICAL PHYSICS**



ESCAPE RATE FROM STRANGE SETS AS AN EIGENVALUE

T. Tél




**INTERNATIONAL
ATOMIC ENERGY
AGENCY**



**UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION**

1986 MIRAMARE-TRIESTE



International Atomic Energy Agency
and
United Nations Educational Scientific and Cultural Organization

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

ESCAPE RATE FROM STRANGE SETS AS AN EIGENVALUE *

T. Tél **

International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

A new method is applied for calculating the escape rate from chaotic repellers or semi-attractors, based on the eigenvalue problem of the master equation of discrete dynamical systems. The corresponding eigenfunction is found to be smooth along unstable directions and to be, in general, a fractal measure. Examples of one and two dimensional maps are investigated.

MIRAMARE - TRIESTE

June 1986

* To be submitted for publication.

** On leave of absence from the Institute for Theoretical Physics, Eötvös University, Budapest, Hungary.

Chaotic behaviour which can be observed on a long but finite time-scale only has attracted recent interest from both theoretical [1]-[15] and experimental [16]-[19] points of view. This transient chaos is related to strange invariant objects (Cantor sets) of the phase space, called chaotic repellers [8] or, if they are partially attracting, semi-attractors [12] or almost attractors [13]. On the other hand, Cantor sets (being e.g. supports of the density of state) have recently been found to play an essential role in certain localization problems [20] in the dynamics on fractal lattices [21], in models of amorphous solids [22] and of quasicrystals [23]. Since these Cantor sets may be interpreted as chaotic repellers or semi-attractors, the investigation of the latter may be of relevance also for the theory of disordered systems.

An important characteristic of the dynamics in the vicinity of such strange sets is the escape rate [7], [8], α , that measures how fast the repulsion occurs. For discrete dynamical systems, which we shall investigate here, the definition is the following. Let us distribute a large number of points in some neighbourhood Γ of the repeller (semi-attractor). The probability that a randomly chosen point has not yet escaped Γ after n steps is denoted by W_n . As n gets large ($n \rightarrow \infty$) one observes, in general, an exponential decay [6]-[8], [11]-[13]:

$$W_n \sim \exp(-\alpha n), \quad (1)$$

where $\alpha > 0$ is the escape rate (known to be independent of Γ). The long time behaviour, however, should then be governed by the largest eigenvalue $1/q = \exp(-\alpha)$ of the master equation describing the evolution of a probability distribution on Γ , which has now no non-trivial stable solution (see also [11]). This eigenvalue develops from the unit eigenvalue associated with the stationary distribution of a chaotic attractor when the latter ceases to exist and is replaced by a repeller or semi-attractor as the control parameter is raised. Our aim is here to illustrate that the concept of the escape rate as an eigenvalue may provide a rather rapid and simple method for determining α .

First, we consider chaotic transients in one-dimensional maps $x' = f(x)$, where $f(x)$ is a single humped function. Γ is then an interval containing the repeller. Let us investigate the iteration scheme

$$C_{n+1}(x') = q \left(\sum_{x \in f^{-1}(x')} \frac{C_n(x)}{|f'(x)|} \right) \quad (2)$$

starting with a smooth positive function $C_0(x)$ on Γ , where f' denotes the derivative of f , and $0 < q < 1$. According to our numerical studies there is, in a broad class of maps, a single value of q , $q = \exp(-\alpha)$, for which the functions $C_n(x)$ converge towards a finite $C(x)$ for $n \rightarrow \infty$. This $C(x)$ is then independent of the initial condition $C_0(x)$. (For a rigorous proof see [3].) Since $C(x)$ fulfils the equation

$$C(x') = q \left(\sum_{x \in f^{-1}(x')} \frac{C(x)}{|f'(x)|} \right), \quad (3)$$

it can be considered as an eigenfunction of the Frobenius-Perron equation [24] associated with the eigenvalue $1/q = \exp(-\alpha)$. The quantity $\int C(\bar{x}) d\bar{x}$ is called the conditionally invariant measure which has been invented and interpreted by Pianigiani and Yorke (first item of Ref.[3]).

It is worth mentioning here that the fractal dimension of the repeller also appears as an eigenvalue of a certain linear equation. The corresponding eigenfunction (see Eq.(4) of [14]), which was shown there to yield the stationary distribution on the repeller, accessible in numerical simulations or in experiments, is to be distinguished from $C(x)$, the density of the conditionally invariant measure. In what follows we show that Eq.(2), just like Eq.(4) of Ref.[14], may converge rather rapidly, and this fact makes it useful for practical calculations.

In order to illustrate the rapid convergence of the procedure with a simple example we consider the map

$$f(x) = \begin{cases} 1 - a_1 x & , \quad x > 0 \\ 1 + a_2 x & , \quad x < 0 \end{cases}, \quad (4)$$

where $a_1, a_2 > 1$ and $a_1^{-1} + a_2^{-1} < 1$ so that a chaotic repeller shows up. The evolution of a linear initial function $C_0(x) = \gamma_0 x + \beta_0$ on an interval Γ , say $(-1,1)$, can then be followed exactly. The result is $C_n(x) = \gamma_n x + \beta_n$ with

$$\gamma_{n+1} = q (a_2^{-2} - a_1^{-2}) \gamma_n, \quad (5a)$$

$$\beta_{n+1} = q (a_1^{-1} + a_2^{-1}) \beta_n - \gamma_{n+1}. \quad (5b)$$

Since $a_1, a_2 > 1$, γ_n tends towards zero. A non-trivial limit for $n \rightarrow \infty$ exists only if β_n remains constant. This specifies the eigenvalue to be

$$1/q = \exp(-\alpha) = a_1^{-1} + a_2^{-1}. \quad (6)$$

It follows from Eq.(5a) that the convergence to the constant limit solution is then exponentially fast (with a critical slowing down for $a_1(2) \rightarrow \infty$, $a_2(1) \rightarrow 1$).

Our next example is the parabola map defined by $f(x) = 1 - ax^2$ in the region $a > 2$. In order to determine the escape rate we considered the iterates of $C_0(x) \equiv 1$. An explicit expression for $C_n(x)$ follows then from (2) in the form

$$C_n(x') = q^n \sum_{x \in f^{-n}(x')} |(f^n(x))'|^{-1}, \quad (7)$$

where f^n and f^{-n} represents the n^{th} iterate of f and its inverse, respectively. We treated $q > 1$ as a free parameter to be adjusted so that a non-trivial limit solution exists. As long as q is too small (large) $C_n(x)$ monotonously decreases (increases) with n at a fixed x_0 ($= 0.5$). If, however, q is appropriately chosen a convergence is found with an accuracy of less than one percent at the fourth iterate. We used this property to obtain a lower (upper) bound for the escape rate as the value $\alpha = \log q$, where $C_5(x_0) - C_6(x_0) < \epsilon$ ($C_6(x_0) - C_5(x_0) > \epsilon$) with a small positive ϵ was first realized when increasing q . By this way a fast algorithm has been found. Fig.1 displays the plot α versus a in the interval $2 < a < 3$ calculated at 300 different values of a ($\epsilon = 10^{-3}$). Error bars typically correspond to the thickness of the curve $\alpha(a)$.

To find a comparison with the results obtained by means of other methods we applied the present procedure at those special values of a (also for $a > 3$), where the escape rate has been determined by Widom et al. [7]. Table I reflects a quantitative agreement. By a more detailed analysis of the convergence of $C_n(x)$ and by going beyond $n = 6$, q could

have been calculated more accurately but this refinement is not the subject of the present paper.

Due to the rapid convergence of the iteration the eigenfunction $C(x)$ can be safely approximated by $C_0(x)$ of Eq.(7). Fig.2 shows the eigenfunction obtained at different values of a . (It is to be noted that the shortest interval covering the repeller is $(-x^*, x^*)$ with $x^* = (1 + (1 + 4a)^{1/2})/(2a) < 1$.) All $C(x)$ are found to be smooth in $(-1, 1)$ since for $a > 2$ the singularity appearing at $x = 1$ is mapped outside this region. The figure makes it evident that the functions $C(x)$ tend to a finite solution for $a \rightarrow \infty$.

This asymptotic form can be easily deduced from Eq.(3). Since $x = \pm ((1 - x')/a)^{1/2}$ and $C(x)$ is smooth, the argument of $C(x)$ on the right-hand side can be set zero for $a \rightarrow \infty$ and one obtains

$$C(x') = \frac{q C(0)}{a^{1/2} |1 - x'|^{1/2}} \quad (8)$$

The requirement of a finite solution then yields

$$\exp(\alpha) = q = a^{1/2} \quad (9)$$

The same result was obtained by means of a different method in [7]. Curve 4 of Fig.2 corresponding to the eigenfunction at $a = 90$ is, in fact, hard to distinguish from that corresponding to Eqs.(8) and (9).

For maps defined by $f(x) = 1 - a|x|^z$, $z > 0$ we obtain in a similar way

$$\exp(\alpha) = q = (z/2)a^{1/z} \quad (10)$$

$$C(x) = C(0) |1 - x|^{1/z-1} \quad (11)$$

as the asymptotic results for $a \rightarrow \infty$.

Finally, we investigate invertible maps of the plane $x' = T(x)$ producing chaotic transients. Now, the eigenvalue problem can be most conveniently formulated in terms of a certain measure, we call c -measure (the analogue of $\int C_n(\bar{x}) d\bar{x}$ of the one-dimensional case). As an extension of Eq.(2), its evolution is defined by

$$(\mu_j^c)' = q T(\mu_j^c) \quad (12)$$

where μ_j^c denotes the c -measure of a region B around a point x_j lying in a neighbourhood Γ of the semi-attractor, and $T(\mu_j^c)$ represents the action of the dynamics on μ_j^c . The transformed c -measure $(\mu_j^c)'$ belongs to the region $T(B)$ around x_j' . Starting e.g. with the Lebesgue measure on Γ , subsequent applications of (12) with $q = \exp(\alpha)$ lead to the conditionally invariant measure [3]. Note that the conditionally invariant measure again differs from the "natural" invariant measure defined in [14].

This procedure will be illustrated on a generalized version of the baker transformation [25] introduced in [14]. The dynamics is given by

$$y' = \begin{cases} sy & , y < c, \\ t(1-y) & , y > c, \end{cases} \quad x' = \begin{cases} ax & , y < c, \\ 1/2 + bx & , y > c, \end{cases} \quad (13)$$

where $0 < a, b, c < 1/2$ and $sc, t(1-c) > 1$. The latter condition ensures the escape along the unstable direction, y . Although (13) is a piecewise linear model it seems to reflect the most typical features of chaotic semi-attractors. By means of this example, general properties of conditionally invariant measures can be studied which have not been discussed in the literature.

Starting with the Lebesgue measure on the unit square (Γ) Eq.(12) leads to a c -measure q/s and q/t on the strips, $0 < x < a$, $0 < y < 1$ and $1/2 < x < 1/2 + b$, $0 < y < 1$, respectively. After n steps, the c -measure on strips of width $a^m b^{n-m}$, $m = 0, 1, \dots, n$ is $(q/s)^m (q/t)^{n-m}$. The total c -measure of the unit square is then $q^n (s^{-1} + t^{-1})^n$. A non-trivial limit exists only if this quantity remains finite for $n \rightarrow \infty$, from which

$$\exp(\alpha) = q = (s^{-1} + t^{-1})^{-1} \quad (14)$$

follows for the escape rate.

The measure obtained for $n \rightarrow \infty$ has then a smooth density along the y direction, it is, however, a fractal measure [26]. By means of the method used in [25] one obtains for the fractal dimension d_0 [27] and the information dimension d_1 [26] of this measure the equations

$$a^{d_0-1} + b^{d_0-1} = 1, \quad (15)$$

and

$$d_1 = 1 + \frac{t \log(q/s) + s \log(q/t)}{t \log(a) + s \log(b)} \quad (16)$$

respectively. The difference between conditionally and natural invariant measures is reflected also in the difference of their dimensions (c.f. [14]).

Finally, we mention that the smoothness along the unstable direction, the fractal structure along the stable direction, and the fractal measure property seem to be general features of conditionally invariant measures of semi-attractors.

ACKNOWLEDGMENTS

Illuminating discussions with Profs P. Grassberger and P. Szépfalusy are highly acknowledged. Thanks are due to Dr. L. Mendoza for a critical reading of the manuscript. The author would also like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste.

REFERENCES

- [1] T. Shimizu and N. Morioka, *Phys. Lett.* 69A, 148 (1978);
J.L. Kaplan and J.A. Yorke, *Comm. Math. Phys.* 67, 93 (1979);
J.A. Yorke and E.D. Yorke, *J. Stat. Phys.* 21, 263 (1979).
- [2] R. Bowen, *Publ. Math. I.H.E.S.* 50, 11 (1979).
- [3] G. Pianigiani and J.A. Yorke, *Trans. Am. Math. Soc.* 252, 351 (1979);
G. Pianigiani, *J. Math. Anal. Appl.* 82, 75 (1981).
- [4] J. Coste, *J. Stat. Phys.* 23, 521 (1980);
F. Manneville, *Phys. Lett.* 90A, 327 (1982).
- [5] D. Ruelle, *Ergod. Th. Dyn. Syst.* 2, 99 (1982);
H. McCluskey and A. Manning, *Ergod. Th. Dyn. Syst.* 3, 251 (1983).
- [6] C. Grebogi, E. Ott and J.A. Yorke, *Physica* 7D, 181 (1983);
C. Grebogi, E. Ott and J.A. Yorke, *Phys. Rev. Lett.* 50, 935 (1983); and
Ergod. Th. Dyn. Syst. 5, 341 (1986).
- [7] M. Widom, D. Bensimon, L.P. Kadanoff and S.J. Shenker, *J. Stat. Phys.* 32, 443 (1983).
- [8] L.P. Skadanoff and C. Tang, *Proc. Nat. Acad. Sci.* 81, 1276 (1984).
- [9] S. Takesue and K. Kaneko, *Prog. Theor. Phys.* 71, 35 (1984);
S.W. McDonald, C. Grebogi, E. Ott and J.A. Yorke, *Physica* 17D, 125 (1985).
- [10] F.T. Arecchi, R. Badii and A. Politi, *Phys. Lett.* 103A, 3 (1984).
- [11] M. Widom and S.J. Shenker in *Chaos and Statistical Methods*, Ed. Y. Kuramoto (Springer-Verlag, New York, 1984);
L.P. Kadanoff in *Regular and Chaotic Motions in Dynamic Systems*, Eds. G. Velo and A.S. Wightman (Plenum Press, New York, 1985).
- [12] H. Kantz and P. Grassberger, *Physica* 17D, 75 (1985).
- [13] J.P. Eckmann and D. Ruelle, *Rev. Mod. Phys.* 57, 617 (1985).
- [14] P. Szépfalusy and T. Tél, to appear in *Phys. Rev. A*.
- [15] S. Takesue, unpublished.
- [16] G. Ahlers and R.W. Walden, *Phys. Rev. Lett.* 44, 445 (1980).

- [17] F.T. Arecchi and F. Lisi, Phys. Rev. Lett. 49, 94 (1982);
 F.T. Arecchi, R. Meucci, G.P. Puccioni and J.R. Tredicce, Phys. Rev. Lett. 49, 1217 (1982);
 F.T. Arecchi and A. Califano, Phys. Lett. 101A, 443 (1984).
- [18] P. Bergé and M. Dubois, Phys. Lett. 93A, 365 (1983).
- [19] M. Gorman, P.J. Widman and K.A. Robbins, Phys. Rev. Lett. 52, 2241 (1984); and Physica 19D, 255 (1986).
- [20] M. Kohmoto, L.P. Kadanoff and C. Tang, Phys. Rev. Lett. 50, 1870 (1983).
- [21] E. Domany, S. Alexander, D. Bensimon and L.P. Kadanoff, Phys. Rev. B28, 3110 (1983);
 C.K. Harris and R.B. Stinchcombe, Phys. Rev. Lett. 50, 1399 (1983);
 R. Rammal, J. Phys. (Paris) 45, 191 (1984);
 J.R. Banavar, L.P. Kadanoff and A.M. Pruisken, Phys. Rev. E31, 1388 (1985);
 R.B. Stinchcombe in Scaling Phenomena in Disordered Systems, Eds. R. Fynn and A. Skjeltorp (Plenum Press, New York, 1985).
- [22] R. Schilling, Phys. Rev. Lett. 53, 2258 (1984);
 P. Reichert and R. Schilling, Phys. Rev. B32, 5731 (1985).
- [23] J.M. Luck and D. Petritis, J. Stat. Phys. 42, 121 (1986);
 J.P. Lu, T. Odagaki and J.L. Birman, Phys. Rev. B33, 4809 (1986);
 M.C. Valsakumar and G. Ananthakrishna, unpublished.
- [24] P. Collet and J.P. Eckmann, Iterated Maps on the Interval as Dynamical Systems (Birkhäuser, Cambridge MA, 1980).
- [25] D. Farmer, E. Ott and J.A. Yorke, Physica 7D, 153 (1983).
- [26] D. Farmer, Z. Naturforschung 37a, 1304 (1982).
- [27] B.B. Mandelbrot, The Fractal Geometry of Nature (Freeman, San Francisco, 1982);
 K.J. Falconer, The Geometry of Fractal Sets (Cambridge University Press, Cambridge, 1985).

TABLE 1

a	q	q-th	q-exp
2.4725	1.3019 ± 2.10^{-4}	1.3020	1.3022
3.75	1.7381 ± 2.10^{-4}	1.7384	1.7380
20.0	4.3888 ± 3.10^{-4}	4.3888	4.3887
48.75	6.9286 ± 5.10^{-4}	6.9278	6.9285
90.0	9.4473 ± 7.10^{-4}	9.4458	9.4473

A comparison with the results of [7]. The second column contains the reciprocal eigenvalue q obtained with the procedure described in the text. At these values of a the convergence of the series $C_n(x_0)$, $n > 3$ seems to be monotonous, therefore, an ϵ as small as 10^{-4} could have been chosen. The data of the last two columns are taken over from Ref.[7] and show the results of a theoretical and an 'experimental' method.




Fig.1 Escape rate $\alpha = \log(q)$ obtained as an eigenvalue of Eq.(2) for the map $x' = 1 - ax^2$, $2 < a < 3$. Near $a = 2$ a power law behaviour with exponent $1/2$ is present (see e.g. Ref.[15]).




Fig.2 Eigenfunction $C(x)$ (not normalized). The curves 1, 2, 3 and 4 belong to control parameter value $a = 2.01, 2.10, 3.75$ and 90.0 , respectively.