



Ontario Hydro

A REVIEW OF THE INORGANIC AND  
ORGANOMETALLIC CHEMISTRY OF ZIRCONIUM

OH-- Report No 85-124-K

A. Kuksis Kalvins  
Chemist  
Atmospheric Research Section  
Chemical Research Department



RESEARCH



**ontario hydro  
research division**

A REVIEW OF THE INORGANIC AND  
ORGANOMETALLIC CHEMISTRY OF ZIRCONIUM

OH-- Report No 85-124-K

A. Kuksis Kalvins  
Chemist  
Atmospheric Research Section  
Chemical Research Department

ABSTRACT

The results of a literature review of the inorganic and organometallic chemistry of zirconium are presented. Compounds with physical and chemical properties compatible with the requirements of an ir laser zirconium isotope separation process have been identified.

job	740635-575-051	file	836.46	date	April 12, 1985	report no.	85-124-K
-----	----------------	------	--------	------	----------------	------------	----------



ontario hydro  
research division

EXECUTIVE SUMMARY

A REVIEW OF THE INORGANIC AND  
ORGANOMETALLIC CHEMISTRY OF ZIRCONIUM

A. Kuksis Kalvins  
Atmospheric Research Section  
Chemical Research Department

This report presents the results of a literature review of the inorganic and organometallic chemistry of zirconium. The goal of the review was to identify candidate compounds for an infrared (ir) laser isotope separation process. In order for a compound to be suitable for ir laser based isotope separation it must have a stretching or bending vibration in its ir spectrum, ie, 9-16  $\mu\text{m}$  (1111 - 625  $\text{cm}^{-1}$ ). The vibration must have an isotopic shift comparable with and preferably greater than its absorption width. The compound must also be relatively volatile at moderate temperatures, thermally robust and possess a relatively low activation energy to unimolecular decomposition or intramolecular reaction. A number of candidate compounds were identified, these include,

- anhydrous  $\text{ZrOCl}_2$  ( $\nu\text{Zr-O}$  877  $\text{cm}^{-1}$ , 11.5  $\mu\text{m}$ )
- $[\text{Zr}(\text{NR}_2)_4]$  ( $\nu\text{Zr-N}$  533-677  $\text{cm}^{-1}$ , 19-15  $\mu\text{m}$ )
- $[\text{Zr}(\text{SOCNR}_2)_4]$  ( $\nu\text{Zr-O}$  552-569  $\text{cm}^{-1}$ , 18-17.6  $\mu\text{m}$ )
- $[\text{Cp}_2\text{Zr}(\text{OR})_2]$  ( $\nu\text{Zr-O}$  1090-1140  $\text{cm}^{-1}$ , 9.2-8.8  $\mu\text{m}$ )
- $[\text{Cp}_x\text{Zr}(\text{NR}_2)_{4-x}]$  ( $\nu\text{Zr-N}$  500-640  $\text{cm}^{-1}$ , 20-15  $\mu\text{m}$ ).

Compounds such as  $[\text{Zr}(\text{hfacac})_4]$ ,  $[\text{Zr}(\text{tfacac})_4]$ ,  $[\text{Cp}_2\text{Zr}(\text{acac})_2\text{Cl}]$  and  $[\text{CpZr}(\text{hfacac})_3]$  (acac = acetylacetonate, hfacac = hexa-fluoroacetylacetonate, tfacac = trifluoroacetylacetonate) may also be suitable however the data required to completely assess their potential, eg, ir spectra, are not available. The compounds  $[\text{ZrOCl}_2]$ ,  $[\text{Zr}(\text{NET}_2)_4]$  and  $[\text{Zr}(\text{tfacac})_4]$  are commercially available.

Synthesis of many of these complexes involves a number of steps and yields are often low. All of the compounds should however be tested in order to assess the feasibility of ir laser based zirconium isotope separation. If the process is viable, methods to improve the yields can be examined.

job	file	date	report no.
740635-575-051	836.46	April 12, 1985	85-124-K

## TABLE OF CONTENTS

	<u>Page</u>
GLOSSARY OF ABBREVIATIONS	ii
1.0 INTRODUCTION	1
2.0 REQUIREMENTS	2
3.0 CHEMICAL AND PHYSICAL PROPERTIES OF ZIRCONIUM	3
3.1 Inorganic Chemistry	5
3.2 Organometallic Chemistry	14
4.0 CONCLUSIONS AND RECOMMENDATIONS	21
REFERENCES	24
DISTRIBUTION	last page

## GLOSSARY OF ABBREVIATIONS

acac	acetylacetonate ion, 2,4-pentanedionate
as	asymmetric
nBu	n-butyl
cm <sup>-1</sup>	wavenumber
C <sub>p</sub>	$\pi$ -cyclopentadienyl, $\eta^5$ -C <sub>5</sub> H <sub>5</sub>
d <sup>n</sup>	formal d-electron configuration
Et	ethyl
hfacac	hexafluoroacetylacetonate ion, 1,1,1,5,5,5-hexafluoro-2,4-pentanedionate
ir	infrared
L	any unidentate ligand
Me	methyl, CH <sub>3</sub>
Ph	phenyl
nPr	n-propyl
R	an alkyl group
RT	room temperature
s, sym	symmetric
tfacac	trifluoroacetylacetonate ion, 1,1,1-trifluoro-2,4-pentanedionate
THF	tetrahydrofuran
X	a halogen
vs	very strong



# ontario hydro research division

To Mr. F.J. Kee  
Director of Research

## A REVIEW OF THE INORGANIC AND ORGANOMETALLIC CHEMISTRY OF ZIRCONIUM

### 1.0 INTRODUCTION

The recent zirconium pressure tube cracking problems in Ontario Hydro's CANDU reactors have renewed interest in the development of a technique for removal of the zirconium-91 isotope from the zirconium isotope mix. Zirconium has five stable isotopes, see Table 1, the majority of which have a very low neutron absorption character. The zirconium-91 isotope however has a thermal neutron capture cross-section of 1.58 barns/1/. Selective removal of this isotope would permit construction and use of more conservative, ie, thicker zirconium pressure tubes without concomitant loss of fuel burn-up due to increased neutron absorption by the metal. Thicker pressure tubes would offer a number of advantages, these include

- increased resistance to hydride cracking,
- creep and elongation problems would be diminished,
- higher fuel channel temperature and pressure operating conditions could be accommodated resulting in a more efficient conversion of heat to power, ie, more power per kilogram of uranium/2/.

A possible route to zirconium isotope separation is one based on a uranium isotope separation technique using infrared lasers reported by Kaldor et al in 1979/3,4/. This method relies on increasing, in an isotopically selective fashion, the internal energy of a target molecule to the point where some thermally allowed intra- or intermolecular reaction, ie bond cleavage, proceeds at a suitable rate. In Kaldor's experiment,  $[\text{UO}_2(\text{hfacac})_2 \cdot \text{THF}]$  was irradiated with a  $\text{CO}_2$  laser. The absorption band due to the asymmetric stretch of the moiety  $\text{UO}_2^{2+}$ ,  $\nu(\text{O-U-O})$ , falls within the range of the  $\text{CO}_2$  laser. Irradiation at this frequency resulted in an isotope specific unimolecular dissociation of the molecule. The U-O bond associated with the THF ligand was broken, ie, the weakest bond in the molecule, resulting in loss of THF.

TABLE 1  
STABLE ISOTOPES OF ZIRCONIUM/1/

Isotope	% Natural Abundance	Atomic Mass	Neutron-Capture Cross-Section (barns)
${}_{40}\text{Zr}^{90}$	51.4	89.9043	$0.10 \pm 0.076$
${}_{40}\text{Zr}^{91}$	11.2	90.9053	$1.58 \pm 0.12$
${}_{40}\text{Zr}^{92}$	17.1	91.9046	$0.25 \pm 0.12$
${}_{40}\text{Zr}^{94}$	17.5	93.9061	$0.075 \pm 0.008$
${}_{40}\text{Zr}^{96}$	2.8	95.9082	$0.05 \pm 0.01$

Application of infrared (ir) laser based isotope separation to removal of the zirconium-91 isotope requires an inorganic or organozirconium complex which contains a zirconium-91-ligand vibration within the operating range of an infrared laser. The purpose of this study was to examine the chemistry of zirconium in order to determine if there are any zirconium complexes, either commercially available or readily synthesized, which possess properties suitable for laser based zirconium isotope separation. In the following, the physical and chemical properties that a compound must possess to be suitable for laser based isotope separation, and the inorganic and organometallic chemistry of zirconium as related to these requirements are examined.

## 2.0 REQUIREMENTS


To be suitable for infrared laser based isotope separation, a complex must have a stretching or bending vibration in its ir spectrum at a frequency within the operating range of an infrared laser, ie, 9-16  $\mu\text{m}$  (1111-625  $\text{cm}^{-1}$ ). The band must be relatively sharp, ie,  $<10 \text{ cm}^{-1}$  fwhm, and in a region free of other absorptions/6/. The vibration must also have an isotopic shift comparable with and preferably greater than its absorption width.

The compound must be relatively volatile at moderate temperatures (~1 torr at 100°C), thermally robust and possess a relatively low activation energy to unimolecular decomposition or intramolecular reaction/3/. The molecule should be polyatomic in order that there is an abundance of vibrational-rotational transitions which may facilitate ready absorption of many photons up to the dissociation limit/5/. There is no such multiple absorption of ir photons in simple (two or three atom) molecules where the number of possible vibrational-rotational transitions is limited.

### 3.0 CHEMICAL AND PHYSICAL PROPERTIES OF ZIRCONIUM

Zirconium is a hard, corrosion resistant, high melting metal (mp 1855°C ± 15°C), resembling stainless steel in appearance/7/. An oxide film on the surface is believed to account for its lack of reactivity at room temperature. At higher temperatures zirconium is slowly attacked by aqua regia (HCl/HNO<sub>3</sub>), phosphoric acid, hydrofluoric acid, sulphuric acid and molten sodium bisulphate, sodium carbonate and sodium peroxide. At higher temperatures it also reacts readily with hydrogen (reversibly), oxygen, nitrogen and the halogens.

Zirconium and hafnium occur together in nature and their chemistries are very similar. Unless specifically purified, zirconium usually contains from 0.5-2% hafnium/8/. Pure zirconium is required for nuclear applications and is prepared commercially via the Kroll process. The Kroll process involves reaction of the vapour of ZrCl<sub>4</sub> with molten magnesium under an atmosphere of pure helium or argon, Equation 1.



The reaction is exothermic,  $\Delta H^0_{1150 \text{ K}} = -78.6 \text{ kcal/8/}$ . Distillation at 925 K in vacuo removes the magnesium chloride and excess magnesium leaving a highly reactive zirconium sponge which is cooled under helium and compacted by arc melting under argon. Extremely pure zirconium is obtained by zone refining.

The most commonly observed oxidation states and stereochemistries of zirconium are summarized in Table 2. Zirconium in its elemental form has an electronic configuration with four valence electrons outside a noble gas core, ie [Ar]5s<sup>2</sup>4d<sup>2</sup>. The chemistry of zirconium is restricted primarily to the reactions of the metal in the 4+ oxidation state. The zirconium 4+ ion is relatively large, spherical and highly charged. The high size to charge ratio of the Zr<sup>4+</sup> ion renders it a relatively hard Lewis acid and thus promotes strong bond formation to small, (nonpolarizable) anions such as oxide or fluoride. The Zr<sup>4+</sup> ion

TABLE 2

## OXIDATION STATES AND STEREOCHEMISTRY OF ZIRCONIUM/7/

Oxidation State	Coordination Number	Geometry	Examples
Zr(0)	6	Octahedral (?)	[Zr(bipy) <sub>3</sub> ](?)
Zr(I) d <sup>3</sup> Zr(II) d <sup>2</sup>		Complex sheet and cluster structures	
Zr(III) d <sup>1</sup>	6	Octahedral	ZrCl <sub>3</sub> , ZrBr <sub>3</sub> , ZrI <sub>3</sub>
Zr(IV) d <sup>0</sup>	4	Tetrahedral	ZrCl <sub>4</sub> (g), Zr(CH <sub>2</sub> C <sub>6</sub> H <sub>5</sub> ) <sub>4</sub>
	6	Octahedral	Li <sub>2</sub> ZrF <sub>6</sub> , Zr(acac) <sub>2</sub> Cl <sub>2</sub> ZrCl <sub>6</sub> <sup>2-</sup> , ZrCl <sub>4</sub> (S)
	7	Pentagonal bipyramidal Capped trigonal prism	Na <sub>3</sub> ZrF <sub>7</sub> , K <sub>2</sub> CuZr <sub>2</sub> F <sub>12</sub> ·6H <sub>2</sub> O (NH <sub>4</sub> ) <sub>3</sub> ZrF <sub>7</sub>
		Tetragonal base-trigonal base polyhedron	ZrO <sub>2</sub>
	8	Square antiprism Dodecahedron	Zr(acac) <sub>4</sub> , Zr(SO <sub>4</sub> ) <sub>2</sub> ·4H <sub>2</sub> O [Zr(C <sub>2</sub> O <sub>4</sub> ) <sub>4</sub> ] <sup>4-</sup> , [ZrX <sub>6</sub> (diars) <sub>2</sub> ] [Zr <sub>4</sub> (OH) <sub>8</sub> (H <sub>2</sub> O) <sub>16</sub> ] <sup>8+</sup>

has no partly filled outer electron shell to give it stereochemical preferences. These properties result in high coordination numbers, eg 6, 7, 8, and a great variety of coordination polyhedra. Discrete, 4-coordinated molecular species are found only in special cases, generally when there is steric hindrance combined with ligand-donor  $\pi$ -bonding, eg  $Zr(BH_4)_4$ ,  $Zr(allyl)_4$ ,  $Zr(NR_2)_4$ . In general, the most common higher coordination number is eight. This is usually promoted by the most electro-negative donors,  $F^-$ ,  $O^{2-}$ ,  $OH^-$ .  $OH_2$  or by polydentate groups.

The following review of the chemistry of zirconium is divided into two parts, inorganic and organometallic complexes. Organometallic compounds are those in which the carbon atoms of organic groups are bound to metal atoms/7/. The term inorganic compound denotes all other complexes, ie, those containing metal-halogen, metal-oxygen, metal-nitrogen bonds, etc.

### 3.1 Inorganic Chemistry

Inorganic complexes of zirconium include zirconium halides, oxides, chalcogenides, phosphates, etc. Zirconium halides, specifically  $ZrCl_4$ , are common starting materials for the preparation of many other inorganic and organometallic zirconium complexes and will be discussed first.

All four zirconium tetrahalides,  $ZrF_4$ ,  $ZrCl_4$ ,  $ZrBr_4$  and  $ZrI_4$  are known and are commercially available/9/. The compounds are tetrahedral monomers in the gas phase and the solids are polymers with bridging halides. The infrared stretching frequencies of Zr-X generally occur at frequencies too low for an infrared laser, eg.

$$\begin{aligned} \nu_{Zr-Cl} \text{ (terminal, gas)} &= 431-388 \text{ cm}^{-1} \text{ (23-26 } \mu\text{m)}; \\ \nu_{Zr-Cl} \text{ (bridge, solid)} &= 310-291 \text{ cm}^{-1} \text{ (32-34 } \mu\text{m)} /8/. \end{aligned}$$

These compounds are therefore not expected to be suitable candidates for an ir laser zirconium isotope separation process. Derivatives of some of the tetrahalides may however be useful.

Zirconium tetrachloride is a white, crystalline polymeric solid subliming at  $331^\circ C$ /7/. The polymer consists of zig-zag chains of  $ZrCl_6$  octahedra.  $ZrCl_4$  fumes in moist air and is vigorously hydrolyzed, Equation 2/7/.



The product of hydrolysis,  $ZrOCl_2 \cdot 8H_2O$ , is a stable oxide chloride containing the polymeric cation  $[Zr_4(OH)_8(H_2O)_{16}]^{8+}$

together with chloride ions. Anhydrous  $ZrOCl_2$  is made by reaction of chlorine monoxide and  $ZrCl_4$ . Its infrared spectrum is interpreted on the basis of an ionic aggregate of  $[Zr=O]^{2+}$  and polymeric  $[ZrOCl_4]_n^{2-}$ . The ir spectrum shows a broad band at  $877\text{ cm}^{-1}$  ( $11.5\ \mu\text{m}$ ) which is assigned to  $\nu(Zr=O)$ . There is however considerable doubt about the existence of a species such as  $[Zr=O]^{2+}/8/$ .

The chemistries of zirconium tetrabromide and zirconium tetraiodide are similar to zirconium tetrachloride. Zirconium tetrafluoride is a white solid which sublimes at  $903^\circ\text{C}/7/$ . It has an eight coordinate structure consisting of square antiprisms which are joined by sharing fluorines. The infrared spectrum of the vapour at  $800^\circ\text{C}$  shows bands at  $668\text{ cm}^{-1}$  ( $15\ \mu\text{m}$ ) and  $190\text{ cm}^{-1}$  ( $52.6\ \mu\text{m}$ ) due to Zr-F stretching and bending modes, respectively. The ir spectrum of the solid shows a Zr-F stretching band at  $550\text{ cm}^{-1}$  ( $18.2\ \mu\text{m}$ ) and ZrF bending vibrations at 320, 295 and  $255\text{ cm}^{-1}$  ( $31.3$ ,  $33.9$  and  $39.2\ \mu\text{m}$ , respectively)/8/.  $ZrF_4$  is the only tetrahalide which combines with  $H_2O$  to form stable hydrates,  $ZrF_4 \cdot xH_2O$ ,  $x = 1$  or  $3$ . The trihydrate has an eight coordinate structure with two bridging fluorides,  $(H_2O)_3F_3ZrF_2ZrF_3(H_2O)_3$  /7/.

The best known zirconium trihalides are  $ZrCl_3$ ,  $ZrBr_3$ , and  $ZrI_3$ . The structures of these compounds consist of close-packed halide layers between which equally spaced zirconium atoms occupy one-third of the octahedral interstices to form infinite linear chains of metal atoms perpendicular to the layers, ie, infinite parallel chains of  $MX_6$  octahedra sharing opposite faces/7/.

There is some evidence suggesting the existence of  $ZrX_x$ ,  $1 < x < 3$ . The structure is believed to consist of stacked, hexagonally packed layers of either all metal atoms or all X atoms, with a stacking sequence XMMX...XMMX. The X...X interlayer distances are normal van der Waals contacts and the M-X distances are appropriate for single bonds/7/. The monohalides have great thermal stability ( $mp > 1100^\circ\text{C}$ ), metallic reflectivity, cleave like graphite and are generally metallic in character.  $ZrCl$  and  $ZrBr$  react with  $H_2$  to give discrete though nonstoichiometric phases, eg  $ZrXH_{0.5}$ ,  $ZrXH$ ,  $X = Cl, Br$ . The hydrogen atoms are inserted through and between double metal layers of the initial halides/7/.

The main oxide formed by zirconium is  $ZrO_2$ . The material has an extremely high melting point,  $2700^\circ\text{C}$ , good mechanical properties and is exceptionally resistant to attack by both acids and alkalis/7/. The compound has a number of structures including the monoclinic form shown in Figure 1/8/. At very high temperatures a number of 'zirconates' may be prepared by reaction of  $ZrO_2$  with metal oxides, metal hydroxides, etc.

A number of zirconium chalcogenides,  $ZrY_x$ ,  $x = 3, 2$ ,  $Y = S, Se, Te$ , also exist. Zirconium exhibits a lower coordination number in  $ZrS_2$  than in  $ZrO_2$ . The disulphide is stable in air or water, but is readily attacked by oxidizing reagents. The lower zirconium selenides and tellurides are similar to the corresponding sulphides. All compounds have layered structures and are intrinsic semiconductors/7/.





Figure 1. Monoclinic  $ZrO_2$  (baddeleyite). Small circle = Zr atom; large circles = oxygen atoms./8/

Zirconium nitrides ( $ZrN$ ), phosphides ( $ZrP_2$ ,  $ZrP$ ,  $Zr_3P$ ), arsenides ( $ZrAs_2$ ), carbides ( $ZrC$ ), silicides ( $ZrSi_2$ ,  $Zr_5Si_4$ ,  $Zr_2Si$ ), germanides and borides are also known. Most of these compounds have very high melting points, eg  $ZrN$ , mp  $2980^\circ C$ ;  $ZrC$ , mp  $3500^\circ C$ ; and complicated structures/8/.

$ZrCl_4$  is a common starting material for the preparation of a number of other inorganic zirconium complexes. Reaction of  $ZrCl_4$  with carboxylic acid derivatives yields zirconium carboxylates, eg



As the alkyl chainlength is increased, solubility in water decreases and solubility in hydrocarbons increases. Some solution ir studies of zirconium carboxylates have been reported/10/. Bands in the  $650\text{ cm}^{-1}$  ( $15.4\ \mu\text{m}$ ) and  $470\text{ cm}^{-1}$  ( $21.3\ \mu\text{m}$ ) region were generally assigned to  $\nu Zr-O$ . This region

of the ir spectra was not however well characterized. Thermal stability of the tetracarboxylates is not great and decomposition to oxycompounds occurs readily/8/. This thermal instability may preclude their utility in ir laser based Zr isotope separation. Recall from Section 2.0, one of the requirements was that the compound be thermally robust.

Tetrakis diketonates such as  $Zr(acac)_4$ ,  $acac = acetylacetonate$ , are also known.  $Zr(acac)_4$  is commercially available/9/. The crystal structure of  $Zr(acac)_4$  reveals that it is eight-coordinate and the diketonate rings are slightly distorted/8/. The ir spectrum of these compounds is well characterized. The zirconium-oxygen stretching frequency is sensitive to the metal coordination number in the series  $Zr(acac)_4$ ,  $ZrCl(acac)_3$ ,  $ZrCl_2(acac)_2$ . The absorption band due to the Zr-O symmetric stretch generally occurs in the region  $449-460\text{ cm}^{-1}$  ( $22.3-21.7\text{ }\mu\text{m}$ ), the Zr-O asymmetric stretch in the region  $421-450\text{ cm}^{-1}$  ( $23.8-22.2\text{ }\mu\text{m}$ ) and  $\nu Zr-X$ , X = halogen, in the region  $<314\text{ cm}^{-1}$  ( $>32\text{ }\mu\text{m}$ ), see Table 3/8/.

Analogous hexafluoroacetylacetonate (hfacac) and trifluoroacetylacetonate (tfacac) zirconium complexes have also been prepared/11/. The fluorine containing derivatives are generally more volatile than the analogous acac compounds and melting points decrease in the series acac, tfacac, hfacac/11/. Infra-red data have been reported for  $Zr(hfacac)_4$  and  $Zr(tfacac)_4$ , however the region below  $1000\text{ cm}^{-1}$ , ie the region where  $\nu(Zr-O)$  would be seen, was not well characterized/11/. In general the absorption bands in the fluorine substituted complexes are shifted to higher wavenumbers relative to those seen for the acac species. Therefore, although the Zr-O stretching vibrations for the acac derivative,  $[Zr(acac)_4]$ , are out of the range of an ir laser,  $\nu Zr-O$  in the analogous fluorine substituted compound may fall within the operating range of an ir laser. Tetrakis(trifluoroacetylacetonate) zirconium is commercially available/9/.

A number of zirconium alkoxy compounds have also been reported. The tetraalkoxides are coordination polymers,  $[M(OR)_4]_n$  ( $n \sim 3.5$  for lower aliphatic R groups) due to bridging by alkoxy groups/8/. Steric hindrance in secondary and tertiary alkoxy groups prevents bridging and leads to volatile, monomeric compounds. The alkoxides are readily hydrolyzed and react with hydroxylic compounds, acids,  $\beta$ -diketonates, etc. The infrared spectra of these compounds have been well characterized, see Table 4. The Zr-O (bridging) stretching vibration generally falls in the region  $400-600\text{ cm}^{-1}$  ( $17-23\text{ }\mu\text{m}$ ). The polymeric nature of the compounds, ie low volatility, decreases their potential utility in an ir laser isotope separation process.

TABLE 3

INFRARED ABSORPTIONS OF SELECTED  
ZIRCONIUM DIKETONATE COMPLEXES/7/


Complex	$\nu(\text{Zr-O})_s$ cm <sup>-1</sup> (μm)	$\nu(\text{Zr-O})_{as}$ cm <sup>-1</sup> (μm)	$\nu(\text{Zr-X})$ cm <sup>-1</sup> (μm)
Zr(acac) <sub>4</sub>		421 (23.8) 301 (33.2)	
ZrCl(acac) <sub>3</sub>	449 (22.3)	432 (23.1)	314 (31.8)
ZrBr(acac) <sub>3</sub>	449 (22.3)	434 (23.0)	164 (61)
ZrI(acac) <sub>3</sub>	452 (22.1)	438 (22.8)	93 (108)
ZrCl <sub>2</sub> (acac) <sub>2</sub>	460 (21.7)	450 (22.2)	335 (30) 325 (31)
ZrBr <sub>2</sub> (acac) <sub>2</sub>	460 (21.7)	450 (22.2)	213 (47)

$$\begin{array}{c} \text{O} \quad \quad \text{O} \\ \parallel \quad \quad \parallel \\ \text{CH}_3\text{C}(\text{O}) \text{CH} \text{C}(\text{O})\text{CH}_3 \end{array}$$

acac = acetylacetonate ligand, CH<sub>3</sub>C(O)CHC(O)CH<sub>3</sub>

s = symmetric stretch  
as = asymmetric stretch

Reaction of ZrCl<sub>4</sub> with N<sub>2</sub>O<sub>5</sub> yields the adduct I, Equation 3/6/,



I

The compound forms colourless sublimable crystals but decomposes at 140°C. The ir spectrum of I suggests that the nitrate groups are bidentate. The far ir and Raman spectra have been interpreted in terms of a tetrachelated, 8-coordinate (dodecahedral) molecule/8/. The Zr-O stretching vibrations were not however reported. A number of basic zirconium nitrates are known. These include [Zr(OH)<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>4</sub>], [ZrO(NO<sub>3</sub>)<sub>2</sub>], [Zr(diket)<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>], and [Zr(NO<sub>3</sub>)<sub>4</sub>]. Zirconium nitrate, in an aqueous solution, is commercially available/9/.

TABLE 4

## PHYSICAL PROPERTIES OF SELECTED ZIRCONIUM ALKOXIDES/8/

Compound	Volatility (°C/mm HG)	Degree of Polymerization	$\Delta H_{\text{vap}}$ (kcal mole <sup>-1</sup> )	IR Spectra ( $\nu_{\text{Zr-O}}$ )	
				cm <sup>-1</sup>	$\mu\text{m}$
$[\text{Zr}(\text{OCH}_3)_4]$	280/0.1	-	-	492, 430	20.3, 23.3
$[\text{Zr}(\text{OC}_2\text{H}_5)_4]^*$	234.8/5.0	3.6	30.2	520, 463, 412	19.2, 21.6, 24.3
$[\text{Zr}(\text{OC}_3\text{H}_7)_4]^*$					
$[\text{Zr}\{\text{OCH}(\text{CH}_3)_2\}_4]^{**}$	203.8/5.0	3.57	31.5	567, 550, 510, 459	17.6, 18.2, 19.6, 21.8
$[\text{Zr}(\text{OC}(\text{CH}_3)_3)_4]$	89.1/5.0	1.0	15.2	557, 540	18.0, 18.5
$[\text{Zr}(\text{OSi}(\text{CH}_3)_3)_4]$	135/0.1	2.05	-	521	19.2
$[\text{Zr}(\text{OC}_4\text{H}_9)_4 \cdot \text{C}_6\text{H}_5\text{OH}]^*$					

\* commercially available/9/

\*\*  $[\text{Zr}(\text{OC}_3\text{H}_7)_4 \cdot \text{C}_3\text{H}_7\text{OH}]$  available from Alfa/9/



TABLE 5

PHYSICAL PROPERTIES OF SELECTED DERIVATIVES OF  $\{Zr(NR_2)_4\}$

R	mp (°C)	Volatility °C/mm Hg	$\nu_{Zr-N}$ cm <sup>-1</sup> (μm)	$\nu_{NC_2}$ sym cm <sup>-1</sup> (μm)
Me (CH <sub>3</sub> )	70	60/0.001	537 s (18.6)	936 vs (10.7)
Et (C <sub>2</sub> H <sub>5</sub> )*	liquid	110/0.01	557 m (17.3)	1000 vs (10)
<sup>n</sup> Pr(C <sub>3</sub> H <sub>7</sub> )	liquid	165/0.1	637 (15.7)	
<sup>n</sup> Bu (C <sub>4</sub> H <sub>9</sub> )	liquid	180/0.1	637 (15.7)	

\* available commercially from Alfa/9/

The thiocarbamate ligands are bidentate and the zirconium complexes are eight coordinate. The compounds are thermally stable, hydrolytically unstable, soluble in  $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ , but insoluble in saturated hydrocarbons. Solubility in water increases with the size of the alkyl group. The compounds are hydrolyzed rapidly upon exposure to air. Their ir spectra show strong bands in the  $1500\text{-}1550\text{ cm}^{-1}$  ( $6.7\text{-}6.5\text{ }\mu\text{m}$ ) region due to the  $\text{C}=\text{O}$  and  $\text{C}=\text{N}$  coupled stretching vibrations and bands in the range  $552\text{-}569\text{ cm}^{-1}$  ( $18.1\text{-}17.6\text{ }\mu\text{m}$ ) due to  $\nu(\text{Zr-O})$ . No strong absorptions which can be attributed to  $\nu(\text{Zr-S})$  were observed in the region  $250\text{-}380\text{ cm}^{-1}$  ( $40\text{-}26.3\text{ }\mu\text{m}$ ). These bands may however be too weak to be observed/13/.

Tetrakis-N,N-dialkyldithiocarbamate zirconium compounds,  $[\text{Zr}(\text{S}_2\text{CNR}_2)_4]$ , have also been reported. Their ir spectra consist of bands in the region  $1488\text{-}1520\text{ cm}^{-1}$  ( $6.7\text{-}6.6\text{ }\mu\text{m}$ ) due to  $\nu(\text{N}=\text{C})$ ,  $994\text{-}998\text{ cm}^{-1}$  ( $\sim 10\text{ }\mu\text{m}$ ) due to  $\nu(\text{C-S})$  and  $\nu(\text{Zr-S})$  is found in the range  $353\text{-}359\text{ cm}^{-1}$  ( $\sim 28\text{ }\mu\text{m}$ ). The dialkyldithiocarbamate ligand behaves as a chelate/8/.

A number of inorganic zirconium complexes have been reviewed in the previous section, however, only a very small fraction of these compounds appear to meet the requirements outlined in Section 2.0. Compounds which have an ir absorption band within or close to the range of an ir laser,  $9\text{-}16\text{ }\mu\text{m}$  ( $1111\text{-}625\text{ cm}^{-1}$ ), include

- anhydrous  $\text{ZrOCl}_2$  ( $\nu\text{Zr-O } 877\text{ cm}^{-1}$ ,  $11.5\text{ }\mu\text{m}$ )
- $[\text{Zr}(\text{NR}_2)_4]$  ( $\nu\text{Zr-N } 533\text{-}677\text{ cm}^{-1}$ ,  $19\text{-}15\text{ }\mu\text{m}$ )
- $[\text{Zr}(\text{SOCNR}_2)_4]$  ( $\nu\text{Zr-O } 552\text{-}569\text{ cm}^{-1}$ ,  $18\text{-}17.6\text{ }\mu\text{m}$ ).

Zirconium diketonates such as  $[\text{Zr}(\text{acac})_4]$  are not suitable because their Zr-O stretching vibrations fall outside the range of an ir laser, ie in the region  $440\text{-}460\text{ cm}^{-1}$  ( $22.7\text{-}21.7\text{ }\mu\text{m}$ ). The Zr-O stretching vibrations of the analogous fluorine substituted compounds, eg  $[\text{Zr}(\text{hfacac})_4]$ , are however expected to be shifted to higher wavenumbers relative to  $\nu\text{Zr-O}$  for  $[\text{Zr}(\text{acac})_4]$ . This shift might bring the  $\nu\text{Zr-O}$  within the range of the ir laser. The zirconium-oxygen stretching and bending vibrations should be characterized for the fluorine substituted zirconium acetylacetonate complexes.

The Zr-O stretching bands in the ir spectra of zirconium carboxylates, eg  $[\text{Zr}(\text{O}_2\text{CR})_4]$ , also lie close to the range of an ir laser, however the compounds are not thermally stable. If the problem of poor thermal stability could be overcome, ie with different alkyl groups (R), the compounds should be tested with an ir laser.

The compounds  $[\text{Zr}(\text{NEt}_2)_4]$  and  $[\text{Zr}(\text{tfacac})_4]$  are commercially available/9/ and could quickly be prepared for testing with an ir laser.


### 3.2 Organometallic Chemistry

The designation organozirconium complex is given to a complex containing a direct zirconium to carbon link. Organozirconium compounds can be divided into four classes:


- i)  $\pi$ -cyclopentadienyls and arenes,
- ii) alkyls and aryls,
- iii) allyls,
- iv) carbonyls, nitrosyls and cyanides /7/.

All syntheses of organozirconium compounds must be carried out under dry, inert atmospheres, ie, under argon. Each class of compound will be reviewed briefly in the following section.

The most predominant organozirconium complexes are those which contain the cyclopentadienyl unit,  $(\eta^5\text{-C}_5\text{H}_5)_2\text{Zr}$ , eg  $\text{Cp}_2\text{ZrHCl}$ , where  $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ . The majority of cyclopentadienyl zirconium complexes are prepared from  $\text{Cp}_2\text{ZrCl}_2$ . The versatility of  $\text{Cp}_2\text{ZrCl}_2$  as a starting material is illustrated in Figure 2.  $\text{Cp}_2\text{ZrX}_2$ ,  $\text{X} = \text{Cl}, \text{Br}$  is prepared from zirconium tetrahalide and the cyclopentadienide Grignard reagent or alkali metal cyclopentadienide, Equations 7,8.



30% yield



81% yield

$\text{Cp}_2\text{ZrCl}_2$  has an essentially tetrahedral geometry. Its infrared spectrum is well characterized. The Zr-Cp stretching vibration occurs at  $358 \text{ cm}^{-1}$  ( $27.9 \mu\text{m}$ ) and the Zr-Cl stretching vibration is found at  $333 \text{ cm}^{-1}$  ( $30 \mu\text{m}$ )/14/. Both stretching frequencies are considerably out of the range of the infrared lasers commercially available at present.

$\text{Cp}_2\text{ZrX}_2$  compounds containing halides other than chlorides are best prepared from the commercially available  $\text{Cp}_2\text{ZrCl}_2$  via halide exchange reactions. Replacement of  $\text{C}_5\text{H}_5$  by  $\text{C}_5\text{H}_4\text{Me}$  and  $\text{C}_5\text{Me}_5$  enhances the solubility and crystallizability of  $[(\eta^5\text{-C}_5\text{H}_5)_2 \text{ZrL}_2]$  type complexes.






Figure 2. Reaction schemes for  $Cp_2ZrCl_2$ , demonstrating its utility as a starting material.


$Cp_2ZrCl_2$  is readily hydrolyzed in the presence of base to give  $Cp_2Zr(Cl)OZr(Cl)Cp_2$ , Equation 9 /14/.



The product, IV, is colourless, sublimes at 200-220°C in vacuo and is stable in air and light for periods greater than one month. It is soluble in all common organic solvents except saturated hydrocarbons. The infrared spectrum shows an intense absorption band in the range  $749-755 \text{ cm}^{-1}$  ( $13.4-13.3 \mu\text{m}$ ) due to the metal-oxygen-metal stretching vibration. The dimeric compound may not however be suitable for laser based isotope separation because of the complication of two zirconium atoms per molecule. The probability of the bimetallic complex having two Zr-91 atoms is very low, therefore the Zr-91 decomposition product may contain significant quantities of all other Zr isotopes. The yield of the Zr-91 depleted product would thereby be considerably reduced. Reaction of  $[Cp_2ZrCl]_2O$  with protic reagents (HX) yields the monomer  $Cp_2Zr(Cl)X$ , Equation 10 /13/.




A number of  $\text{Cp}_2\text{Zr}$  complexes containing oxygen bonded organic ligands have also been synthesized. These include alkoxides, phenoxides,  $\beta$ -diketonates, carboxylates, sulfinates and sulfonates. To illustrate,



Bis(cyclopentadienyl)alkoxides and phenoxides, eg  $[\text{Cp}_2\text{Zr}(\text{OC}_6\text{H}_5)_2]$ , are white crystalline compounds readily soluble in most organic solvents. In contrast to the tetraalkoxide derivatives discussed in Section 3.1, these species are not polymers. The ir spectra of these compounds have been well characterized. The C-O stretching vibrations for the phenoxides are found in the 1280-1300  $\text{cm}^{-1}$  (7.8-7.7  $\mu\text{m}$ ) region and in the 1090-1140  $\text{cm}^{-1}$  (9.2-8.8  $\mu\text{m}$ ) range for alkoxides. Bands between 730-765  $\text{cm}^{-1}$  (13.7-13.1  $\mu\text{m}$ ) have been tentatively assigned to Zr-O bending modes while those in the 420-570  $\text{cm}^{-1}$  (23.8-17.5  $\mu\text{m}$ ) region were considered to arise from Zr-O stretching vibrations/14/. These infrared bending and stretching vibrations of the biscyclopentadienylzirconium alkoxides fall within or close to the operating range of ir lasers. The synthesis of these compounds however requires a number of steps and yields vary from 20 to 98%. The compounds may however be worth testing just to see if laser based isotope separation is possible.


Reaction of  $\text{Cp}_2\text{ZrCl}_2$  with  $\beta$ -diketones (except the tri- or hexafluorinated derivatives) yields the bis( $\beta$ -diketonate) complex V. Equation 12 /14/.



V


94-98% yield

With either hexafluoro- or trifluoroacetylacetone the tris(diketonato) complex is obtained, Equation 13 /14/.




Zirconium-oxygen infrared stretching frequencies have not been reported for the acac, hfacac nor tfacac derivatives.

Reaction of  $\text{Cp}_2\text{ZrCl}_2$  with excess lithium borohydride yields  $\text{Cp}_2\text{Zr}(\text{BH}_4)_2$ . Treatment of  $\text{Cp}_2\text{Zr}(\text{BH}_4)_2$  with a tertiary amine, eg trimethylamine yields the dihydride, Equation 14 /14/.




$\text{Cp}_2\text{ZrH}_2$  is a polymer, nonvolatile and completely insoluble in all solvents. It is hydrolyzed only slowly in air. The zirconium-hydrogen stretching frequency occurs in the region  $1300\text{--}1540\text{ cm}^{-1}$  ( $7.79\text{--}6.59\text{ }\mu\text{m}$ ). The monohydride is similarly prepared, Equation 15 /14/.



The monohydride, in contrast to the dihydride, is a monomer, air reactive, sublimes in vacuo at  $60^\circ\text{C}$  and is soluble in organic solvents. The zirconium-hydrogen stretching frequency occurs at  $1945\text{ cm}^{-1}$  ( $5.1\text{ }\mu\text{m}$ ). The Zr-H vibration is found at the other end of the working range of an ir laser, relative to  $\nu\text{Zr-O}$ ,  $\nu\text{Zr-N}$  and  $\nu\text{Zr-Cl}$ .


Reaction of  $\text{Cp}_2\text{ZrCl}_2$  with allylmagnesium chloride yields the bisallyl complex, Equation 16 /14/.



#### VI


VI is unstable and decomposes after several hours at room temperature in light. The C=C stretching vibrations in the ir spectra,  $1533\text{ cm}^{-1}$  ( $6.5\text{ }\mu\text{m}$ ) and  $1589\text{ cm}^{-1}$  ( $6.3\text{ }\mu\text{m}$ ), suggest that one allyl ligand is  $\sigma$ -bonded and the other  $\pi$ -bonded to the metal/15/.

Sulfur and selenium derivatives are prepared as shown in Equation 17,



The sulfur and analogous selenium complexes are glassy solids, extremely air and moisture sensitive. The compounds are generally monomers when freshly prepared but in solution, or with aging in the solid state, polymerize with loss of solubility/14/.

Cyclopentadienyl zirconium complexes containing amido groups have also been reported. Reaction of tetrakis(dialkylamido)zirconium with a stoichiometric amount of cyclopentadiene yields the mono or biscyclopentadienyl derivative, Equation 18 /14/.



$x = 1, 2$

The amide derivatives can also be prepared via reaction between  $\text{Cp}_2\text{ZrX}_2$  ( $X = \text{Cl}, \text{Br}$ ) and a lithium or potassium amide. The physical and chemical properties of selected compounds are given in Table 6. Most of the compounds are air-sensitive and some are pyrophoric. The few Zr-N stretching bands which have been reported fall in the range  $500\text{--}640 \text{ cm}^{-1}$  ( $20\text{--}15 \mu\text{m}$ ).


The nitrate derivative,  $[\text{CpZr}(\text{NO}_3)_2]$ , is prepared by the reaction of tetrakis(cyclopentadienyl)zirconium with nitric acid. The compound is benzene soluble and explodes on heating to  $185^\circ\text{C}$  /15/. The mononitrate,  $\text{Cp}_2\text{Zr}(\text{Cl})\text{NO}_3$ , is more heat stable, mp  $190.5^\circ\text{C}$ . It is made from nitric acid and the dichloride. No ir data ( $\nu\text{Zr-N}$ ) are available.


Zirconium forms elimination stabilized alkyls,  $\text{ZrR}_4$ ,  $R = \text{benzyl}$  or allyl. The instability of zirconium alkyl and aryl derivatives is due to facile decomposition by low activation energy pathways such as  $\alpha$  and  $\beta$  elimination of zirconium hydride, homolysis, or coupling of ligands at the zirconium atom. The tetramethyl derivative,  $\text{Zr}(\text{CH}_3)_4$ , is unstable and decomposes at  $T < -15^\circ\text{C}$ . The neopentyl derivative,  $[\text{Zr}\{\text{CH}_2\text{C}(\text{CH}_3)_3\}_4]$ , mp  $103^\circ\text{C}$ , in contrast to  $[\text{Zr}(\text{CH}_3)_4]$  is a thermally stable tetraalkylated species. No ir data,  $\nu\text{Zr-C}$ , are available.

Few organometallic zirconium (III) compounds are known, no alkyl or aryl complexes have been reported. The only well-characterized Zr(III) complexes are the phosphido bridged dimers, VII, prepared as shown in Equation 19 /16/.

TABLE 6  
SELECTED AMIDO COMPLEXES OF ZIRCONIUM/14/


Compound	Colour	Bp or Sublimation Temp (°C)	$\nu_{\text{Zr-N}}$ $\text{cm}^{-1}$ ( $\mu\text{m}$ )	Yield
$\text{Cp}_2\text{Zr}[\text{N}(\text{CH}_3)_2]_3$	yellow, slightly viscous liquid	94-96/0.05 mm	-	33
$\text{Cp}_2\text{Zr}[\text{N}(\text{CH}_3)_2]_2$	yellow, shining crystals	110-120/0.05 mm	-	54
$\text{Cp}_2\text{Zr}[\text{N}(\text{C}_2\text{H}_5)_2]_2$	orange-yellow solid	120-130/0.03 mm	-	64
$\text{Cp}_2\text{Zr}[\text{N}(\text{C}_4\text{H}_5)_2]_2$	orange crystals	mp 180	507 (19.7)	56
$\text{Cp}_2\text{Zr} \left[ \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{N} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right]_2$	yellow-green	mp 195 (dec)	640 (15.6)	68
$\text{Cp}_2\text{Zr} \left[ \begin{array}{c} \text{---} \\ \diagup \quad \diagdown \\ \text{N} \\ \diagdown \quad \diagup \\ \text{---} \end{array} \right]_2$	orange solid	mp 59-62	550 (18.2)	38
$\text{Cp}_2\text{Zr}(\text{C}_{12}\text{H}_8\text{N})_2^*$	orange-red solid	mp 216-219	530 (18.9)	31





Compound VII is thermally stable, not readily soluble, sublimes under high vacuum and is rapidly oxidized by air,  $\text{I}_2$  and methyl iodide.

The only Zr(II) complex known is a dicarbonyl derivative prepared as shown,



The CO ir stretching frequencies occur at 1849 and 1945  $\text{cm}^{-1}$  (5.4 and 5.1  $\mu\text{m}$ , respectively).

A number of organozirconium complexes have been reviewed in this section. The majority of these compounds contain the unit  $\text{Cp}_2\text{Zr}$ . The synthesis of organometallic zirconium compounds requires a number of steps and yields are often low. Organometallic compounds which have an ir absorption within or close to the range of an ir laser, 9-16  $\mu\text{m}$  (1111-625  $\text{cm}^{-1}$ ), include

- biscyclopentadienylzirconium alkoxides,  $[\text{Cp}_2\text{Zr}(\text{OR})_2]$ ,  $\nu(\text{Zr-O})$  bend 730 - 765  $\text{cm}^{-1}$  (13.7 - 13.1  $\mu\text{m}$ ); and
- cyclopentadienyl(dialkylamido)zirconium complexes,  $[\text{Cp}_x\text{Zr}(\text{NR}_2)_{4-x}]$ ,  $\nu(\text{Zr-N})$  500-640  $\text{cm}^{-1}$ , 20 - 15  $\mu\text{m}$ .

In the case of the amido complexes, it may be possible, with careful selection of the alkyl groups to shift the Zr-N stretch to a value within the range of the ir laser. The amido compounds are however air sensitive and some are pyrophoric, which may cause handling problems.

The compound  $\text{Cp}_2\text{Zr}(\text{H})\text{BH}_4$ ,  $\nu(\text{Zr-H})$  1945  $\text{cm}^{-1}$  (5.1  $\mu\text{m}$ ), might in principle be a candidate for an ir laser isotope separation

process, but the Zr-H stretching vibration probably exhibits little isotopic discrimination since it is the relatively light hydrogen which vibrates and the heavy Zr scarcely moves.

Other candidates may be the  $\beta$ -diketonate derivatives of  $Cp_2ZrCl_2$ . The zirconium-oxygen infrared stretching frequencies have not however been reported for the acac, hfacac or tfacac derivatives. These should be examined and the suitability of the compounds reassessed.

#### 4.0 CONCLUSIONS AND RECOMMENDATIONS

The inorganic and organometallic chemistry of zirconium has been reviewed with the goal of identifying complexes which possess properties suitable for ir laser based zirconium isotope separation. The compounds identified as candidates, Sections 3.1 and 3.2, have Zr-ligand vibrations which occur very close to the outside edge of the range of an ir laser, 1111- 625  $cm^{-1}$  (9 - 16  $\mu m$ ). With careful selection of the composition of the ligands surrounding the zirconium, ie those not directly associated with the Zr-ligand vibration to be irradiated, it may be possible to shift  $\nu$ (Zr-ligand) farther into the practical range of an ir laser.

The candidate compounds are listed in Table 7. Of the compounds recommended for testing, those containing a Zr-O bond appear to be most promising. The extreme air sensitivity of the Zr-N containing species may necessitate more rigorous handling procedures than in the case of the Zr-O containing compounds.

Synthesis of a number of the candidate molecules requires many steps and product yields can be low, thereby making the compounds unattractive for industrial scale processes. All of the candidate molecules should however be tested, regardless of complexity and yields, in order to verify the ir laser isotope separation process. If the technique is viable, ways of improving product yields can be examined.

Candidate inorganic zirconium compounds outnumber organometallic species. Preparation of the inorganic compounds generally requires fewer steps than synthesis of organozirconium compounds. Organometallic compounds, specifically those containing the unit " $Cp_2Zr$ ", however, often have higher volatilities. This is an important property because the compounds must be in the gaseous state for irradiation.

An alternative to irradiating the Zr-ligand vibration may be to irradiate a bond whose vibration is influenced by the Zr atom, but does not directly involve Zr. To illustrate, irradiation of bond X-Y in the complex  $[L_xZr-X-Y]$  instead of bond Zr-X. If

TABLE 7

## SUMMARY OF CANDIDATE MOLECULES

Compound	R	Mp (°C)	Volatility (°C/mmHg)	IR Spectra		Comments	
				cm <sup>-1</sup>	$\nu(\text{Zr-O})$ ( $\mu\text{m}$ )		cm <sup>-1</sup>
ZrOCl <sub>2</sub> (anh)				877	(11.5)		
Zr(hfacac) <sub>4</sub> Zr(tfacac) <sub>4</sub>		130-131	N/A N/A			commercially available	
Zr(NR <sub>2</sub> ) <sub>4</sub>	Et nPr nBu	11q 11q 11q	110/0.01 165/0.1 180/0.1			557 (17.3) 637 (15.7) 637 (15.7)	commercially available
Zr(O <sub>2</sub> CR) <sub>4</sub>				650 470	(15.4) (21.3)		thermally unstable
Zr(SOCNR <sub>2</sub> ) <sub>4</sub>	Me Et			552- 569	(18- 17.6)		
Cp <sub>2</sub> Zr(OR) <sub>2</sub>	C <sub>6</sub> H <sub>5</sub> Me			730-765 420-520	(~13) bend (24-17) stretch		
Cp <sub>2</sub> Zr(acac)Cl				N/A			
CpZr(hfacac) <sub>3</sub> CpZr(tfacac) <sub>3</sub>				N/A N/A			
Cp <sub>2</sub> Zr(NR <sub>2</sub> ) <sub>2</sub>		195 (dec)				640 (15.6)	

anh = anhydrous

N/A = not available

the influence of the Zr atom on bond X-Y is sufficient to achieve isotope separation, then the number of candidate complexes would increase considerably.

Approved by:

Prepared by:



O.A. Kupcis  
Manager  
Chemical Research Department



A. Kuksis Kalvins  
Chemist  
Atmospheric Research Section

Reviewed by:



O.T. Melo  
Supervising Engineer  
Atmospheric Research Section

AK:lg/ljh

## REFERENCES

1. CRC Handbook of Chemistry and Physics, 63rd edition, 1982-1983, R.C. Weast, editor, CRC Press Inc., Boca Raton, Florida.
2. O'Neill, J.A., First Zirconium Isotope Separation Workshop: Summary of Proceedings, OHRD Report No 84-306-K.
3. Cox, D.M., R.B. Hall, J.A. Horsley, G.M. Kramer, P. Rabinowitz and A. Kaldor, *Science*, 205, 390 (1979).
4. Kaldor, A., R.B. Hall, D.M. Cox, J.A. Horsley, P. Rabinowitz and G.M. Kramer, *J. Am. Chem. Soc.*, 101, 4465 (1979).
5. Letokhov, V.S., *Nature* 277, 605 (1979).
6. A.J. Carty - communication
7. Cotton, F.A. and G. Wilkinson, *Advanced Inorganic Chemistry - A Comprehensive Text*, 4th ed., Wiley-Interscience, New York, 1980.
8. Bradley, D.C. and P. Thornton, Chapter 33, Zirconium and Hafnium, in *Comprehensive Inorganic Chemistry*, 3, 419 (1973), J. Bailar, ed., Pergamon Press, Great Britain.
9. Alfa Catalogue, Morton Thiokol Inc. 1983-1984, Danvers, Massachusetts.
10. Prozorovskaya, Z.N., K.I. Petrov and L.N. Komissarova, *Russ. J. Inorg. Chem.* 13, 505 (1968).
11. Chatteraj, S.C., C.T. Lynch and K.S. Mazdiasni, *Inorg. Chem.* 7, 2501 (1968).
12. Wengrovius, J.H., R.R. Schrock and C.S. Day, *Inorg. Chem.* 20, 1845 (1981).
13. Hawthorne, S.L., A.H. Bruder and R.C. Fay, *Inorg. Chem.* 17, 2114 (1978).
14. Wailes, P.C., R.S.P. Coutts and H. Weigold, *Organometallic Chemistry of Titanium, Zirconium and Hafnium*, Academic Press, New York, 1974.
15. Martin, H.A., P.J. Lemaire and F. Jellinek, *J. of Organometallic Chem.* 14, 149 (1968).

16. Gell, K.I., T.V. Harris and J. Schwartz, *Inorg. Chem.* 20,  
481 (1981).