

INSTITUTE FOR NUCLEAR STUDY  
UNIVERSITY OF TOKYO  
Tanashi, Tokyo 188  
Japan

INS-Rep.-746  
TMUP-HEL 8904  
April 1989

**The Universe as a Topological Defect  
in a Higher-Dimensional Einstein-Yang-Mills Theory**

**A. Nakamura**

Department of Physics, Tokyo Metropolitan University  
Setagaya, Tokyo 158, Japan

and

**K. Shiraishi**

Institute for Nuclear Study, University of Tokyo  
Midori-cho, Tanashi, Tokyo 188, Japan

The Universe as a Topological Defect  
in a Higher-Dimensional Einstein-Yang-Mills Theory

A. Nakamura

Department of Physics, Tokyo Metropolitan University  
Setagaya, Tokyo 158, Japan

and

K. Shiraishi\*

Institute for Nuclear Study, University of Tokyo  
Midori-cho, Tanashi, Tokyo 188, Japan

Abstract

An interpretation is suggested that a spontaneous compactification of space-time can be regarded as a topological defect in a higher-dimensional Einstein-Yang-Mills (EYM) theory. We start with D-dimensional EYM theory in our present analysis. A compactification leads to a D-2 dimensional effective action of Abelian gauge-Higgs theory. We find a "vortex" solution in the effective theory. Our universe appears to be confined in a center of a "vortex", which has D-4 large dimensions. In this paper we show an example with SU(2) symmetry in the original EYM theory, and the resulting solution is found to be equivalent to the "instanton-induced compactification". The cosmological implication is also mentioned.

---

\* JSPS Fellow.

Recently there has been much interest in the study of higher-dimensional theories. This is because the candidates for unified theory can be easily and naturally formulated in higher dimensions [1][2]. If the theories with more than four dimensions are to be taken seriously, a mechanism which brings about dynamical compactification of the extra dimensions is needed. Usually the size of the extra space is considered to be of order of the Planck length in order that we cannot observe the extra space experimentally in the existing laboratory. Various mechanisms of compactification are proposed in the literature, using a variety of "forces" to curl up the extra dimensions [1][2].

Rubakov and Shaposhnikov offered a novel mechanism [3]. They considered a self-interacting scalar theory in higher dimensions and discussed the possibility that "we live inside a domain wall". It is implicitly suggested that there are many three-dimensional "worlds" in the higher dimensions in their scenario. The possibility of "many worlds" is of great interest particularly in the cosmological context. However, their analysis did not include the connection to gravity. It is conceivable that no static solution coupled with gravity can be found and such an energetic structure, which is adequately measured in the Planck mass, will soon collapse. In addition, the existence of elementary scalar field with appropriate self-couplings has no compelling reason supported by unification

theories.

On the other hand, it is shown by Ref. [4] that gauge boson-Higgs scalar systems are derived from the dimensional reduction of higher-dimensional Yang-Mills theory. Unfortunately, they consider the dimensional reduction as a mere device to obtain various breaking patterns of Higgs mechanism.

Our scenario for compactification is the following: First we consider that the space-time is partially compactified in higher-dimensional Einstein-Yang-Mills (EYM) theory. Then we obtain an effective gauge-Higgs theory. Second, we consider a topologically non-trivial structure in the effective theory coupled to Einstein gravity. We suppose that stable static solutions can be constructed in this way.

Because we start with EYM theory and consider the configuration with non-trivial topology, it is hopeful to solve the fermion problem in Kaluza-Klein theory [5]. In fact, a simple example has been known. Simplest effective (neutral) scalar theory can be derived from SU(2) gauge theory compactified onto  $S^3$ , three-dimensional sphere [6]. There is an exact analytic solution of "kink" or domain wall. This solution turns out to be equivalent to "instanton-induced compactification" presented by Ranjbar-Daemi, Salam and Strathdee [7]. Accordingly, the stability has already been guaranteed in this case.

By imagining the two-step compactification, or "double

compactification", we anticipate some kind of phase transition in the early history of the universe. As is a usual Higgs potential, the effective potential is dependent on temperature and the deformation of the shape of the potential leads to phase transition [8]. During the phase transition, some topological defects are formed and we were born and live inside one of them.

In this paper we adopt SU(2) gauge theory coupled to Einstein gravity in D dimensional space-time as an example. First we consider compactification on  $S^2$ , two dimensional sphere, and then yields effective Abelian Higgs model in (D-2) dimensions. Next we construct a vortex solution with a unit-quantum of flux. The classical solution is coupled to gravity. Here the term "vortex" means that the topological defect has (D-5) dimensional spatial extension in (D-3) dimensional space; the extension is two dimension less than the space it can move.

In the following analysis, we set  $D=8$  in practice. The dimensionality of the resulting flat space-time we live in can be taken as an arbitrary number, but in this paper the dimensionality of the flat space-time is set to four.

We begin with the following Einstein-Yang-Mills action

$$S = \int d^D x \sqrt{-g} \left\{ -\frac{1}{2\kappa^2} R + \frac{1}{4e^2} \text{tr}(F_{MN}F^{MN}) + \Lambda \right\}. \quad (1)$$

Here  $\kappa^2=8\pi G$ ;  $G$  is Newton constant;  $e$  is a gauge coupling constant;  $\Lambda$  is a cosmological constant. The scalar curvature of  $S^N$  with unit radius is defined as  $R=N(N-1)$ . The suffices  $M$  and  $N$

run from 0 to 7.

The gauge symmetry group under consideration is SU(2). This may be regarded as a subgroup of a large unified symmetry group.

The field equations are Yang-Mills equation

$$D_n F^{nk} = \nabla_n F^{nk} + i[A_n, F^{nk}] = 0, \quad (2)$$

where the field strength is given by

$$F_{nk} = \partial_n A_k - \partial_k A_n + i[A_n, A_k]. \quad (3)$$

and Einstein equations

$$R_{MN} = \frac{\kappa^2 \Lambda}{3} g_{MN} + \kappa^2 \left( T_{MN} - \frac{1}{8} T g_{MN} \right), \quad (4)$$

where

$$T_{MN} = \frac{1}{e^2} \text{tr} \left( F_{MP} F_N^P - \frac{1}{4} F_{PQ} F^{PQ} g_{MN} \right), \quad (5)$$

$$\text{and } T = T^M_M. \quad (6)$$

Here  $R_{MN}$  is a Ricci tensor derived from the metric  $g_{MN}$ .

To solve the solution coupled to gravity, we take an ansatz for the form of the metric:

$$ds^2 = ds^2(M^4) + g^2(\rho) d\rho^2 + a^2(\rho) d\psi^2 + b^2(\rho) d\Omega^2(S^2), \quad (7)$$

where  $d\Omega^2(S^2) = d\theta^2 + \sin^2\theta d\phi^2$  and  $0 \leq \psi < 2\pi$ ,  $0 \leq \theta \leq \pi$  and  $0 \leq \phi < 2\pi$ . We assume the four dimensional space-time which is to be the dimensions we live in admits flat Minkowski metric.

The form of gauge fields on  $S^2$  is assumed to be

$$A_\theta = \Phi_1 \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} e^{i\psi} - i \frac{e^{-i\psi}}{0} \\ + \Phi_2 \frac{1}{2} \begin{pmatrix} 0 & 1 \\ e^{i\psi} & 0 \end{pmatrix}, \quad (8a)$$

$$A_\phi = -\Phi_1 \frac{1}{2} \begin{pmatrix} 0 & 1 \\ e^{i\psi} & 0 \end{pmatrix} \sin\theta \\ + \Phi_2 \frac{1}{2} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} e^{i\psi} \sin\theta \\ + \frac{1}{2} \begin{pmatrix} 1 - \cos\theta & 0 \\ 0 & -(1 + \cos\theta) \end{pmatrix}. \quad (8b)$$

Here  $\Phi_1$  and  $\Phi_2$  are the function of the coordinates but independent of the  $S^2$  coordinates  $\theta$  and  $\phi$ . Further, we assume the "U(1) gauge field" in six dimensional space-time, in general, of the form:

$$A_\mu = A_\mu(x^\mu) \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (9)$$

where  $A_\mu(x^\mu)$  depends on only the coordinate of six dimensions.  $\mu$  runs from 0 to 5. Note that here we use a coordinate basis associated with the metric and not an orthonormal one.

When the ansatz for the form of gauge configurations is substituted to the Yang-Mills equation (2), the equations of motion for  $\Phi_1$ ,  $\Phi_2$  and  $A_\mu$  closely resemble the equations of motion in Abelian Higgs model considered by Nielsen and Olesen [8]. The equations of motion are

$$D^\mu D_\mu \hat{\Phi} + \frac{1}{B^2} (1 - |\hat{\Phi}|^2) \hat{\Phi} = 0, \quad (10a)$$

$$\nabla_{\mu}(b^2 F^{\mu\nu}) + i(\dot{\Phi} \cdot D^{\nu} \dot{\Phi} - \dot{\Phi} D^{\nu} \dot{\Phi}) = 0, \quad (10b)$$

where  $\dot{\Phi} = \Phi_1 + i\Phi_2$  and  $F_{\mu\nu} = \nabla_{\mu} A_{\nu} - \nabla_{\nu} A_{\mu}$ . The covariant derivative is defined as  $D_{\mu} = \nabla_{\mu} + iA_{\mu}$  with  $\nabla_{\mu}$  is the covariant derivative associated with six-dimensional metric  $g_{\mu\nu}$ .

Of course the equations coincide with theirs when  $g^2=1$ ,  $a^2=\rho^2$  and  $b^2=\text{constant}$ . When gravity is coupled, the solution obtained by Nielsen and Olesen is modified except for near the origin,  $\rho=0$ . We impose some ansatz for the form of a vortex solution. They are the following [8]:

$$\dot{\Phi} = |\dot{\Phi}(\rho)| e^{i\psi}, \quad (11a)$$

$$A_{\mu} = 0 \text{ except for } A_{\psi}(\rho) \text{ (function on } \rho). \quad (11b)$$

Furthermore we use physically plausible properties to solve the equation: In the limit  $\rho \rightarrow \infty$ ,  $|\dot{\Phi}| \rightarrow 1$  and  $b \rightarrow 0$ , and at  $\rho \rightarrow 0$ ,  $g^2=1$ ,  $a^2=\rho^2$  (usual cylindrical coordinates) and the "magnetic flux"  $\int A_{\psi} d\psi \rightarrow 0$ .

At last, we find the following solution of vortex-type:

$$|\dot{\Phi}(\rho)| = \frac{\rho/B}{\sqrt{1+(\rho/B)^2}}, \quad (12a)$$

$$A_{\psi}(\rho) = \frac{1}{\sqrt{1+(\rho/B)^2}} - 1, \quad (12b)$$

$$g^2(\rho) = \frac{1}{(1+(\rho/B)^2)^2}, \quad (12c)$$

$$a^2(\rho) = \frac{\rho^2}{1+(\rho/B)^2}, \quad (12d)$$

$$b^2(\rho) = \frac{B^2}{1+(\rho/B)^2}, \quad (12e)$$

where  $B^2 = \kappa^2/(2e^2)$  and the flatness of the large dimensions realizes provided that  $\Lambda = 8e^2/\kappa^4$ . These algebraic relations remain unchanged when the dimensionality of the flat space-time left untouched varies.

The flux is quantized as expected:

$$\frac{1}{4\pi} \frac{1}{2\pi} \int_0^{2\pi} A_{\psi} d\psi = -1. \quad (13)$$

Note that here we obtain an analytic expression of the solution.

For the natural values of the couplings,  $B$  is small of order of the Planck length. Thus the size of the core of the vortex is nearly the Planck scale. The direction of  $\rho$  is contracted by gravitational effect, and then it becomes "compact". At first sight the metric of the solution is a little bizarre, but the geometry of the space spanned by  $\rho$ ,  $\psi$ ,  $\theta$  and  $\phi$  turns out to be the one of  $S^4$  after the rewriting of the coordinates.

Next, we examine the energy of the configuration of Yang-Mills fields. The energy is appreciated easily in the formulation of eight-dimensional Yang-Mills theory.

Soon one can find

$$F_{\rho\psi} = - \frac{\rho/B^2}{(1+(\rho/B)^2)^2} \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad (14a)$$

$$F_{\theta\phi} = \frac{\sin\theta}{1+(\rho/B)^2} \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad (14b)$$

$$F_{\rho\theta} = \frac{1/B}{(1+(\rho/B)^2)^{3/2}} \frac{1}{2} \begin{pmatrix} 0 & e^{-1(\phi-\psi)} \\ e^{i(\phi-\psi)} & 0 \end{pmatrix}, \quad (14c)$$

$$F_{\psi\phi} = \frac{\rho \sin\theta/B}{1+(\rho/B)^2} \frac{1}{2} \begin{pmatrix} 0 & e^{-1(\phi-\psi)} \\ e^{i(\phi-\psi)} & 0 \end{pmatrix}, \quad (14d)$$

$$F_{\rho\phi} = -\frac{\sin\theta/B}{(1+(\rho/B)^2)^{3/2}} \frac{1}{2} \begin{pmatrix} 0 & e^{-1(\phi-\psi)} \\ e^{i(\phi-\psi)} & 0 \end{pmatrix}, \quad (14e)$$

$$F_{\psi\theta} = \frac{\rho/B}{1+(\rho/B)^2} \frac{1}{2} \begin{pmatrix} 0 & e^{-1(\phi-\psi)} \\ e^{i(\phi-\psi)} & 0 \end{pmatrix}, \quad (14f)$$

If one utilizes the vierbeins to treat the suffices, he can immediately see the anti-self duality of the solution. Thus the energy density per unit three-dimensional large spatial volume is given by

$$\frac{1}{4e^2} \int d^4y \sqrt{g(y)} \text{tr} F^2 = \frac{1}{4e^2} \left| \int d^4y \text{tr} \tilde{F} F \right| = \frac{4\pi^2}{e^2}, \quad (15)$$

where  $y^a = \{\rho, \psi, \theta, \phi\}$  and  $g(y) = \det g_{ab}$ .

The stability of the solution is expected from this relation, though the full stability analysis is in need including the perturbation of the metric and the mixing with the modes from graviton. The relation in the form of the gauge configuration exhibits the equivalence of the well-known solution to the one of the "instanton-induced compactification" [7]. We emphasize, however, that when the technique of the type we showed is applied to an EYM theory with other gauge group and compactification, such as  $SU(m+1)$  on  $CP^m$  ( $m \geq 2$ ) [4], we can construct various type of solutions. We will not explore it further here.

In the model of the present type, we cannot take arbitrary values for coupling constants of the scalar  $\Phi$ . It can be read from the equation of motion that the effective "Higgs self-coupling" is the same as the gauge coupling up to an adequate normalization [9]. The fact suggests the existence of a static multi-vortex solution or a configuration of  $n$ -superimposed vortices. The study of multi-vortex system is the most important task when we wish to investigate a cosmological scenario (see later).

To conclude this paper, we should mention the cosmological implication. As mentioned before, we wish to consider "double compactification" as a dynamical phase transition, not merely a technical formulation of compactification. The investigation of the high-temperature phase of the model and the study of dynamical evolution of the scale factors are of great importance if one want to consider inflation or cosmological aspects.

Another remarkable concept is the possibility of "many worlds". If topological defects can copiously produced after the phase transition, many lower dimensional "worlds" can form networks. The event of "worlds in collision" may occur in higher dimensional space. In analogy with cosmic strings [10], self-crossing of a vortex as a "world" may lead to a "closed" world. The questions now arise: then does it collapse? If we live in the vortex, how about the effect of the collision or crossing on our world? The above questions must be considered not only by

analysing classical solutions of multi-defect system but also by including quantum effects of matter and gravity. The selection of the manifold on which the first-step compactification produces the gauge-Higgs system may also be concerned with quantum effects.

As a variation of the scenario, it can be considered that we live outside the defects and then we can find the extra space only inside the topological defect in our universe. This possibility is also worth studying.

If the solution obtained here is included in some series of solutions, we must find the method to obtain the series of the solutions and extend the solutions to apply various physical situations. We want to investigate other gauge configurations on other type of the compact manifolds in future works.

#### Acknowledgements

One of the authors (K.S.) would like to thank S. Saito and H. Yamakoshi for some comments.

This work is supported in part by the Grant-in-Aid for Encouragement of Young Scientist from the Ministry of Education, Science and Culture (# 63790150).

One of the authors (K.S.) is grateful to the Japan Society for the Promotion of Science for the fellowship. He also thanks Iwanami Fūjukai for financial aid.

#### References

- [1] T. Appelquist, A. Chodos and P. G. O. Freund, Modern Kaluza Klein Theories (Benjamin-Cummings, New York, 1987).
- [2] D. Bailin and A. Love, Rep. Prog. Phys. 50 (1987) 1087.
- [3] V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. 125B (1983) 136.
- [4] Yu. A. Kubyshin, J. M. Mourao and I. P. Volobujev, Phys. Lett. 203B (1988) 349.
- [5] E. Witten, in "Proceedings of the Shelter Island Conference" (MIT Press, Cambridge, Mass. 1985).
- [6] A. Nakamura and K. Shiraishi, preprint INS-Rep.-740 and TMUP-HEL 8902, March 1989.
- [7] S. Randjbar-Daemi, A. Salam and J. Strathdee, Nucl. Phys. B242 (1984) 447.
- [8] H. B. Nielsen and P. Olesen, Nucl. Phys. B81 (1973) 46.
- [9] L. Jacobs and C. Rebbi, Phys. Rev. B19 (1979) 4486.
- [10] A. Albrecht and N. Turok, preprint FERMILAB-Pub-89/42-A and PUPT-89-1119, March 1989, and references there in.