



IC/89/290

INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

THE COADJOINT ORBIT SPACES OF $\widehat{\text{Diff}}(S^1)$ AND TEICHMÜLLER SPACES

Subhashis Nag

and

Alberto Verjovsky



**INTERNATIONAL
ATOMIC ENERGY
AGENCY**



**UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION**

1989 MIRAMARE - TRIESTE



International Atomic Energy Agency
and
United Nations Educational Scientific and Cultural Organization
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**THE COADJOINT ORBIT SPACES OF $\widehat{\text{Diff}}(S^1)$
AND TEICHMÜLLER SPACES**

Subhashis Nag and Alberto Verjovsky
International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

Precisely two of the homogeneous spaces that appear as coadjoint orbits of the group of string reparametrizations ($\widehat{\text{Diff}}(S^1)$ ^{*)} carry in a natural way the structure of infinite dimensional, holomorphically homogeneous complex analytic Kähler manifolds. These are $N = \text{Diff}(S^1)/\text{Rot}(S^1)$ and $M = \text{Diff}(S^1)/\text{Möb}(S^1)$. Note that N is a holomorphic disc fiber space over M . Now, M can be naturally considered as embedded in the classical universal Teichmüller space $T(1)$, simply by noting that a diffeomorphism of S^1 is a quasisymmetric homeomorphism. $T(1)$ is itself a homomorphically homogeneous complex Banach manifold. We prove in the first part of the paper that the inclusion of M in $T(1)$ is *complex analytic*.

In the latter portion of this paper it is shown that the *unique* homogeneous Kähler metric carried by $M = \text{Diff}(S^1)/SL(2, \mathbb{R})$ induces precisely the *Weil-Petersson metric* on the Teichmüller spaces. This is via our identification of M as a holomorphic submanifold of universal Teichmüller space. Now recall that every Teichmüller space $T(G)$ of finite or infinite dimension is contained canonically and holomorphically within $T(1)$. Our computations allow us also to prove that every $T(G)$, G any infinite Fuchsian group, projects out of M *transversely*. This last assertion is related to the "fractal" nature of G -invariant quasicircles, and to Mostow rigidity on the line.

Our results thus connect the loop space approach to bosonic string theory with the sum-over moduli (Polyakov path integral) approach.

MIRAMARE - TRIESTE

September 1989

^{*)} The Virasoro group (see Witten [16]).

INTRODUCTION

PART I: *The complex structures*

The group $\text{Diff}(S^1)$ and its universal central extension, the Virasoro group, occurs in string theory as the space of reparametrizations of a closed string. Two coadjoint orbit spaces of $\text{Diff}(S^1)$, namely, $N = \text{Diff}(S^1)/S^1$ and $M = \text{Diff}(S^1)/SL(2, \mathbb{R})$, have occurred in the physics literature as critically important because precisely these two carry the structure of infinite dimensional, holomorphically homogeneous, complex (Kähler) manifolds (Witten [16]). The complex structures on these spaces are obtained by placing a natural physics-motivated almost complex structure (arising from Kirillov-Kostant representation theory) on the appropriate spaces of real vector fields on S^1 . This complex structure, arising from "conjugation" of Fourier series, appears, for example, in Pressley [18].

Now, considering diffeomorphisms of S^1 as quasimetric homeomorphisms one can naturally identify M as embedded in the classical universal Teichmüller space $T(1)$. $T(1)$ is a holomorphically homogeneous complex Banach domain from the famous Ahlfors-Bers theory of the Teichmüller spaces. Our first main result is that this inclusion of M into $T(1)$ is *complex analytic*. In fact, M is one leaf of a holomorphic foliation of $T(1)$. Also, since M_1 is a holomorphic disc fiber space over M , it seems to us that this naturally and directly connects the string reparametrization complex manifolds with the complex analytic moduli of Riemann surfaces. It appears to have been an important question (see, for example, Bowick [5], Bowick and Rajeev [6]) to relate these reparametrization spaces with the spaces of moduli of Riemann surfaces because that would connect the loop space ("geometrical quantization") approach to string theory with the path integral ("sum over moduli") approach. Indeed, $T(1)$ contains canonically within itself as complex submanifolds all the Teichmüller spaces of arbitrary Riemann surfaces or Fuchsian groups. If, therefore, strings are reparametrized using the more general quasimetric homeomorphisms of the circle (rather than only by smooth diffeomorphisms), then the corresponding $SL(2, \mathbb{R})$ orbit space is the universal Teichmüller space of Riemann surfaces.

Our method of proof is to show that the almost complex structure obtained by the physicists (Bowick and Rajeev [6], [7]; Bowick and Lahiri [8]) on real vector fields on S^1 modulo the Möbius vector fields coincides with the almost complex structure of $T(1)$ at the origin. The holomorphic homogeneity of both M_2 and $T(1)$ under the action of (right-) translation then implies that the complex structures are compatible everywhere.

PART II: *The Kähler structures*

The infinite dimensional holomorphically homogeneous complex manifold $M = \text{Diff}(S^1)/SL(2, \mathbb{R})$ is shown in Part I to be naturally embedded in universal Teichmüller space, $T(1)$, as one leaf of a holomorphic foliation of $T(1)$. That result showed that these two

“universal moduli spaces” are intimately related. There is a *unique* (up to a scaling factor) homogeneous Kähler metric, g , on M , this metric and its curvature have been studied intensively by many physicists including Bowick, Rajeev, Lahiri, Zumino, Kirillov (see [5], [6], [7],[8] [12],[13]). The chief result now is that this metric g produces precisely the *Weil-Petersson* Kähler metric on the Teichmüller spaces ^{*)}.

Let us be more precise. The metric g assigns a hermitian inner product on smooth real vector fields on the unit circle S^1 . (S^1 is to be thought of as the boundary of the open unit disk Δ .) Now, tangent vectors to the Teichmüller space are represented by Beltrami coefficients on Δ (modulo the infinitesimally trivial ones). The vector field on S^1 corresponding to a Beltrami coefficient μ is $\dot{w}[\mu](z) \frac{\partial}{\partial z}$, $z \in S^1$. We are able to express the metric g as a pairing on these Beltrami coefficients; indeed, the formula is:

$$(*) \quad g(\mu, \nu) = \iint_{\Delta} \iint_{\Delta} \frac{\mu(z) \overline{\nu(\zeta)}}{(1 - z\bar{\zeta})^4} d\xi d\eta \cdot dx dy$$

This formula converges (as it must) whenever $\mu, \nu \in L^\infty(\Delta)$ represent *smooth* ($C^{2+\epsilon}$ is enough smoothness) vector fields on S^1 . However, as it stands, (*) must diverge whenever μ, ν represent non-zero tangent vectors to any Teichmüller space $T(G) (\subset T(1))$, where G is any infinite Fuchsian group. This implies that each $T(G)$ sits in $T(1)$ *intersecting transversely* the leaves of the foliation of $T(1)$ by M and its $\text{Mod}(G)$ translates.

On the other hand, *the formula (*) for the physicists' metric g can still be used to recover the Weil-Petersson metric on any of the finite dimensional Teichmüller spaces $T(G)$* , (which is where the Weil-Petersson metric is defined classically). This is accomplished via a simple regularization of the improper integral (*) - as explained in Sec.II.3. Thus, the unique Kähler structure of $\text{Diff}(S^1)/SL(2, \mathbf{R})$ really does tie up with the Weil-Petersson Kähler metric on the Teichmüller spaces in a very convincing fashion.

The transversality we have mentioned above, of each $T(G)$ with M , relates intimately to various facts about the relationship of smooth diffeomorphisms to general quasimetric homeomorphisms (on S^1). Recall (see Sec.II.2) that the universal Teichmüller space $T(1)$ can be thought of as the space of all (Möbius-normalized) quasidisks. Now, the quasidisks corresponding to the analytic subset M are precisely the ones with C^∞ boundaries. Bowen [4] had proved that the boundary of any quasidisk corresponding to a (non-origin) point of $T(G)$, (where Δ/G is a compact Riemann surface), must be very non-smooth - indeed “fractal”. These results (discussed in Sec.II.4) are thus compatible with, and shed new light on, what we have proved in this paper.

^{*)} The Kähler form of the metric g is precisely the symplectic form that M carries by virtue of being a coadjoint orbit manifold. See Witten [16] and Kirillov [19].

PART I: THE COMPLEX STRUCTURES

I.1 THE COMPLEX STRUCTURE OF $\text{Diff}(S^1)/S^1$ AND $\text{Diff}(S^1)/\text{SL}(2, \mathbb{R})$

Let N and M denote respectively these two orbit spaces. We will think of them as right coset spaces. Using $S^1 = \text{Rot}(S^1)$ and $\text{SL}(2, \mathbb{R}) = \text{Möb}(S^1)$ to normalize a given C^∞ diffeomorphism (by following the given diffeomorphism by a normalizing one) we can identify N (and M) as those diffeomorphisms of S^1 that fix one (respectively, three) points of S^1 .

The Lie algebra of the Fréchet Lie group $\text{Diff}(S^1)$ is the algebra of C^∞ smooth real vector fields on S^1 (see Goodman [10]). The complexification of this Lie algebra is the Virasoro algebra generated by the $L_n = e^{in\theta} \frac{\partial}{\partial \theta} = iz^{n+1} \frac{\partial}{\partial z}$, $n \in \mathbb{Z}$. (Here $z = e^{i\theta}$.) A tangent vector to N at its origin is a linear combination:

$$\vartheta = \sum_{m \neq 0} \vartheta_m L_m, \quad \bar{\vartheta}_m = \vartheta_{-m}, \quad (1)$$

where $\vartheta = u(\theta) \frac{\partial}{\partial \theta}$ is the corresponding smooth real vector field on the circle and the ϑ_m are the Fourier coefficients of $u(\theta)$ (The ϑ_k decay faster than any negative power of k since $u(\theta)$ is C^∞ . See Katznelson [11], p.24.) For M , at its origin, a tangent vector will be of the form

$$\vartheta = \sum_{m \neq -1, 0, 1} \vartheta_m L_m, \quad \bar{\vartheta}_m = \vartheta_{-m} \quad (2)$$

Here one loses the coefficients $\vartheta_{-1}, \vartheta_0, \vartheta_1$ because an infinitesimal Möbius transformation of Δ (the unit disk) allows one to normalize precisely these coefficients. One may also check that the Lie algebra generated by L_{-1}, L_0, L_1 is precisely (the complexification of) $\mathfrak{sl}(2, \mathbb{R})$, as would be expected.

The natural *almost complex structure* \bar{J} at the origin of these two spaces is then defined (in each case) by (motivated from representation theory); see Auslander–Kostant [17]:

$$\bar{J}\vartheta = \sum_m -i \text{sgn}(m) \vartheta_m L_m. \quad (3)$$

See Pressley [18], Bowick and Rajeev [7], and Bowick and Lahiri [8]. The formula (3) is, of course, the classic formula known in the theory of Fourier series as “conjugation”. See, for example, Katznelson [11] Chapter III. One now follows [7], [8] to define the almost complex structure everywhere on these (right–) coset spaces M_i by right–translation invariance. As explained in [7] using the rather obvious involutivity of the (1,0) vector fields this \bar{J} is seen to be integrable and the right translations by elements of $\text{Diff}(S^1)$ act as biholomorphic automorphisms on N and M . (It is possible to get fairly explicit holomorphic coordinates on N and M as explained by Bruno Zumino in his July 1988 lectures at the ICTP.) See Zumino [20].

Remark We are purposely using right translations and right-invariant objects in order to finally coincide with the usual version of the theory of Teichmüller spaces. It is of course possible, as indicated in the last remark of the next section, to modify the definition of the Teichmüller spaces so that the left-invariant theory of the M and N works compatibly.

I.2 THE UNIVERSAL TEICHMÜLLER SPACE $T(1)$:

Let $\text{Homeo}_{qs}(S^1)$ denote the group of quasimetric homeomorphisms of the unit circle. These are the ones which allow some quasiconformal extension into the unit disc Δ bounded by S^1 . By a well-known characterization due to Ahlfors (see [1] or [14]) these are the homeomorphisms that alter cross ratios of points on S^1 by a bounded ratio. Now, Bers' universal Teichmüller space is

$$T(1) = \text{Homeo}_{qs}(S^1)/SL(2, \mathbb{R}) \quad (4)$$

Again, $SL(2, \mathbb{R}) = \text{Möb}(\Delta)$ can be thought of as normalizing a homeomorphism by following it by a Möbius transformation so that the composition fixes $+1, -1$ and $-i$ (say) on S^1 .

The complex analytic structure of $T(1)$ comes by thinking of it as equivalence classes of proper Beltrami coefficients on Δ . These Beltrami coefficients comprise the unit ball $L^\infty(\Delta)_1$ of the complex Banach space $L^\infty(\Delta)$. Given any $\mu \in L^\infty(\Delta)_1$ one solves the Beltrami equation

$$w_{\bar{z}} = \mu w_z \quad (5)$$

to get a quasi conformal self-homeomorphism $w = w_\mu$ of Δ . The boundary values of w_μ on S^1 (which always exist) is the quasimetric homeomorphism of S^1 representing the equivalence class $[\mu]$ in $T(\Delta)$. Thus,

$$T(1) = L^\infty(\Delta)_1 / \sim \quad (6)$$

where \sim is the equivalence relation saying $\mu \sim \nu$ if and only if w_μ and w_ν (normalized as explained by post-composition with Möbius transformations) have identical boundary values on S^1 .

Remark The way to get w_μ given μ in Δ is explained in Sec.II.2. Much more about the Teichmüller spaces will be needed in Part II of this paper – see the Section quoted above.

Bers proved that $T(1)$ inherits the structure of a complex Banach manifold from the complex structure of the unit ball $L^\infty(\Delta)_1$. Namely there is a unique induced complex structure on $T(1)$ such that the quotient projection $\Phi : L^\infty(\Delta)_1 \rightarrow T(1)$ becomes a *holomorphic submersion*. For complete proofs see Nag [14].

Notice that $T(1)$ is a group (though *not* a topological group). In fact, composition of quasimetric homeomorphisms corresponds to the following group law on Beltrami coefficients (see [14], p. 54–55 and p. 227–228)

$$\begin{aligned} \lambda \cdot \mu &= \text{Beltrami coefficient of } (w_\lambda \circ w_\mu) \\ &= \frac{\mu + (\lambda \circ w_\mu)\gamma_\mu}{1 + \bar{\mu}(\lambda \circ w_\mu)\gamma_\mu} \quad \text{where } \gamma_\mu = \frac{(w_\mu)_z}{(w_\mu)_{\bar{z}}} \end{aligned} \quad (7)$$

Since formula (7) depends holomorphically on λ we see that right translations act as biholomorphic automorphisms on $T(1)$ (and on $L^\infty(\Delta)_1$).

Remark If we redefine universal Teichmüller space by associating to $\mu \in L^\infty(\Delta)_1$ the boundary values of w_μ^{-1} , then the *left* translations act biholomorphically. The usual conventions in the physics literature regarding the orbit spaces of $\text{Diff}(S^1)$ can then be retained. We prefer to stick to the classical conventions in Teichmüller space theory.

I.3 $M \hookrightarrow T(\Delta)$ IS A HOLOMORPHIC INCLUSION:

It is well-known that every diffeomorphism of S^1 extends to a diffeomorphism of the closed disk $\Delta \cup S^1$. So diffeomorphisms are certainly quasimetric. Consequently, $M = \text{Diff}(S^1)/\text{Möb}(S^1)$ sits canonically inside $T(1) = \text{Homeo}_{qs}(S^1)/\text{Möb}(S^1)$.

Theorem I.1 The natural inclusion $M \hookrightarrow T(1)$ is holomorphic. M can be thought of as one leaf of a holomorphic foliation of $T(1)$ by injectively and holomorphically immersed leaves. (The leaves, which are copies of M , are not closed in $T(1)$.)

Using the holomorphic homogeneity of both M and $T(1)$ under right translations, one only needs to check the identity of the almost complex structures at the origin. The first problem is therefore to get a description of the almost complex structure, J , of $T(1)$ at the origin, (so as to be able to compare it with the \tilde{J} of Sec. I.1).

Acknowledgment: The pretty description of J on $T(1)$ given in the Proposition below is essentially an idea of S. Kerckhoff. The idea was explained to the first author in oral communication by C.J. Earle at Cornell University (1987–88).

A quasimetric real vector field on S^1 , to be thought of as an arbitrary tangent vector at the origin of $T(1)$, is obtained from a one-parameter flow of quasimetric homeomorphism $w_{t\mu}$, for any $\mu \in L^\infty(\Delta)$. The vector field on S^1 is then $\vartheta = \dot{w}[\mu] \frac{\partial}{\partial z}$ where $w_{t\mu}$ has the perturbation expansion:

$$w_{t\mu}(z) = z + t\dot{w}[\mu](z) + o(t), \quad t \rightarrow 0. \quad (8)$$

The problem is to obtain $J\vartheta$ on S^1 , where

$$J\vartheta = \dot{w}[i\mu](z) \frac{\partial}{\partial z}, \quad \text{given } \vartheta = \dot{w}[\mu](z) \frac{\partial}{\partial z}. \quad (9)$$

(Recall that the complex structure of $T(1)$ is inherited from the complex structure of the space of μ 's, as explained in Sec. I.2. So J corresponds to sending μ to $i\mu$.)

Proposition Using θ as coordinate on S^1 , $z = e^{i\theta}$, we can write $\vartheta = u(\theta) \frac{\partial}{\partial \theta}$ where $\dot{w}[\mu](z) = izu(z)$, $z \in S^1$. Then, $J\vartheta = u^*(\theta) \frac{\partial}{\partial \theta}$ where $\dot{w}[i\mu](z) = izu^*(z)$, $z \in S^1$. The formula for u^* is:

$$u^*(z) = \text{Im}(D(z)) + (cz + \bar{c}\bar{z} + b) \quad \text{on } S^1 \quad (10)$$

for a certain $b \in \mathbb{R}$, $c \in \mathbb{C}$. Here $D(z)$ is a member of the disc algebra $A(\Delta)$ (namely, functions holomorphic in Δ and continuous on $\Delta \cup S^1$) such that $\text{Re} D = u$ on S^1 .

Remark Notice that $u(z)$ is simply the *magnitude* of the vector field ϑ at the point $z \in S^1$. Note also that u^* is being shown to be essentially *the Hilbert transform* of u .

Proof The first variation term $\dot{w}[\mu]$ can actually be explicitly written down (see [14], or Eq.(24) of Part II here) in the form

$$\dot{w}[\mu](z) = \int \int_{\mathbb{C}} \bar{\mu}(\zeta) R(z, \zeta) d\zeta \wedge d\bar{\zeta} \quad (11)$$

where $\bar{\mu}$ is the extension of μ to the whole plane by reflection across S^1 , as explained in Part II, Sec.II.2. (Explicitly, $\bar{\mu}(\frac{1}{\bar{w}}) = \overline{\mu(w)} \frac{w^2}{\bar{w}^2}$ for w in Δ .) Here $R(z, \zeta)$ is a certain rational function. The main feature of $\dot{w}[\mu]$ (from which actually formula (11) can be derived) is that

$$\bar{\partial} \dot{w}[\mu] = \mu \text{ a.e. on } \Delta, \quad (\text{here } \bar{\partial} = \partial/\partial \bar{z}). \quad (12)$$

See Nag [14] p.39–40 and p.171 for a proof of this critical property. We also note for later use that formula (11) implies, since μ is L^∞ , that $u(z) = \frac{\dot{w}[\mu](z)}{iz}$ (on S^1) satisfies a Hölder condition $(|u(z_1) - u(z_2)| \leq c|z_1 - z_2|^\lambda, 0 < \lambda < 1)$ – in fact with λ arbitrarily close to 1.

Construct the function

$$F(z) = \dot{w}[i\mu](z) - \dot{w}[\mu](z), \text{ on } \Delta \cup S^1. \quad (13)$$

By formula (12) we see that $\bar{\partial} F = 0$ on Δ , so F is in the disc algebra $A(\Delta)$. Therefore the critical fact is:

$$izu^*(z) - zu(z) = F(z) \text{ on } S^1, \text{ for } F \in A(\Delta). \quad (14)$$

It is easy to derive the proposition from formula (14) as follows. Define $G \in A(\Delta)$ by $F(z) = F(0) + zG(z)$. Then (14) becomes

$$u(z) + iu^*(z) = G(z) + F(0)\bar{z} \text{ on } S^1. \quad (15)$$

We therefore have

$$\begin{aligned} u(z) &= \text{Re}(G(z) + F(0)\bar{z}) \text{ on } S^1 \\ &= \text{Re}(G(z) + \overline{F(0)}z) \text{ on } S^1 \end{aligned} \quad (16)$$

Of course, $D(z) = G(z) + \overline{F(0)}z$ is also in $A(\Delta)$, and the above calculation shows that $D(z)$ solves the “holomorphic Dirichlet problem” for the real boundary values u on S^1 .

Eq.(15) now allows us to relate u^* with u as desired:

$$\begin{aligned} u^*(z) &= \text{Im}(G(z) + F(0)\bar{z}), \text{ on } S^1 \\ &= \text{Im}(D(z) - \overline{F(0)}z + F(0)\bar{z}), \text{ on } S^1 \\ &= \text{Im}(D(z)) + kz + \bar{k}\bar{z}, \quad z = e^{i\theta}. \end{aligned} \quad (17)$$

Note that our $D(z)$ must be of the form

$$D(z) = \frac{1}{2\pi} \int_0^{2\pi} u(\theta) \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} \right) d\theta + ib, \quad b \in \mathbb{R},$$

(the ‘‘Schwarz kernel’’ formula for the Dirichlet problem). The fact that u is Hölder is well known (see, for example, Gakhov [9]) to guarantee that D is in $A(\Delta)$, and therefore $Im(D(z))$ on S^1 is well-defined.

The Proposition is fully proved. \square

Remark Notice that the normalization of w_μ (to fix +1, -1 and i) implies that the vector fields $u(\theta)$, $u^*(\theta)$ must vanish at these three points. The constants b, c occurring in formula (10) can be related at least partly to the enforcement of this normalization.

To prove our Theorem we need to show that J and \tilde{J} act identically on the smooth real vector fields ϑ of S^1 . Let, $\vartheta = u(\theta) \frac{\partial}{\partial \theta}$, with Fourier expansion

$$u(\theta) = \sum_{n=-\infty}^{\infty} a_n e^{in\theta} \quad (18)$$

Then,

$$D(z) = a_0 + 2 \sum_{n=1}^{\infty} a_n z^n \quad (19)$$

is in $A(\Delta)$ with $Re D = u$ on S^1 clearly.

By the Proposition we obtain therefore: $J\vartheta = u^*(\theta) \frac{\partial}{\partial \theta}$ where, for certain b, c

$$\begin{aligned} u^*(\theta) &= Im(D(e^{i\theta})) + (b + ce^{i\theta} + \bar{c}e^{-i\theta}) \\ &= \sum_{n=2}^{\infty} (-ia_n) e^{in\theta} + \sum_{n=2}^{\infty} \overline{(-ia_n)} e^{-in\theta} \\ &\quad + (\beta + \gamma e^{i\theta} + \bar{\gamma} e^{-i\theta}) \end{aligned} \quad (20)$$

The β and γ get normalized to zero via the $SL(2, \mathbb{R})$ normalization. Thus, comparing (20) with formula (3) for \tilde{J} , we see that $J\vartheta \equiv \tilde{J}\vartheta$. The theorem is proved. \square

PART II: THE KÄHLER STRUCTURES:

II.1 THE KÄHLER METRIC g ON M

Recall from Sec I.1 that the Lie algebra of the Frechet Lie group $Diff(S^1)$ consists of $Vect^\infty(S^1)$ (= smooth real vector fields on S^1). The homogeneous space of (right-) cosets $M =$

$\text{Diff}(S^1)/SL(2, \mathbb{R})$ has, as its tangent space at the origin, those smooth real vector fields v which are of the form

$$(1) \quad v = v(\theta) \frac{\partial}{\partial \theta} = \sum_{m \in \mathbb{Z} - \{-1, 0, 1\}} v_m L_m, \quad v_{-m} = \overline{v_m}$$

Here the $L_m = e^{im\theta} \frac{\partial}{\partial \theta}$, $m \in \mathbb{Z}$, generate the complexification of $\text{Vect}^\infty(S^1)$. Note that

$$(2) \quad [L_m, L_n] = i(n - m)L_{m+n}$$

There exists a *unique homogeneous Kähler metric* g on M , ([6],[7],[8]). By homogeneity, one needs to determine the Kähler form ω only at the origin of M . The requirement $d\omega = 0$ forces (see [7]):

$$(3) \quad \omega([L_m, L_n], L_p) + \omega([L_n, L_p], L_m) + \omega([L_p, L_m], L_n) = 0$$

Also, ω must vanish whenever one of its arguments is L_{-1}, L_0 or L_1 - since these vector fields (which generate $\mathfrak{sl}(2, \mathbb{R})$) give the zero tangent vector on M . From these conditions one finds that *the only possible homogeneous Kähler form* ω is given at the origin by:

$$(4) \quad \omega(L_m, L_n) = a(m^3 - m)\delta_{m,-n}, \quad m, n \in \mathbb{Z} - \{\pm 1, 0\}$$

(a is a non-zero complex constant.) Elsewhere on M , of course, ω is transported by translations. This Kähler form ω is precisely the Kirillov–Kostant symplectic form that exists on M since M is a coadjoint orbit of $\text{Diff}(S)$. Compare Witten [16].

Remark Interestingly, the formula (4) has occurred in more than one other context. Segal ([15], p.321) in calculating $H^2(\text{Vect}^\infty(S^1), \mathbb{C})$ gets (4), where (3) becomes interpreted as a cocycle condition. Again, (4) occurs in conformal field theory where the interpretation of (3) is basically as the Jacobi identity.

Proposition Let $v = \sum v_m L_m$ and $w = \sum w_m L_m$ (of the form (1)) represent two real tangent vectors to M at the origin. Then the Kähler metric g , whose Kähler form ω was determined above, assigns the inner product

$$(5) \quad g(v, w) = -2ia \text{Re} \left[\sum_{m=2}^{\infty} v_m \overline{w_m} (m^3 - m) \right]$$

The infinite series in (5) converges absolutely whenever the vector fields v and w are $C^{\frac{1}{2}+\epsilon}$ on S^1 (any $\epsilon > 0$). In particular C^2 vector fields produce convergence.

Proof Recall that the Kähler 2-form ω is related to the corresponding pairing g by $\omega(v, w) = g(v, \bar{J}w)$. So $g(v, w) = -\omega(v, \bar{J}w)$. It is therefore trivial to deduce (5) from (4). The convergence assertion follows from the fact that the Fourier coefficients of a $C^{k+\epsilon}$ function on S^1 decay at least

as fast as $\frac{1}{n^{2+\epsilon}}$. (See Katznelson [11], p. 24-25). Indeed, apply the Cauchy-Schwarz inequality to (5). Convergence is guaranteed if it is in ℓ^2 , and this happens when the vector field ϕ is $C^{\frac{3}{2}+\epsilon}$ smooth. \square

Note: Since we want $g(v, v) > 0$ we must have $a = ib$ with $b > 0$.

II.2 THE TEICHMÜLLER SPACES

We recall the basic facts we need from Teichmüller theory. See Nag [14] for complete proofs. This section is necessarily a trifle long since we have to understand the relation between various equivalent definitions of Teichmüller space.

The universal Teichmüller space, $T(1)$, is a holomorphically homogeneous complex Banach manifold which contains within itself, as complex submanifolds, all the Teichmüller spaces $T(G)$, (for arbitrary Fuchsian groups G operating on Δ). $T(G)$ parametrizes the various complex structures on the Riemann surface $X = \Delta/G$. The ball of proper Beltrami differentials, $L^\infty(\Delta)_1$, is fundamental: it is the open unit ball in the complex Banach space of L^∞ functions on Δ . The G -invariant Beltrami differentials constitute the closed complex subspace:

$$(6) \quad L^\infty(G) = \{ \mu \in L^\infty(\Delta) : \mu(gz)\overline{g'(z)}/g'(z) = \mu(z) \text{ a.e. on } \Delta \text{ for all } g \text{ in } G \}.$$

The unit ball $L^\infty(G) \cap L^\infty(\Delta)_1$ is denoted $L^\infty(G)_1$.

The chief construction is to solve the Beltrami equation

$$(7) \quad w_{\bar{z}} = \mu w_z$$

for any $\mu \in L^\infty(\Delta)_1$. One needs to look at two solutions of (7), namely,

$[w_\mu]$: The quasiconformal homeomorphism of \mathbb{C} which is μ -conformal (i.e. solves (7)) in Δ , fixes ± 1 and $-i$, and keeps Δ and Δ^* (= exterior of Δ) both invariant. This w_μ is obtained by applying the existence and uniqueness theorem of Ahlfors-Bers (for (7)) to the Beltrami coefficient which is μ on Δ and extended to Δ^* by reflection ($\tilde{\mu}(\frac{1}{\bar{z}}) = \overline{\mu(z)} z^2 / \bar{z}^2$ for $z \in \Delta$).

$[w^\mu]$: The quasiconformal homeomorphism on \mathbb{C} which is μ -conformal on Δ and conformal on Δ^* , (fixing ± 1 and $-i$ again, say). w^μ is obtained by applying the Ahlfors-Bers theorem to the Beltrami coefficient which is μ on Δ and zero on Δ^* .

Now one defines, for any Fuchsian group G including $G = \{1\}$, the *Teichmüller space*,

$$(8) \quad T(G) = L^\infty(G)_1 / \sim$$

where $\mu \sim \nu$ if and only if $w_\mu = w_\nu$ on $\partial\Delta = S^1$, which happens if and only if $w^\mu = w^\nu$ on $\Delta^* \cup S^1$. The space $T(G)$ carries uniquely the structure of a complex Banach manifold induced from the complex structure of the open ball $L^\infty(G)_1$ - i.e., such that the quotient projection

$$(9) \quad \Phi : L^\infty(G)_1 \rightarrow T(G)$$

becomes a *holomorphic submersion*.

How does this relate to moduli? If $\mu \in L^\infty(G)_1$ then w_μ conjugates G to another Fuchsian group

$$(10) \quad G_\mu = w_\mu G w_\mu^{-1}$$

The equivalence class of μ in $T(G)$ represents the Riemann surface $X_\mu = \Delta / G_\mu$.

Alternatively, one can utilize w^μ to conjugate G to a quasi-Fuchsian group

$$(11) \quad G^\mu = w^\mu G (w^\mu)^{-1}$$

so that G^μ operates discontinuously on the quasidisks $\Delta^\mu = w^\mu(\Delta)$ and its exterior $\Delta^{*\mu} = w^\mu(\Delta^*)$. X_μ is represented by Δ^μ / G^μ , (whereas $\Delta^{*\mu} / G^\mu$ is the fixed Riemann surface Δ^* / G – since w^μ was conformal on Δ^*).

From the w_μ picture it is clear that

$$(12) \quad T(G) = \left\{ \begin{array}{l} \text{quasisymmetric homeomorphisms of } S^1 \\ \text{compatible with } G \end{array} \right\} / \text{Möb}(S^1)$$

Here a homeomorphism f of S^1 is called quasisymmetric if it has some quasiconformal extension into Δ , and f is compatible with G if fGf^{-1} is again a group of (restriction to S^1 of) Möbius transformations. The relation between (8) and (12) is by associating to $\mu \in L^\infty(G)_1$ the homeomorphism w_μ restricted to S^1 .

Remark We want to draw attention to the fundamental fact that the moduli of two-dimensional Riemann surfaces are hereby determined by homeomorphisms of just the unit circle. *Thus reparametrizations of a closed string (S^1) are intimately related to moduli of complex structures.* Indeed, the homomorphic embedding of M in $T(1)$ exhibited in Part I is simply the inclusion $\text{Diff}(S^1) \hookrightarrow \text{Homeo}_{qs}(S^1)$.

Utilizing the w^μ picture, one can associate to the Teichmüller class of μ the quasidisk Δ^μ . Consequently:

$$(13) \quad T(1) = \left\{ \begin{array}{l} \text{quasidisks on the complex sphere} \\ \text{normalized by Möbius transformations} \end{array} \right\}$$

$T(G)$ comprises those quasidisks on which some quasi-Fuchsian conjugate of G acts discontinuously.

How does one pass back and forth between the descriptions (12) and (13) of Teichmüller space? Given a quasisymmetric homeomorphism f on S^1 , extend it to *any* quasiconformal homeomorphism F of $\Delta \cup S^1$. Then, if μ is the Beltrami coefficient $\bar{\partial}F/\partial F$ of F on Δ , the quasidisk corresponding to f is $D = \Delta^\mu$. D does *not* depend on the choice of the extension F .

Conversely, note that w_μ and w^μ are both μ -conformal on Δ – and consequently $\rho^\mu = w^{\mu\circ} w_\mu^{-1}$ is a (normalized) Riemann mapping of Δ onto Δ^μ . It follows that if a quasidisk D is supplied the corresponding quasisymmetric homeomorphism f on S^1 is

$$(14) \quad f = \rho^{-1} \circ \sigma$$

where ρ and σ are respectively the (normalized) Riemann mappings of Δ onto D and Δ^* onto D^* (= exterior of D).

We recall now Teichmüller's Lemma, which identifies the tangent space to any $T(G)$ at the origin. The fundamental pairing $L^\infty(G) \times A_2(G) \rightarrow \mathbb{C}$ between L^∞ Beltrami coefficients and integrable holomorphic quadratic differentials for G is given by

$$(15) \quad \langle \mu, \varphi \rangle = \iint_{\Delta/G} \mu \varphi.$$

Teichmüller's Lemma: Let $\Phi : L^\infty(G)_1 \rightarrow T(G)$ be the defining quotient projection (9). The kernel of the derivative of Φ at 0 is the subspace

$$(16) \quad N(G) = \{ \mu \in L^\infty(G) : \langle \mu, \varphi \rangle = 0, \text{ for all } \varphi \in A_2(G) \}$$

Consequently, the tangent space at origin of $T(G)$ is $L^\infty(G)/N(G)$. □

We are equipped now to define the *Weil-Petersson (W-P)* inner product on $T(G)$. Given $\mu, \nu \in L^\infty(G)$ one sets

$$(17) \quad \text{W-P}(\mu, \nu) = \iint_{\Delta/G} \times \iint_{\Delta} \frac{\mu(z) \overline{\nu(\zeta)}}{(1 - z\bar{\zeta})^4} d\xi d\eta \cdot dx dy$$

We want to explain this formula. From Ahlfors [1] one recalls the fundamental map

$$L^\infty(G) \rightarrow B_2(G)$$

given by

$$(18) \quad \nu \mapsto \varphi[\nu](z) = \iint_{\Delta} \frac{\overline{\nu(\zeta)}}{(1 - z\bar{\zeta})^4} d\xi d\eta$$

Here $B_2(G)$ are all the "Nehari-bounded" holomorphic quadratic differentials for G , (i.e. $\|\varphi(z)(1 - |z|^2)\|_\infty < \infty$ on Δ). The kernel of (18) is known to be precisely $N(G)$, and consequently the W-P inner product (17) becomes:

$$(19) \quad \text{W-P}(\mu, \nu) = \langle \mu, \varphi[\nu] \rangle$$

However, $\varphi[\nu](\in B_2(G))$ is not necessarily in $A_2(G)$ for general Fuchsian group G . So the W-P formula is usually defined only for the finite dimensional $T(G)$ - when Δ/G is on finite conformal type - because for such G one knows $B_2(G) \equiv A_2(G)$. For such G the formula (17), equivalently (19), converges.

IL3 THE KÄHLER METRIC ON $\text{Diff}(S^1)/\text{SL}(2, \mathbf{R})$ IS WEIL-PETERSSON

The main program is to calculate the physicists' Kähler metric on M in terms of Beltrami differentials. Let $\mu \in L^\infty(\Delta)$ represent any tangent vector to $T(1)$. The corresponding "quasisymmetric" vector field on S^1 is $v = \dot{w}[\mu](z) \frac{\partial}{\partial z}$ for $z = e^{i\theta}$. Consequently,

$$(20) \quad v(\theta) \frac{\partial}{\partial \theta} = \frac{\dot{w}[\mu](e^{i\theta})}{ie^{i\theta}} \frac{\partial}{\partial \theta}$$

Here $\dot{w}[\mu]$ is the first variation term in the solution theory of the Beltrami equation (7):

$$(21) \quad w_{t\mu}(z) = z + t\dot{w}[\mu](z) + o(t), \quad t \rightarrow 0.$$

Theorem II.1 Let $\mu, \nu \in L^\infty(\Delta)$ represent two tangent vectors at the origin of $T(1)$. Then the Bowick-Rajeev-Lahiri hermitian inner product g of formula (5) becomes formally:

$$(22) \quad g(\mu, \nu) = -\frac{ia}{3\pi^2} \iint_{\Delta} \times \iint_{\Delta} \frac{\mu(z)\overline{\nu(\zeta)}}{(1-z\bar{\zeta})^4} d\xi d\eta \cdot dx dy$$

This integration is convergent whenever μ and ν represent (see (20)) at least $C^{\frac{1}{2}+\epsilon}$ smooth vector fields on S^1 . (Thus C^2 vector fields will certainly produce convergence.)

Clearly we first need the Fourier coefficients of $v(\theta)$ so that we may apply formula (5). We have

Lemma $\mu \in L^\infty(\Delta)$ corresponds to the real vector field $v(\theta) \frac{\partial}{\partial \theta}$ on S^1 as per (20). The Fourier coefficients of $v(\theta)$ are:

$$(23) \quad v_k = \frac{i}{\pi} \iint_{\Delta} \overline{\mu(z)} \bar{z}^{k-2} dx dy, \quad \text{for } k \geq 2$$

and $v_k = \overline{v_{-k}}$ for $k \leq -2$.

Proof The explicit formula for $\dot{w}[\mu]$ is

$$(24) \quad \dot{w}[\mu](\zeta) = -\frac{(\zeta-1)(\zeta+1)(\zeta+i)}{\pi} \left\{ \iint_{\Delta} \frac{\mu(z) dx dy}{(z-1)(z+1)(z+i)(\zeta-z)} + \iint_{\Delta} \frac{i\overline{\mu(z)} dx dy}{(\bar{z}-1)(\bar{z}+1)(\bar{z}-i)(1-\zeta\bar{z})} \right\}$$

This is the formula of Sec. 1.2.12 of Nag [14] adapted to the disk. Or see formula (1.9) of Ahlfors [2]. Note that $\dot{w}[\mu]$ vanishes at ± 1 and $-i$, as it must by the normalization enforced on w_μ .

In (24) we substitute $\zeta = e^{i\theta}$ and make a straight calculation (in which some remarkable cancellations occur) to get the Fourier coefficients as desired.

□

Proof of the Theorem The coefficients v_k and w_k are determined respectively from μ and ν by the preceding Lemma. To apply formula (5) for the (hermitian) inner product on M we calculate

$$(25) \quad \sum_2^\infty \bar{v}_m w_m (m^3 - m) = -\frac{1}{\pi^2} \iint_{\Delta} \times \iint_{\Delta} \mu(z) \overline{\nu(\zeta)} \left(\sum_2^\infty z^{m-2} \bar{\zeta}^{m-2} (m^3 - m) \right) d\xi d\eta \cdot dx dy$$

We have interchanged order of sum and integral above. This is justified whenever there is absolute convergence – for example when the vector fields are $C^{2+\epsilon}$ as noted before.

Now,

$$(26) \quad \sum_{m=2}^\infty (m^3 - m) x^{m-2} = \frac{-1}{6(1-x)^4}, \text{ for } |x| < 1,$$

(as may be seen by differentiating $\sum_0^\infty x^m = (1-x)^{-1}$ three times). Using (5), (25), (26) we get the Theorem easily.

□

Remarks Since the choice of α is at our disposal (up to a positive multiple) we will henceforth normalize so that the factor outside the integral in (22) becomes unity. Notice that formula (22) is now precisely the W-P formula (17) for the case $G = \{1\}$, i.e. for universal Teichmüller space.

Corollary Let G be any infinite Fuchsian group. Then $T(G)$ intersects $M = \text{Diff}(S^1)/SL(2, \mathbb{R})$ transversely. Indeed, every non-null tangent vector to $T(G)$ at the origin, described by $\mu \in (L^\infty(G) - N(G))$ produces a vector field on S^1 that cannot be even $C^{\frac{1}{2}+\epsilon}$ smooth (for any $\epsilon > 0$). (Thus, $\dot{w}[\mu]$ cannot be C^2 on S^1 if $\mu \in (L^\infty(G) - N(G))$, whereas the tangent vectors to M are C^∞ smooth vector fields.)

Remark This Corollary is an infinitesimal form of Mostow rigidity on the line. See Sec. II.4 below.

Proof Formula (22) says that

$$(27) \quad g(\mu, \nu) = \iint_{\Delta} \mu \cdot \varphi[\nu]$$

where $\varphi[\nu]$ is defined in Ahlfors' map (18). But $\iint \mu \varphi$ over any union of N fundamental regions for G is $N\langle \mu, \varphi \rangle$, by G -invariance. Therefore, for $\mu, \nu \in (L^\infty(G) - N(G))$ formula (22) and (27) must diverge whenever G has infinitely many elements. All the assertions are now clear.

Remark If μ is a G -invariant Beltrami differential and φ is any (integrable) holomorphic function on Δ then the classic Poincaré series formula is

$$(28) \quad \iint_{\Delta} \mu \cdot \varphi = \iint_{\Delta/G} \mu \cdot \theta_2(\varphi),$$

where $\theta_2(\varphi) = \sum_{g \in G} (\varphi \circ g)(g')^2$. (See [14], p. 73, p. 174, p. 231, etc.)

But if $\nu \in L^\infty(G)$ then $\varphi[\nu]$ is *already* a G -invariant quadratic differential. So the Poincaré series is simply (order of G) $\times \varphi$, and the divergence of (27) is again manifest.

In spite of this divergence for $T(G)$ of the Bowick-Rajeev et al. metric, formula (22) one can still recover the W-P metric on every finite dimensional $T(G)$ from formula (22). The idea is to simply regulate passage to the limit in the improper integral. For $\mu, \nu \in L^\infty(G)$ rewrite (22) as

$$(29) \quad g(\mu, \nu) = \lim_{r \rightarrow 1^-} g_r(\mu, \nu)$$

where

$$(30) \quad g_r(\mu, \nu) = \iint_{D_r} \times \iint_{\Delta} \frac{\mu(z) \overline{\nu(\zeta)}}{(1 - z\bar{\zeta})^4} d\xi d\eta \cdot dx dy$$

or, equivalently, (see Eqs.(19) and (27) above),

$$(30') \quad g_r(\mu, \nu) = \iint_{D_r} \mu \cdot \varphi[\nu]$$

where $D_r = \Delta_r = \{|z| \leq r\}$, $r < 1$, is simply the disk of radius r .

Theorem II.2 The W-P inner product in any finite dimensional Teichmüller space $T(G)$, for $\mu, \nu \in L^\infty(G)$, is related to the Bowick-Rajeev inner product g by the formula:

$$(31) \quad \frac{W - P(\mu, \nu)}{W - P(\nu_0, \nu_0)} = \lim_{r \rightarrow 1^-} \frac{g_r(\mu, \nu)}{g_r(\nu_0, \nu_0)}$$

where $\nu_0 \in L^\infty(G) - N(G)$ represents any nonzero tangent vector to $T(G)$ at its origin.

Proof The group G is finitely generated of the first kind. If a function f on the hyperbolic unit disk Δ is C -automorphic then it is known that

$$(32) \quad \iint_{\Delta/G} f \cdot dA = \lim_{r \rightarrow \infty} \left(\frac{\iint_{B_r(z)} f \cdot dA}{\iint_{B_r(x)} dA} \right) \times \text{area}(\Delta/G)$$

where dA is the hyperbolic area element and $B_r(x)$ is the closed hyperbolic ball of radius r centered at any $x \in \Delta$. (In particular, note that the limit involved is independent of x .) The formula (32) follows from S.J. Patterson's results on the distribution in Δ of lattice points for G , this was kindly communicated to the first author by J. Velling.

Applying (32) to the automorphic function $f = \mu(z)\varphi[\nu](z)(1 - |z|^2)^2$ we see the theorem immediately. □

Remark The idea, of course, is that $g_r(\mu, \nu)$ is "essentially" $W - P(\mu, \nu)$ times the number of copies of a fundamental domain Δ/G sitting inside Δ_r . Note that the right-hand-side of (31) is an ∞/∞ form and L'Hospital's rule can be applied to advantage.

Remark Since the metric g was Kähler (Sec. II.1) to start with, this sheds new light on the Kählericity of the Weil-Petersson metric. That W-P is Kähler on $T(G)$ was first shown by Ahlfors in [2]. The negative curvature of W-P also fits in neatly with the corresponding fact known for g . Utilizing the natural relationship exhibited in formula (31) between the Kahler metric g on M and the usual Weil-Petersson metrics, one should now be able to *derive* the Kahlericity and the negative curvature of W-P from the corresponding facts for g .

II.4 ON THE TRANSVERSALITY OF $T(G)$ WITH M

The Corollary of Theorem II.1 of the previous section can be prettily interpreted by considering the deep question of which quasidisks correspond to points of M , and which to points of $T(G)$. We are now thinking of Teichmüller space as the set of (normalized) quasidisks - as explained in Sec. II.2.

Bowen [4] had proved that if G uniformizes a compact Riemann surface, then *every non-origin point of $T(G)$ corresponds to a quasidisk with fractal boundary* (i.e., Hausdorff dimension of the bounding quasicircle is strictly greater than unity).

On the other hand, *the quasidisks corresponding to the points of $M = \text{Diff}(S^1)/SL(2, \mathbf{R})$ are the ones with C^∞ boundaries.*

In fact, recall from Sec. II.2 the relationship between the two ways (12) and (13) of considering Teichmüller space. If the quasidisk is smoothly bounded then the Riemann mappings ρ and σ both extend smoothly to the boundaries. Hence, by formula (14) certainly the C^∞ quasidisks correspond to points of M . That they comprise *all* of M is shown in Kirillov [10] using an idea of Sullivan.

It is also easy to see that a G -compatible $f \in \text{Homeo}_{qs}(S^1)$ must be non-smooth (if f is not itself a Möbius transformation). For example, it is known that if f is compatible with a compact surface group G , and f is C^1 at even a single point of S^1 , then f must represent the origin of $T(G)$. One idea is to note that $fGf^{-1} = G_\mu$, and f assumed C^1 on the circle, implies, on direct

differentiation, that the hyperbolic marked length spectra of Δ/G and Δ/G_μ are the same. Hence G_μ is not a non-trivial deformation of G - so that f represents the origin of $T(G)$, as desired. These assertions are forms of Mostow rigidity on the line. See Agard [21].

We thus see that the fact that each $T(G)$ sticks transversely out of M is intimately related to the *non-smoothness* of the vector fields, or the quasisymmetric homeomorphisms, or the quasidisks corresponding to the (non-origin) points of $T(G)$.

Acknowledgments

One of the authors (S.N.) would like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste. Fruitful discussions with Caroline Series and James Eells are gratefully acknowledged.

REFERENCES

- [1] L. Ahlfors, *Lectures on Quasiconformal Mappings*, (Van Nostrand, Princeton, NJ, 1966).
- [2] L. Ahlfors, "Some remarks on Teichmüller's space of Riemann surfaces", *Annals of Math.* **74** (1961) 171–191.
- [3] A. Beardon, *The Geometry of Discrete Groups*, (Springer-Verlag, Berlin, 1983).
- [4] R. Bowen, "Hausdorff dimension of quasicircles", *Publ. Math IHES*, **50** (1979) 259-273.
- [5] M.J. Bowick, "The geometry of string theory", Eighth Workshop on Grand Unification, Syracuse, NY (MIT Preprint CTP No. 1492), (1987).
- [6] M.J. Bowick and S. Rajeev, "The holomorphic geometry of closed bosonic string theory and $\text{Diff}(S^1)/S^1$ ", *Nucl. Phys.* **B293** (1987) 348.
- [7] M.J. Bowick and S. Rajeev, "String theory as the Kähler geometry of loop space", *Phys. Rev. Letters* **58** (1987) 535–538.
- [8] M.J. Bowick and A. Lahiri, "The Ricci curvature of $\text{Diff}(S^1)/SL(2, \mathbb{R})$ ", Syracuse University preprint SU-4238-377 (February 1988).
- [9] F.D. Gakhov, *Boundary Value Problems*, (Pergamon Press, Oxford, 1966).
- [10] R. Goodman, "Positive energy representations of the group of diffeomorphisms of the circle", in *Infinite Dimensional Groups with Applications*, MSRI Berkley series No. 4, (Springer-Verlag, 1985).
- [11] Y. Katznelson, *An Introduction to Harmonic Analysis*, (John Wiley, New York, 1969).
- [12] A.A. Kirillov, "Kähler structures on K -orbits of the group of diffeomorphisms of a circle", *Funct. Anal. Appl.* **21** (1987) 122–125.
- [13] A.A. Kirillov and D.V. Yur'ev, "Kähler geometry on the infinite-dimensional homogeneous manifold $\text{Diff}_+(S^1)/\text{Rot}(S^1)$ ", *Funkt. Anal. Appl.*, **20** (1986) 322–324.
- [14] S. Nag, *The Complex Analytic Theory of Teichmüller Spaces*, (Wiley-Interscience, New York, 1988).
- [15] G. Segal, "Unitary representations of some infinite dimensional groups", *Comm. Math. Phys.* **80** (1981) 301–392.
- [16] E. Witten, "Coadjoint orbits of the Virasoro group", *Comm. Math. Phys.* **114** (1988), 1–53.
- [17] L. Auslander and B. Kostant, "Quantization and representations of solvable Lie groups", *Bull. Amer. Math. Soc.*, September 1967, 692–695.

- [18] A.N. Pressley, "*The energy flow on the loop space of a compact Lie group*", J. London Math. Soc. (2) 26 (1982) 557–566.
- [19] A. A. Kirillov, *Elements of the Theory of Representations* (Springer Verlag, Berlin, 1972), (Second Part, Articles 14 and 15).
- [20] B. Zumino, "*The geoemtry of the Virasoro group for physicists*", in *Cargise 1987*, Ed. R. Gastmans, (Plenum Press, New York and London), preprint LBL–24319, University of California, Berkley–PTH–87/48).
- [21] S. Agard, "*Mostow rigidity on the line*", *Holomorphic Functions and Moduli Proc. II*, MSRI Berkley (Springer, 1988), Ed. D. Drasin et al. pp. 1–12.

Stampato in proprio nella tipografia
del Centro Internazionale di Fisica Teorica