


**INTERNATIONAL CENTRE FOR  
THEORETICAL PHYSICS**



**SOME DISCUSSION ON THE ACCELERATION  
MECHANISM OF PARTICLES  
IN THE TYPE-I PLASMA COMET**

Li Zhongyuan

and

Guo Sheyu




**INTERNATIONAL  
ATOMIC ENERGY  
AGENCY**



**UNITED NATIONS  
EDUCATIONAL,  
SCIENTIFIC  
AND CULTURAL  
ORGANIZATION**

**MIRAMARE-TRIESTE**



International Atomic Energy Agency  
and  
United Nations Educational Scientific and Cultural Organization  
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

SOME DISCUSSION ON THE ACCELERATION MECHANISM  
OF PARTICLES IN THE TYPE-I PLASMA COMET

Li Zhongyuan \* and Guo Sheyu \*\*  
International Centre for Theoretical Physics, Trieste, Italy.

MIRAMARE – TRIESTE

July 1991

\* Permanent address: Department of Earth and Space Science, University of Science and Technology of China, Hefei 230026, People's Republic of China.

\*\* Permanent address: Institute of Plasma Physics, Hefei 230031, People's Republic of China.

1. Early, the large accelerations of plasma were already observed in type-I tail. The accelerations can be achieved  $300 \text{ cm/sec}^2$ . Recently, the direct measurements for comet G-Z showed that the energy of particles reaches  $2 \times 10^5 \text{ eV}$ . The energy is much higher than the initial energy of comet particles ( $\leq 2 \times 10^4 \text{ eV}$ ). So there should be an accelerated process in the comet.

The first order Fermi acceleration is the main accelerated mechanism to upper reaches particles in magnetosphere. Amata and Formisano have ever emphasized to point out that it is also a main accelerated mechanism in comet bow shock. We think it is possible that the acceleration exists in comet; but we also think that the Mach number of comet shock is very small ( $M \approx 2$ ), so the accelerated process in comet shock is not as effective as in magnetosphere shock. The reconnection of magnetic field lines also produces high energy particles possibly. But Ip has not agreed with that the reconnection is an effective accelerated mechanism in comet, (and it should also be discussed further, whether or not the steady-state reconnection occurs in any type-I comet). We support Ip's opinion, because some research showed us that the highest energy of comet particles,  $E_{max}$  (by "the reconnection") is only  $ZeU_A BL \ll 10^5 \text{ eV}$ . Besides, many people also put forward the concept of stochastic acceleration in coma, the second Fermi acceleration and so on. However, the knottiness of acceleration are still very much, and not solved.

We think the acceleration mechanisms may be varied and not be limited in one or two models. The comet can result in many kinds of plasma instabilities. There are many acceleration sources of particles in the comet, such as the non-steady-state reconnection of magnetic field, the couple of waves with particles, a lot of the turbulence forms of magnetic field and so on. They should be a synthetical process and effect. As below, we will also analyze the accelerated mechanisms in type-I comet, and it will be for the discussion further.

2. The flow lines in type-I comet are both the ion trace and the magnetic field lines. It is possible that the part plasma turbulences in comet may take transverse disturbed magnetic field to appear in some area. The disturbance perpendicular to magnetic field lines, will be able to hang these lines. Under the pressure of gas around, a magnetic neutral current sheet can be formed there. Fig.1 shows that the boundary clips the reversed polarity region (the magnetic diffusion region and the magnetic reconnection region) and cut the magnetic fields of reversed polarity into two. In the diffusion region, the magnetic lines take to diffuse. The magnetic fields annihilation occurs. In the reconnection region, the magnetic field lines merge again. The "annihilation" and the "reconnection" are nearly occurred in the same twinkling. We have already used the mechanism when discussing the type-I tail disruptions.

Both the dissipation and the convection take part in the reversed region. We should adopt the generalized Ohm law in the region,

$$\mathbf{E} + \frac{1}{c} \mathbf{v} \times \mathbf{B} = \eta \mathbf{j} + \frac{m_e}{ne^2} \nabla \cdot (\mathbf{j}_\perp + \mathbf{v} \mathbf{j}) - \frac{1}{ne} \nabla p + \frac{1}{ne} \mathbf{j} \times \mathbf{B} \quad (1)$$

Of course, the magnetic field in comet is till "frozen in" field except the reversed region, we may

use equation

$$\underline{E} + \frac{1}{c} \underline{v} \times \underline{B} = 0 \quad (2)$$

The mass equaiton of continuity may be written here

$$v_i \cdot \chi = v_z(x) \cdot Z(x), \quad (3)$$

where  $v_i$  is the velocity flowing into the reversed region,  $v_z$  is the velocity (the component  $x$ -direction) in the reversed region,  $Z(x)$  is the slow shock shape delineating the boundary of the reversed region. The component of  $x$ -direction of momentum flowing into the reversed region is  $[-\frac{d}{dx}(\rho v_x^2 Z)]$ . We suppose the plasma is unable to be compressed. Striding over the slow shock wave, the total pressure (magnetic pressure and plasma gas pressure) is continuous. The total pressure of the external field is well-distributed, therefore, the total pressure acting on the surface of the region is constant, it does not influence upon the momentum. Only the magnetic stress is able to give a contribution of  $\chi$ -direction of momentum to the surface, and the value is  $[-\frac{B_0 B_{on}}{4\pi} \Delta \chi]$ . Here,  $B_0$  is the external field,  $B_{on}$  is a component of the normal direction of  $B_0$ . Obviously, the magnetic field of the normal direction is continuous from beginning to end, therefore, we have

$$\frac{-B_0 B_{on}}{4\pi} \Delta \chi \simeq -\frac{B_0 B_z}{4\pi} \Delta \chi. \quad (4)$$

Thus, the momentum conservation requires

$$\frac{d}{dx}(\rho v_x^2 Z) = \frac{B_0 B_z}{4\pi}, \quad (5)$$

where,  $B_z$  is the internal field in the reversed region.

The plasmas will be unable as the ideal gas in the magnetic neutral current sheet. We should adopt the equation

$$\underline{E} = \underline{j}/\sigma - \frac{1}{c} \underline{j} \times \underline{B}. \quad (6)$$

In the diffusion region, the magnetic fields annihilate and the energy transforms. According to the Ampere law

$$\nabla \times \underline{B} = \frac{4\pi}{c} \underline{j}, \quad (7)$$

the magnetic energy flowing into the reversed region in the unit time can be gained easily

$$\begin{aligned} -\frac{c}{4\pi} \int \int \underline{E} \times \underline{B} \cdot \underline{n} ds &= \int \int \int j^2/\sigma + \frac{1}{c} \underline{j} \times \underline{B} \cdot \underline{v} d\tau \\ &= +\frac{cEB_0 X}{4\pi} \left[1 - \frac{8}{\pi} M_A G\right], \end{aligned} \quad (8)$$

where

$$G = \left[\frac{1}{2} \ell \pi (1 + L_z^2/X^2) + \frac{L_z}{2} \cot^{-1} \frac{L_z}{2}\right]. \quad (9)$$

The magnetic energy flowing out of the reversed region in the unit time as well, it is

$$\frac{cEB_0 X}{4\pi} \cdot \frac{8}{\pi} M_A \left[\frac{1}{2} \ell \pi (1 + \frac{L_z^2}{X^2}) + \frac{L_z}{X} \cot^{-1} \frac{L_z}{X}\right]. \quad (10)$$

Where  $L_z$  is the thickness of the reversed region,  $X$  is  $\frac{1}{4}$  of the diameter of the reversed region (see fig.2),  $M_A$  is the magnetic field merging rate.

Basing on Eqs.(1)-(5), we can get the precise solution as follows (here, the deduced process has been left out),

$$v_z = v_A M_A \chi / Z(x), \quad (11)$$

$$Z(x) = (M_A^2 \chi^2 + \lambda_e^2)^{1/2}, \quad (12)$$

$$B_z = B_0 M_A [2 M_A \chi / Z(x) - M_A^3 \chi^3 / Z(x)^3]. \quad (13)$$

Where  $v_A$  is Alfvén velocity,  $\lambda_e$  is the electron inertial length,

$$\lambda_e = c(\pi m_e / 4\pi n e^2)^{1/2}. \quad (14)$$

Eqs.(8), (9) and (10) show us that a part of the magnetic energy can be changed into Joule heat and dissipated, a part can be changed into the kinetic energy of plasma, and the rest of magnetic energy will flow out of the reversed region.

The non-steady-state reconnection process caused by the transverse magnetic field turbulences is stochastic. Through the process, the cometary particles will be able to be accelerated. In fact, the disturbed waves in the comet are Alfvén disturbances mainly. Given the Alfvén wavelength is  $\lambda$ . Moving through a distance  $\lambda$ , the Alfvén disturbed waves can make the magnetic neutral current sheet restore in terms of MHD principle. So the reconnection occurs in stochastic and restores by itself. Thus, it changes a part of the magnetic energy into the heat and kinetic energy of plasma in comet. Adopting the new data of comet G-Z, we can get the effect of the acceleration mechanism by the equation above. In our calculation, the effect results from the stochastic process is very small in one time. But the energy accumulative total will be effective if the magnetic field turbulences appear repeatedly. The energy of particles may reach  $5 \times 10^4$  eV and higher. This is nearly correspondence with the results observed in the comets. In general, it is not difficult that the kind of magnetic field turbulence process is occurred in the comets.

Of course, the above analysis is only a preliminary study, but it is a new idea. The mechanism can explain the acceleration phenomena in comets. The further work and a great quantity of the numerical simulation calculation has been considered, it will be a research of the next step.

3. Now we discuss the other possible acceleration mechanism basing on plasma dynamics again. When the solar wind encounters the plasmas of the comet, whether the "two stream" instability exists in the comet or not; there are different views about it. Now we discuss the problem here. The Volasov equation without collision is

$$\frac{\partial f_\alpha}{\partial t} + \underline{v} \cdot \nabla f_\alpha + \frac{e_\alpha}{m_\alpha} (\underline{E} + \frac{1}{c} \underline{v} \times \underline{B}) \cdot \nabla_v f_\alpha = 0. \quad (15)$$

Maxwellian equations are as follows,

$$\nabla \cdot \underline{E} = 4\pi\rho_E + 4\pi \sum_{\alpha} e_{\alpha} \int f_{\alpha} dv, \quad (16)$$

$$\nabla \times \underline{E} = -\frac{1}{c} \frac{\partial \underline{B}}{\partial t} \quad (17)$$

$$\nabla \cdot \underline{B} = 0, \quad (18)$$

$$\nabla \times \underline{B} = \frac{1}{c} \frac{\partial \underline{E}}{\partial t} + \frac{4\pi}{c} \underline{j} + \frac{4\pi}{c} \sum_{\alpha} e_{\alpha} \int f_{\alpha} \underline{v} dv. \quad (19)$$

These equations ((15)-(19)) are appropriate and complete. Where  $f_{\alpha}$  is particles distribution of component  $\alpha$ . It is the  $f_{\alpha 0}(v)$  (the distribution in equilibrium state) plus the  $f_{\alpha 1}(\tau, v, t)$  (the disturbance term),

$$f_{\alpha}(\underline{r}, \underline{v}, t) = f_{\alpha 0}(\underline{v}) + f_{\alpha 1}(\tau, u, t). \quad (20)$$

The electric field should be the similar form to  $f_{\alpha}$ , we have

$$\underline{E}(\underline{r}, t) = \underline{E}_0 + \underline{E}_1(\underline{r}, t). \quad (21)$$

Substituting the formulae (20) and (21) into (15), Eq. (15) can be linearized after the high order terms are dropped. We may get

$$\frac{\partial f_{\alpha 1}}{\partial t} + \underline{v} \cdot \nabla f_{\alpha 1} + \frac{e_{\alpha}}{m_{\alpha}} \underline{E}_0 \cdot \nabla_v f_{\alpha 1} = -\frac{e_{\alpha}}{m_{\alpha}} \underline{E}_1 \cdot \nabla_v f_{\alpha 0}. \quad (22)$$

Substituting the formulae (20) and (21) into Eqs. (16)-(19). Similarly, we can get the linearized equations about them. We omit to write them here.

Now we use the harmonious model and principle value integral. First we expand  $f_{\alpha 1}$  to Fourier function in time and space

$$f_{\alpha 1}(\underline{r}, \underline{v}, t) = \int \int f_{\alpha 1}(\underline{k}, \underline{v}, \omega) \cdot \exp(i\underline{k} \cdot \underline{r} - i\omega t) d\underline{k} d\omega. \quad (23)$$

Transforming  $\underline{E}_1 = -\nabla \phi_1$ , however, into the similar form like the formula (23), we have

$$\phi_1(\tau, t) = \int \int \phi_1(\underline{k}, \omega) \cdot \exp(i\underline{k} \cdot \underline{r} - i\omega t) d\underline{k} d\omega. \quad (24)$$

Substituting the forms (23) and (24) into the previous equations, and both the calculating them and the deducing them in detail we can get

$$\frac{1}{k^2} \int_{-\infty}^{+\infty} \frac{F_0'(v_x) dv_x}{v_x - \omega/k} = 1, \quad (25)$$

$$F_0'(v_x) = \int \int \sum_{\alpha} \left( \frac{4\pi n_{\alpha} e_{\alpha}^2}{m_{\alpha}} \right) f_{\alpha 0} dv_x dv_y \quad (26)$$

Here, Eq.(25) is the instability criterion of the plasmas. In order to apply more easily, the criterion (25) can be written the following equivalent equation,

$$U(u) = P \int_{-\infty}^{+\infty} \frac{F_0'(v_x) dv_x}{v_x - u} > 0. \quad (27)$$

where  $u = \omega/k$ ,  $P$  is the principle value. The plasmas are instability if  $U(u) > 0$ .

We adopt Maxwell distribution to positive (or negative) particles in the solar wind and comet. The main components of ions in comet are  $H_2O^+$ ,  $O^+$ ,  $CO^+$  and  $CO_2^+$ . The ions of solar wind are protons. Given  $v_0$ , the velocity of solar wind, and given  $\xi = v_x/v_0$ ; thus we have

$$F_0(v_x) = F_0(v_0\xi) = F(\xi). \quad (28)$$

If  $F'(\xi_0) = 0$ ,  $F''(\xi_0) > 0$ ; the criterion of plasma instability can be arrived at

$$U(\xi_0) = P \int_{-\infty}^{+\infty} \frac{F''}{\xi - \xi_0} d\xi > 0. \quad (29)$$

Of course, Eq.(29) is not a simple function. Adopting the obtained results of the comet G-Z and other some new data, we have calculated and got the curve of  $F(\xi)$  shown in Fig.3. From Fig.3, we know the minimum of  $F(\xi)$  is at  $\xi_0 = 0.873$ . At this point,  $U(\xi_0) > 0$ . So "two-stream" instability should exist in comet.

Hoyle et al. had shown the "two stream" instability was able to exist in comet. But they thought it is only a very brief period and a transient effect. Once it is occurred, it will be calmed down swiftly. Some scholars (Cheng et al.) thought there was not "two-stream" instability in the comets. In our view, the kind of plasma instability may occur in the inner of comet. We have calculated the data from transition region to tail axis, and discovered the occurs more easily if the distance to axis is closer. So the solar wind can transfer its energy into the ions of the comet and the ions will be able to be accelerated and heated. Especially, the acceleration mechanism is very effective when the solar active strengthen or the velocity of solar wind is speeded up.

#### Acknowledgments

The authors would like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste.

## REFERENCES

- Amata, E. and Formisano, V., *Planet Space Sci.*, vol. 33, p. 1243, 1985.
- Cheng, D., Liu, L. and Clifford Stoll, *Science in China, (Series A)*, No. 11, p. 1350, 1981.
- Hoyle, F. and Harwit, M., *Astrophys. J.*, Vol. 135, p. 867, 1962.
- Ip, W.-H. and Axford, W.I., in *Comet*'(ed. L.L. Wilkening), University of Arizona Press, p. 588, 1982.
- Ip, W.-H., in *Magnetotial Physics* (ed. A.T.Y. Lui), The Johns Hopkins University Press, p. 367, 1987.
- Ip, W.-H. and Axford, W.I., *Planet Space Sci.*, vol. 34, p. 1061, 1986.
- Li Zhongyuan, *Science in China (Series A)*, vol. 33, No. 4, p. 446, 1990.
- Li Zhongyuan, *Chinese J. of Space Sci.*, vol. 10, p. 96, 1990.
- Li Zhongyuan and Guo S., *Kexue Tongbao (Sci. Bul.)*, vol. 36, No. 6, p. 435, 1991.
- Mendis, D.A., et al., *Rev. Geophys. Space Sci.*, vol. 20, p. 883, 1982.
- Petschek, H.E. et al., *Astrophys. J.* vol. 149, p. 1157, 1967.
- Smith, E.J. et al., *Science*, Vol. 232, p. 382, 1986.
- Siscore, G.L., *Geophys. Res. Lett.*, vol. 13, p. 287, 1986.
- Von Rosenving, T.T. et al., *Science*, vol. 232, p.353, 1986.




Fig.1

The magnetic neutral current sheet caused by the transversal turbulence magnetic field.




Fig.2

A schematic diagram of the reversal region of the magnetic field.




Fig.3

A correlation curve of the function  $F(\xi)$  with the  $\xi$ .

