



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ
ДУБНА

JINR - E2-90-280

L. G. Zastavenko

THE HAMILTONIAN OF QED. ZERO MODE

Submitted to "International Journal
of Modern Physics A"

1990

1. The QED Lagrangian is

$$\mathcal{L} = -\frac{1}{4}(\partial_\mu A_\nu - \partial_\nu A_\mu)^2 - \psi^* \gamma_4 [\gamma_\mu (\partial_\mu - ieA_\mu) + m] \psi. \quad (1)$$

Let us introduce new variables \vec{B} , ϕ :

$$\vec{B} = \vec{A}(x, t) + \text{grad } \Lambda,$$

$$\Lambda(x, t, t_0) = \int_{t_0}^t A_0(x, \tau) d\tau, \quad (2)$$

$$\phi = e^{-ie\Lambda} \psi.$$

In these variables one has

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} (\vec{B})^2 - \frac{1}{2} (\text{rot } \vec{B})^2 - \\ & - \phi^* \gamma_4 \left[\sum_1^3 \gamma_j (\partial_j - ieB_j) + \gamma_4 \partial_4 + m \right] \phi. \end{aligned} \quad (3)$$

This Lagrangian gives the Hamiltonian

$$h = \int \left[\frac{1}{2} (\vec{\pi})^2 + \frac{1}{2} (\text{rot } \vec{B})^2 + \phi^* \gamma_4 \left(\sum_1^3 \gamma_j (\partial_j - ieB_j) + m \right) \phi \right] d^3x, \quad (4)$$

here

$$\pi_j(x) = -i\delta / \delta B_j(x), \quad j = 1, 2, 3,$$

(4a)

$$[\phi_\alpha^*(x), \phi_\beta(y)]_+ = \delta_{\alpha\beta} \delta(x-y).$$

Let us note that if

$$A_\mu(x, t) \rightarrow A'_\mu(x, t) = A_\mu(x, t) + e \partial \lambda(x, t) / \partial x_\mu, \quad \text{it} = x_4,$$

$$\psi \rightarrow \psi' = \psi e^{ie\lambda(x, t)},$$

then

$$\vec{B}(x, t) \rightarrow \vec{B}(x, t) + e \text{grad } \lambda(x, t_0), \quad (5)$$

$$\phi(x, t) \rightarrow \phi(x, t) e^{i e \lambda(x, t_0)}$$

The Hamiltonian h is invariant under transformation (5). Let us introduce a cubic periodicity volume V . Let us also introduce the decomposition

$$\vec{B}(x) = \text{grad } \gamma(x) + \vec{C}/\sqrt{V} + \vec{B}^{\text{tr}}(x), \quad (6)$$

$$\int \vec{B}^{\text{tr}} dx = \int \gamma dx = 0, \quad \vec{C} = \int \vec{B}(x) dx / \sqrt{V}, (\text{c.f. eq. (13)}).$$

The quantities \vec{C} , \vec{B}^{tr} do not change under transformation (5). Operator

$$\eta(x) = \phi(x) e^{-i e \gamma(x)} \quad (7)$$

gets a constant phase under this transformation. Let us consider the Schrödinger equation

$$(h - E)\Omega = 0. \quad (8)$$

Here

$$\Omega = \Omega(\phi, \phi^*, B), \quad (9)$$

see eqs. (4) and (4a). It is possible to express Ω as a function of variables $\eta, \eta^*, \vec{B}^{\text{tr}}, \vec{C}, \gamma$:

$$\Omega(\phi, \phi^*, B) = \bar{\Omega}(\eta, \eta^*, \vec{B}^{\text{tr}}, \vec{C}, \gamma). \quad (10)$$

Let us denote

$$h = h_1 + \frac{1}{2} \int (\vec{\pi})^2 dx. \quad (11)$$

One has

$$h_1 = h_1(\phi, \phi^*, \vec{B}) = h_1(\eta, \eta^*, \vec{B}^{\text{tr}} + \vec{C}/\sqrt{V}). \quad (12)$$

Equation (6) entails the decomposition

$$\vec{B}(x) = \frac{1}{\sqrt{V}} \left(\sum_{\substack{k \neq 0 \\ \lambda = 1, 2}} \vec{e}(k, \lambda) q_{\lambda}(k) e^{ikx} + \sum_{k \neq 0} ik q_3(k) e^{ikx} + \vec{C} \right). \quad (13)$$

We chose

$$\vec{e}(\mathbf{k}, \lambda) = \vec{e}(-\mathbf{k}, \lambda), \quad (\vec{e}(\mathbf{k}, \lambda) \cdot \mathbf{k}) = 0. \quad (14)$$

Then

$$-f(\vec{\pi})^2 dx = f\left(\frac{\delta}{\delta \vec{B}(\mathbf{x})}\right)^2 dx = \left(\frac{\partial}{\partial \vec{C}}\right)^2 + \sum_{\mathbf{k} \neq 0} \left(\frac{1}{k^2} \frac{\partial^2}{\partial \mathbf{q}_3(\mathbf{k}) \partial \mathbf{q}_3(-\mathbf{k})} + \sum_{\lambda=1,2} \frac{\partial^2}{\partial \mathbf{q}_\lambda(\mathbf{k}) \partial \mathbf{q}_\lambda(-\mathbf{k})}\right). \quad (15)$$

The derivative $\partial/\partial \mathbf{q}_3(\mathbf{k})$ acts on the functional $\bar{\Omega}(\eta, \eta^*, \vec{B}^{\text{tr}}, \vec{C}, \gamma)$ twice: on γ function and on η, η^* operators (7):

$$\begin{aligned} \frac{\partial}{\partial \mathbf{q}_3(\mathbf{k})} &= \left(\frac{\partial}{\partial \mathbf{q}_3(\mathbf{k})}\right) - ie \int ds \frac{\partial \gamma(s)}{\partial \mathbf{q}_3(\mathbf{k})} z(s) = \\ &= \left(\frac{\partial}{\partial \mathbf{q}_3(\mathbf{k})}\right) - ie \int ds e^{i\mathbf{k}s} z(\mathbf{x}) / \sqrt{V}, \end{aligned} \quad (16)$$

$$z(s) = \eta(s) \frac{\delta}{\delta \eta(s)} - \eta^*(s) \frac{\delta}{\delta \eta^*(s)}. \quad (17)$$

Let us consider, along with the operator $z(s)$, the operator $\rho(s)$,

$$\rho(s) = \eta^*(s) \eta(s). \quad (18)$$

One has

$$\begin{aligned} [\rho(s), \eta(y)] &= -\eta(s) \delta(s-y), \\ [\rho(s), \eta^*(y)] &= \eta^*(s) \delta(s-y), \\ [z(s), \eta(y)] &= \eta(y) \delta(s-y), \\ [z(s), \eta^*(y)] &= -\eta^*(y) \delta(s-y), \end{aligned} \quad (19)$$

Equation (19) provokes to expect

$$\rho(s) = -z(s). \quad (20)$$

Thus we have got new expression for the QED Hamiltonian:

$$\begin{aligned}
h = & \frac{1}{2} \sum_{\substack{k \neq 0 \\ \lambda = 1,2}} \left(- \frac{\partial^2}{\partial q_{\lambda}(k) \partial q_{\lambda}(-k)} + k^2 q_{\lambda}(k) q_{\lambda}(-k) \right) + \\
& + f \eta^* \gamma_4 \left(\sum_1^3 \gamma_j (\partial_j - ie B_j^{tr}) + m \right) \eta(x) d^3x - \\
& - \frac{1}{2} \left(\frac{\partial}{\partial \vec{C}} \right)^2 - ie \frac{\vec{C}}{\sqrt{V}} \int \eta^*(x) \gamma_4 \vec{\gamma} \eta(x) d^3x -
\end{aligned} \tag{21}$$

$$\begin{aligned}
& - \frac{1}{2} \sum_{k \neq 0} \frac{1}{k^2} \left[\frac{\partial^2}{\partial q_3(k) \partial q_3(-k)} + \right. \\
& + \left. \frac{2ie}{\sqrt{V}} \bar{\rho}(-k) \frac{\partial}{\partial q_3(k)} - \frac{e^2}{V} \bar{\rho}(k) \bar{\rho}(-k) \right], \\
& \bar{\rho}(x) = \int \rho(x) e^{ikx} d^3x.
\end{aligned} \tag{22}$$

Equation (21) essentially coincides with the standard QED Hamiltonian. It contains, however, an important new (zero mode) term,

$$- \frac{1}{2} (\partial / \partial \vec{C})^2 - e \vec{C} \int \eta^*(x) \vec{\alpha} \eta(x) d^3x / \sqrt{V}.$$

This term makes the Hamiltonian (21) unbounded from below (see a subsequent paper^{15/}).

We have got also a new derivation of the QED Hamiltonian. This derivation may be applied to the QCD.

Let us note that introduction of straight-forward cut-off in the Hamiltonian (21)

$$(\eta(x)) = \frac{1}{\sqrt{V}} \sum_{|p| < L} u_{p\sigma\lambda} e^{ipx} a_{p\sigma\lambda},$$

$$\vec{B}^{tr}(x) = \frac{1}{\sqrt{V}} \sum_{\substack{|k| < L \\ k \neq 0, \lambda = 1,2}} q_{\lambda}(k) \vec{e}_{\lambda}(k) e^{ikx}$$

does not contradict the Gauge invariance (see the remarks after eqs.(6) and (7)).

2. Now let me comment on the consideration of item 1. As a matter of fact, one can find the derivation of the QED Hamiltonian (mainly) in old textbooks (by Heitler^{11/}, Schiff^{12/},

Dirac ^{3/}, Wentzel^{4/}). All these derivations use, instead of eq. (6), the representation

$$\vec{B}(x) = \text{grad } \gamma + \text{rot } \vec{\delta}, \quad (6a)$$

$\text{rot } \vec{\delta} = \vec{B}^{\text{tr}}$, of the vector field $\vec{B}(x)$, thus omitting the term \vec{C}/\sqrt{V} which gives rise to new terms in the QED Hamiltonian.

This term was neglected in order to ensure the property $B(x) \rightarrow 0$ as $x \rightarrow \infty$ of the field $\vec{B}(x)$. Which of the representations, eq.(6) or eq.(6a) is correct? As much as I do understand, eq.(6) gives the standard commutation rule

$$[\pi_j(x), B_\ell(y)] = -i\delta(x-y)\delta_{j\ell}, \quad (4b)$$

while eq.(6a) gives a wrong commutator

$$[\pi_j(x), B_\ell(y)] = -i\delta_{j\ell} [\delta(x-y) - 1/V], \quad (4c)$$

for eq.(6a) implies the decomposition

$$B_j(y) = V^{-1/2} \sum_{k \neq 0} q_j(k) e^{ikx}.$$

2.1. Thus I dare think that all past derivations of the QED Hamiltonian possess an essential mistake.

2.2. Transformation (2) enabled me to avoid using a supplementary condition for elimination of the field $A_0(x, t)$. I am indebted to Dr. V.L.Ljuboshits who has explained me that transformation (2) is connected with the De Witt gauge.

2.3. I do understand that my proof of eq. (20) is not exhaustive.

2.4. Let me stress once again that it seems to be promising to apply the method of the present work to the problem of the QCD Hamiltonian derivation.

ACKNOWLEDGEMENT

I am deeply indebted to Dr. N.P.Ilieva, whose remark gave rise to this work. I am also indebted to Dr. G.N.Afanasiev and Professors B.M.Barbashov, G.V.Efimov, A.V.Efremov and V.N.Pervushin for discussions and criticism.

REFERENCES

1. Heitler W. - Quantum Theory of Radiation, Oxford, Clarendon Press, 1955.
2. Schiff L.I. - Quantum Mechanics, Mc.Graw-Hill Company INC., New-York - Toronto - London, 1955.
3. Dirac P.A.M. - The Principles of Quantum Mechanics, Oxford, Clarendon Press, 1958.
4. Wentzel G. - Quantum Theory of Fields, Interscience Publ. Inc., New York, 1949.
5. Zastavenko L.G. - JINR E2-90-281, Dubna, 1990.

Received by Publishing Department
on April 20, 1990.

WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Prices -- in US \$, including the packing and registered postage.

D18-85-793	Proceedings of the XII International Symposium on Nuclear Electronics, Dubna, 1985.	14.00
D4-85-851	Proceedings of the International School on Nuclear Structure Alushta, 1985.	11.00
D1,2-86-668	Proceedings of the VIII International Seminar on High Energy Physics Problems, Dubna, 1986 (2 volumes)	23.00
D8,4,17-86-747	Proceedings of the V International School on Neutron Physics. Alushta, 1986.	25.00
D9-87-105	Proceedings of the X All-Union Conference on Charged Particle Accelerators. Dubna, 1986 (2 volumes)	25.00
D7-87-38	Proceedings of the International School-Seminar on Heavy Ion Physics. Dubna, 1986.	25.00
D2-87-123	Proceedings of the Conference "Renormalization Group-86". Dubna, 1986.	12.00
D4-87-692	Proceedings of the International Conference on the Theory of Few Body and Quark-Hadronic Systems. Dubna, 1987.	12.00
D2-87-788	Proceedings of the VIII International Conference on the Problems of Quantum Field Theory. Alushta, 1987.	10.00
D14-87-799	Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1987.	13.00
D17-88-95	Proceedings of the IV International Symposium on Selected Topics in Statistical Mechanics. Dubna, 1987.	14.00
E1,2-88-426	Proceedings of the 1987 JINR-CERN School of Physics. Varna, Bulgaria, 1987.	14.00
D14-88-838	Proceedings of the International Workshop on Modern Trends in Activation Analysis in JINR. Dubna, 1988	8.00
D18-88-988	Proceedings of the XIII International Symposium on Nuclear Electronics. Varna, 1988	13.00
D10-89-70	Proceedings of the International School on the Problems of Use of Computers in Physical Research. Dubna, 1988	8.00
D9-89-52	Proceedings of the XI All-Union Conference on Charged Particle Accelerators. Dubna, 1988 (2 volumes)	30.00
D4,6,15-89-638	Proceedings on the International Conference on Selected Topics in Nuclear Structure. Dubna, 1989	14.00

Orders for the above-mentioned books can be sent at the address:
 Publishing Department, JINR
 Head Post Office, P.O.Box 79 101000 Moscow, USSR

**SUBJECT CATEGORIES
OF THE JINR PUBLICATIONS**

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Заставенко Л.Г.

E2-90-280

Гамильтониан КЭД. Нулевая мода

Мы исходим из стандартного лагранжиана КЭД. Дан новый вывод гамильтониана КЭД (важным новым моментом является учет нулевой моды). Предлагаемое рассмотрение безусловно с точки зрения калибровочной инвариантности. Оно дает важные поправки к стандартному гамильтониану КЭД. Можно надеяться распространить наше рассмотрение на случай КХД.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна 1990

Zastavenko L.G.

E2-90-280

The Hamiltonian of QED. Zero Mode

We start with the standard QED Lagrangian. New derivation of the spinor QED Hamiltonian is given. We have taken into account the zero mode. Our derivation is faultless from the point of view of gauge invariance. It gives important corrections to the standard QED Hamiltonian. Our derivation of the Hamiltonian can be generalized to the case of QCD.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna 1990

13 коп.

Редактор Э.В.Ивашкевич. Макет Т.Е.Попеко.
Набор Е.М.Граменицкой.

Подписано в печать 18.06.90.

Формат 60x90/16. Офсетная печать. Уч.-изд.листов 0,83.

Тираж 490. Заказ 43640.

Издательский отдел Объединенного института ядерных исследований.
Дубна Московской области.