



**СООБЩЕНИЯ  
ОБЪЕДИНЕННОГО  
ИНСТИТУТА  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ  
ДУБНА**

E2-91-19

V. V. Nesterenko

**CANONICAL QUANTIZATION  
OF A RELATIVISTIC PARTICLE  
WITH CURVATURE AND TORSION**

**1991**



## 1 Introduction

Recently interest has been aroused in the generalizations of relativistic particle model, the action of which depends not only on the length of the world curve but also on its curvature [1,3-6] or its torsion [2,7-9]. The models with curvature are used, for example, in the polymer theory [10]. And they can be treated as the one-dimensional version of the rigid string [11]. Investigation of the fermion-boson transmutations of the charged particles in an external Chern-Simons field reduced to the quantization of the relativistic particle with torsion [2,7]. So far the models containing only the curvature or torsion have been considered. For the completeness one has to investigate the model containing at the same time the curvature and torsion of the world trajectory. The present paper is devoted to this problem. The layout of the paper is as follows. In the second section the generalized Hamiltonian formalism for a relativistic particle with curvature and torsion in a  $D$ -dimensional space-time is constructed. A complete set of constraints in the phase space is found, their separation into the first class and the second-class constraints is fulfilled. The third section is devoted to the canonical quantization of this model. At first the general scheme of quantization in a  $D$ -dimensional space-time is considered. Further the case of a three-dimensional space-time is investigated in detail. In the sector with positive mass squared we obtain a spectrum determined by an equation involving the parameters of the model, the mass and the spin of a state. The possibility to describe in the framework of this model the states

with integer, half-odd-integer and continuous spins is shown ( $D = 3$ ). By making use of the Casimir operators of the Poincaré group we construct the wave equation and investigate it in detail in the rest frame. In the fourth section, the interaction with an external Abelian gauge field is introduced in the geometrical way. In Conclusion (Sect 5), the obtained results are briefly discussed and the problems waiting for their solutions are outlined. In Appendix some details of the calculation of the Poincaré group invariant  $W$  on the constraint surface are presented.

## 2 Generalized Hamiltonian formalism for relativistic particle with curvature and torsion

We shall investigate the model defined by the action

$$S = -m \int ds - \alpha \int k(s) ds - \beta \int \kappa(s) ds, \quad (2.1)$$

where  $k(s)$  is a curvature of the world curve of the particle,  $\kappa(s)$  is a torsion of this curve,  $m$  is a constant with the mass dimension,  $\alpha$  and  $\beta$  are dimensionless constants. If  $x^\mu(\tau)$ ,  $\mu = 0, 1, \dots, D-1$  is a parametric representation of the world trajectory, then the action (2.1) can be rewritten in the form [15]

$$S = -m \int d\tau \sqrt{\dot{\mathbf{x}}^2} - \alpha \int d\tau \frac{\sqrt{\bar{g}}}{\dot{\mathbf{x}}^2} - \beta \int d\tau \frac{\sqrt{\dot{\mathbf{x}}^2} d}{\bar{g}}, \quad (2.2)$$

where

$$\dot{\mathbf{x}} \equiv d\mathbf{x}/d\tau, \quad \bar{g} = (\dot{\mathbf{x}}\dot{\mathbf{x}})^2 - \dot{\mathbf{x}}^2 \ddot{\mathbf{x}}^2, \quad d = \det(d_{\alpha\beta}),$$

$$d_{\alpha\beta} = x^{(\alpha)} x_{\beta}, \quad \dot{\mathbf{x}} \equiv d^\alpha x/d\tau^\alpha, \quad \alpha, \beta = 1, 2, 3.$$

In the  $D$ -dimensional space-time the metric with the signature  $(+, -, \dots, -)$  is used.

To eliminate the superlight velocities in the model under consideration, we assume that  $\dot{\mathbf{x}}^2 > 0$ . Putting  $x^0(\tau) = \tau$  we can deduce from here the following conditions  $\ddot{\mathbf{x}}^2 < 0$  and  $\dot{\mathbf{x}}^2 < 0$ . If all these conditions are satisfied then the radicands in eq. (2.2) are positive.

The Lagrangian function in action (2.2) depends on  $\dot{\mathbf{x}}$ ,  $\ddot{\mathbf{x}}$  and  $\bar{\mathbf{x}}$ . Therefore the resulting Euler-Lagrange equations are ordinary differential equations of the sixth order. The action (2.2) is invariant under the Poincaré transformations in the ambient space-time and under the reparametrization  $\bar{\tau} = f(\tau)$ . As a consequence, the Lagrangian in this model is singular. Let us construct the generalized Hamiltonian description of this model following the papers [1, 2, 13]. To begin with we introduce the canonical variables

$$q_1 = \mathbf{x}, \quad q_2 = \dot{\mathbf{x}}, \quad q_3 = \ddot{\mathbf{x}}, \quad (2.3)$$

$$p_1 = -\frac{\partial L}{\partial \dot{x}} - \frac{dp_2}{d\tau}, \quad (2.4)$$

$$p_2 = -\frac{\partial L}{\partial \dot{x}} - \frac{dp_3}{d\tau}, \quad (2.5)$$

$$p_3 = -\frac{\partial L}{\partial \dot{x}}, \quad (2.6)$$

where  $L$  is the Lagrangian function in (2.2). The Lorentz indices in eqs (2.3)-(2.6) are omitted for simplicity. We shall do so further if misunderstanding does not appear.

We shall need the explicit form of the canonical momentum  $p_3$  only. It is given by

$$p_3^\mu = \beta \frac{\sqrt{\dot{x}^2}}{g} \sqrt{d} \sum_{\alpha=1}^3 d^{3\alpha} x^\mu, \quad (2.7)$$

where  $d^{\alpha\beta}$  is the matrix inverse to  $d_{\alpha\beta}$ :  $d_{\alpha\beta} d^{\beta\gamma} = \delta_\alpha^\gamma$ . From (2.7) we deduce three primary constraints

$$\phi_1^{(1)} = p_3^2 + \beta^2 \frac{q_2^2}{g} = 0, \quad (2.8)$$

$$\phi_2^{(1)} = p_3 q_2 = 0, \quad (2.9)$$

$$\phi_3^{(1)} = p_3 q_3 = 0, \quad (2.10)$$

where  $g = (q_2 q_3)^2 - q_2^2 q_3^2$ . They have the same form as primary constraints in the theory of the relativistic particle with torsion [2].

The Poisson brackets will be defined as follows

$$\{f, g\} = \sum_{\alpha=1}^3 \left( \frac{\partial f}{\partial p_\alpha^\mu} \frac{\partial g}{\partial q_{\alpha\mu}} - \frac{\partial f}{\partial q_\alpha^\mu} \frac{\partial g}{\partial p_{\alpha\mu}} \right). \quad (2.11)$$

The primary constraints (2.8)-(2.10) are mutually in involution

$$\{\phi_1^{(1)}, \phi_2^{(1)}\} = 0, \quad \{\phi_1^{(1)}, \phi_3^{(1)}\} = 2 \phi_1^{(1)} \approx 0, \quad \{\phi_2^{(1)}, \phi_3^{(1)}\} = \phi_2^{(1)} \approx 0. \quad (2.12)$$

The sign  $\approx$  means a weak equality [14]. The canonical Hamiltonian is

$$H = -p_1 \dot{x} - p_2 \dot{x} - p_3 \dot{x} - L = -p_1 q_2 - p_2 q_3 + m \sqrt{\dot{x}^2} + \alpha \frac{\sqrt{g}}{q_2^2}. \quad (2.13)$$

The equations of motion in the phase space are written as follows

$$\frac{df}{d\tau} = \frac{\partial f}{\partial \tau} + \{f, H\} + \sum_{\alpha=1}^3 \lambda_\alpha \{f, \phi_\alpha^{(1)}\}, \quad (2.14)$$

where  $f$  is a function of the canonical variables and evolution parameter  $\tau$ .

We now proceed to find the secondary constraints by making use of the Dirac method [1,14]. Demanding the stationarity of the primary constraints

$$\frac{d}{d\tau} \phi_a^{(1)} = \{ \phi_a^{(1)}, H_T \} \approx 0, \quad a = 1, 2, 3, \quad (2.15)$$

where  $H_T = H + \sum_{b=1}^3 \lambda_b \phi_b^{(1)}$  we obtain three new constraints

$$\phi_1^{(2)} = p_2 p_3 - \beta^2 \frac{q_2 q_3}{g} = 0, \quad (2.16)$$

$$\phi_2^{(2)} = p_2 q_2 = 0, \quad (2.17)$$

$$\phi_3^{(2)} = p_2 q_3 - \frac{\alpha}{q_2^2} \sqrt{g} = 0. \quad (2.18)$$

The requirement of the stationarity of the constraints (2.16)–(2.18) on the equations of motion

$$\frac{d}{d\tau} \phi_a^{(2)} = \{ \phi_a^{(2)}, H \} + \sum_{b=1}^3 \lambda_b \{ \phi_a^{(2)}, \phi_b^{(1)} \} \approx 0, \quad a = 1, 2, 3 \quad (2.19)$$

results in three additional constraints with the canonical Hamiltonian (2.16) become them

$$\phi_1^{(3)} = p_1 p_3 + p_2^2 + \alpha \frac{p_2 q_3}{\sqrt{g}} - \beta^2 \frac{q_3^2}{g} = 0, \quad (2.20)$$

$$\phi_2^{(3)} = H - p_1 q_2 - p_2 q_3 + m \sqrt{q_2^2} + \frac{\alpha}{q_2^2} \sqrt{g} = 0, \quad (2.21)$$

$$\phi_3^{(3)} = -p_1 q_3 + m \frac{q_3^2}{\sqrt{q_2^2}} = 0. \quad (2.22)$$

At this stage the process of generation of constraints is complete. The requirement of the stationarity of the last constraints (2.20)–(2.22) enables us to determine the Lagrange multipliers  $\lambda_1$  and  $\lambda_2$  in the total Hamiltonian

$$H_T = m \frac{(q_3 q_3)^2 - \alpha q_2^2 q_3^2}{2(p_1 p_3) q_3^2 \sqrt{q_2^2}} - \lambda_1 \frac{p_1 p_3}{p_1 p_3} \quad (2.23)$$

Now we have to solve all the constraints (1)–(3) and (2)–(3) constraints. For this purpose we construct the matrix  $M$  with the elements

$$M_{ij} = \{ \phi_i, \phi_j \} = \{ \phi_i, \phi_j \} \quad (2.24)$$

where  $\theta_{3(b-1)+a} = \phi_a^{(b)}$ ,  $a, b = 1, 2, 3$ . The matrix  $\Omega$  can be rewritten in a block form

$$\Omega = \begin{pmatrix} 0 & 0 & A \\ 0 & B & C \\ -A^t & -C^t & D \end{pmatrix}, \quad (2.25)$$

where 0 is the (3x3)-zero matrix and

$$A = \begin{pmatrix} 0 & 0 & -2p_1 p_3 \\ 0 & 0 & 0 \\ -p_1 p_3 & 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 & -p_1 p_2 \\ 0 & 0 & 0 \\ p_1 p_3 & 0 & 0 \end{pmatrix}, \quad (2.26)$$

$$C = \begin{pmatrix} c & 0 & -p_1 p_2 \\ -2p_1 p_3 & 0 & 0 \\ -2p_1 p_2 & 0 & \frac{mg}{\sqrt{q_1^2 p_3^2}} \end{pmatrix}, \quad D = \begin{pmatrix} 0 & -3p_1 p_2 & d \\ 3p_1 p_2 & 0 & \frac{mg}{q_2^2 \sqrt{q_2^2}} \\ -d & -\frac{mg}{q_2^2 \sqrt{q_2^2}} & 0 \end{pmatrix},$$

where

$$c = \beta^2 \frac{p_1 q_2}{g} - \alpha \frac{p_1 p_3}{\sqrt{g}} + 2\alpha \frac{p_2^3 q_2^2}{q_2^2 \sqrt{g}},$$

$$d = m^2 - p_1^2 + \alpha \frac{m\sqrt{g}}{(q_2^2)^{3/2}}.$$

The sign  $^t$  means a transposition. As known [15], the number of the first class constraints equals  $\text{Dim ker } \Omega$ . If the vector  $\xi \in \text{ker } \Omega$ ,  $\xi = \{\xi_1, \dots, \xi_9\}$  then

$$\xi_4 = \xi_6 = \xi_7 = \xi_8 = 0,$$

$$(p_1 p_3) \xi_3 + 2(p_1 p_3) \xi_5 - 3(p_1 p_2) \xi_8 = 0, \quad (2.27)$$

$$2(p_1 p_3) \xi_1 - \frac{mg}{q_2^2 \sqrt{q_2^2}} \xi_8 = 0.$$

Thus we get

$$\text{Dim ker } \Omega = 3.$$

Therefore in the model under consideration there are three first-class constraints and six second-class constraints. The number of physical degrees of freedom equals obviously  $3D - 6$ . The first-class constraints can be separated by the formula [15]

$$\Phi_a = \sum_{i=1}^9 \zeta_i^{(a)} \theta_i, \quad a = 1, 2, 3, \quad (2.28)$$

where  $\zeta_i^{(a)}$ ,  $a = 1, 2, 3$  are the basis vectors of  $\text{ker } \Omega$ . By making use of eq.(2.27) one can easily construct these vectors up to an arbitrary factor for each  $\zeta_i^{(a)}$ . They have the following nonzero components

$$\zeta_2^{(1)} = 1; \quad \zeta_3^{(2)} = -2; \quad \zeta_8^{(2)} = 1,$$

$$\xi_1^{(3)} = \frac{mg}{2q_2^2 \sqrt{q_2^2 p_1 p_3}}, \quad \xi_3^{(3)} = 2 \frac{p_1 p_2}{p_1 p_3}, \quad \xi_8^{(3)} = 1. \quad (2.29)$$

Taking into account (2.28) and (2.29) we obtain the first-class constraints

$$\Phi_1 = p_3 q_2 = 0, \quad (2.30)$$

$$\Phi_2 = p_2 q_2 - 2p_3 q_3 = 0, \quad (2.31)$$

$$\Phi_3 = \frac{mg}{2q_2^2 \sqrt{q_2^2 p_1 p_3}} \phi_1^{(1)} + 2 \frac{p_1 p_2}{p_1 p_3} \phi_3^{(1)} + H = 0. \quad (2.32)$$

As the second-class constraints  $\omega_s$ ,  $s = 1, \dots, 6$  one can take six arbitrary constraints from the set  $\{\theta_i, i = 1, \dots, 9\}$  with  $\det \|\{\omega_s, \omega_{s'}\}\| \neq 0$ ,  $s, s' = 1, \dots, 6$ . This can be done in many ways. For example, one may put

$$\omega_a = \phi_a^{(2)} = 0, \quad \omega_{3+a} = \phi_a^{(3)} = 0, \quad a = 1, 2, 3. \quad (2.33)$$

In this case the Hamiltonian (2.13) is considered to be the second-class constraint.

However we can substitute  $H = \omega_c$  in (2.33) by  $\phi_1^{(1)}$ . At the quantum level we shall consider both these possibilities.

### 3 Quantum theory

At first we consider the general scheme of the canonical quantization of this model in the  $D$ -dimensional space-time. We are dealing with a generalized Hamiltonian system in the  $6D$ -dimensional phase space with three first-class constraints (2.30)-(2.32) and six second-class constraints (2.33). The state vectors will be defined by the conditions

$$\Phi_a |\psi\rangle = 0, \quad a = 1, 2, 3. \quad (3.1)$$

The commutators of the operators  $q_a$  and  $p_a$ ,  $a = 1, 2, 3$  should be determined by the Dirac brackets constructed by means of the second-class constraints  $\omega_s$ . After this the constraints  $\omega_s$  will vanish at the quantum level identically. Therefore they can be omitted in conditions (3.1). As a result, the wave equations (3.1) can be rewritten only in terms of the primary constraints

$$\phi_a^{(1)} |\psi\rangle = 0, \quad a = 1, 2, 3. \quad (3.2)$$

If the canonical Hamiltonian is substituted in the set of the second-class constraints by  $\phi_1^{(1)}$ , then the same substitution will take place in the wave equations (3.2).

The number of the wave equations (3.2) can be reduced by introducing the gauge conditions. For example, the condition

$$X_1 = q_2 q_3 = 0 \quad (3.3)$$



details considerable simplification. From (3.3) it follows that

$$q_2^2 = \text{const.} \quad (3.4)$$

Thus eq (3.3) is, in fact, the proper time gauge. This gauge eliminates completely the functional freedom in the equations of motion (2.14), and the last Lagrange multiplier turns out to be

$$\lambda_2 = q_3^2/q_2^2. \quad (3.5)$$

In principle, we can impose one or two gauge conditions in addition to (3.3)

$$\lambda_c(q_a, p_a, \tau) = 0, \quad c = 2, 3$$

demanding that

$$\det \|\{\chi_a, \Phi_b\}\| \neq 0, \quad a, b = 1, 2, 3, \quad (3.6)$$

$$\frac{\partial \chi_c}{\partial \tau} + \{\chi_c, H\} + \sum_{a=1}^3 \lambda_a \{\chi_c, \phi_a^{(1)}\} = 0, \quad c = 2, 3, \quad (3.7)$$

where  $\lambda_a$ ,  $a = 1, 2, 3$  are determined in (2.33) and (3.5).

Further simplification is achieved when  $D = 3$ . In this case three vectors  $q_2, q_3, p_3$  form, by virtue of the constraints (2.8) (2.10) and gauge condition (3.3), (3.4), a complete orthogonal basis. The velocity  $q_2^\mu$  is a time-like vector while the acceleration  $q_3^\mu$  and the momentum  $p_3^\mu$  are the space-like vectors. The constraints  $\phi_a^{(2)} = 0$ ,  $a = 1, 2, 3$  enables us to obtain in this basis the expansion for the momentum  $p_2^\mu$  in the form

$$p_2^\mu = \alpha \frac{q_3^\mu}{\sqrt{-q_2^2 q_3^2}}, \quad \mu = 0, 1, 2. \quad (3.8)$$

The constraints  $\phi_a^{(3)} = 0$ ,  $a = 1, 2, 3$  and gauge condition (3.3) determine the projection of the momentum  $p_1^\mu$  on the basis vectors  $q_2^\mu$ ,  $q_3^\mu$  and  $p_3^\mu$

$$p_1 \sigma_2 = m \sqrt{q_2^2}, \quad p_1 q_3 = 0, \quad p_1 p_3 = \beta^2 / q_2^2. \quad (3.9)$$

From here we deduce

$$p_1^\mu = q_2^\mu \frac{m}{\sqrt{q_2^2}} + p_3^\mu \frac{q_3^2}{q_2^2}. \quad (3.10)$$

Squaring eq. (3.10) we obtain

$$p_1^2 = M^2 = m^2 + \beta^2 \frac{q_3^2}{(q_2^2)^2}, \quad (3.11)$$

where  $M^2$  is the mass of the particle with the action (2.1). From (3.11) it follows that

$$M^2 < m^2 \quad (3.12)$$

and  $M^2$  is not positive definite because  $q_3^2 < 0$ . Thus in the model under consideration the squared mass is determined by the initial conditions ( by the Cauchy data ) for variables  $q_3^\mu$  and it can be either positive, or negative, or it can vanish. <sup>1</sup> Further we shall confine our consideration to the sector in this model, where  $p_1^2 = M^2 > 0$ .

Let us examine the angular momentum in this model

$$M_{\mu\nu} = \sum_{\alpha=1}^3 (q_{\alpha\mu} p_{\alpha\nu} - q_{\alpha\nu} p_{\alpha\mu}). \quad (3.13)$$

At the quantum level the algebra of the operators  $M_{\mu\nu}$  should be determined by the commutators of the operators  $q_\alpha$  and  $p_\alpha$ ,  $\alpha = 1, 2, 3$ . In their turn these commutators are defined, as mentioned above, by the corresponding Dirac brackets. But the requirement of the Poincare invariance of the theory under consideration determines the algebra of the operators  $p_{1\mu}$  and  $M_{\mu\nu}$  completely. This algebra must be the same as the algebra of the Poincare group. Without calculating the corresponding Dirac brackets we assume that the Poincare invariance takes place. As the scalar Casimir operators of the Poincare group we take the following ones [16]

$$p_1^2 = p_1^\mu p_{1\mu}, \quad (3.14)$$

$$W = \frac{1}{2} M_{\mu\nu} M^{\mu\nu} p_1^2 - (M_{\mu\sigma} p_1^\sigma)^2. \quad (3.15)$$

In the four-dimensional space-time the invariant  $W$  is the squared Pauli-Lubanski vector with sing minus,  $W = -w_\mu w^\mu$ , where  $w_\mu = (1/2)\epsilon_{\mu\nu\rho\sigma} M^{\nu\rho} p_1^\sigma$ . The obvious advantage of the definition (3.15) is the possibility to use it for arbitrary  $D$ .

The relativistic invariance requires that the physical state vectors  $|\psi\rangle$  should be the eigenvectors of the operators (3.14) and (3.15)

$$p_1^2 |\psi\rangle = M^2 |\psi\rangle. \quad (3.16)$$

As mentioned above, we suppose that  $M^2 > 0$ . In this case we can go to the rest frame where  $p_1^\mu = (p_1^0 = M, \mathbf{p}_1 = 0)$ . Here we have

$$W = (p_1^0)^2 M_{12} M^{12} = \frac{M^2}{2} C_2(SO(2)), \quad (3.17)$$

<sup>1</sup>In the ordinary theory of the relativistic particle with the action  $S = -m \int dt \sqrt{|\dot{\mathbf{x}}|^2}$  the squared mass of the particle  $p^2$  is positive only for the initial data obeying the inequality  $\dot{\mathbf{x}}^2 > 0$ . It is important that this condition, satisfied at the initial moment will be fulfilled always. In the general case for arbitrary sing of  $\dot{\mathbf{x}}^2$  we have

$$p_\mu = -\frac{\partial L}{\partial \dot{x}^\mu} = m \frac{\dot{x}_\mu}{\sqrt{|\dot{\mathbf{x}}|^2}} \text{sign}(\dot{\mathbf{x}}^2).$$

Squaring of this equation gives

$$p^2 = m^2 \text{sign}(\dot{\mathbf{x}}^2).$$

where  $C_2(SO(2))$  is the squared Casimir operator of the  $SO(2)$  group (see for example [18]). As known [17] the group of rotations on the plane  $SO(2)$  has three different representations with integer, half-odd-integer and continuous values for spin  $j$ . In all these cases we have [18]

$$C_2(SO(2)) = 2j^2. \quad (3.18)$$

It should be noted here that in the model under consideration we have to deal with the tensor representations of the  $SO(2)$  group but not with spinor ones because the initial action (2.1) contains no spin degrees of freedom. Thus the eigenvalues of the Casimir operator  $W$  are

$$W = M^2 j^2, \quad M^2 > 0, \quad j \geq 0. \quad (3.19)$$

We assume, in addition to (3.16), that the state vector  $|\psi\rangle$  is the eigenvector of the Casimir operator  $W$

$$W |\psi\rangle = M^2 j^2 |\psi\rangle, \quad M^2 > 0, \quad j \geq 0. \quad (3.20)$$

Let us proceed to the consideration of the wave equation. First we take as the second-class constraints the set (2.33). In addition to the proper time gauge (3.3) we introduce one more gauge condition that transforms the constraints  $\phi_3^{(1)}$  into the second-class constraint. As a result, only one wave equation survives in (3.2)

$$\phi_1^{(1)} |\psi\rangle = (p_3^2 q_3^2 - \beta^2) |\psi\rangle = 0. \quad (3.21)$$

If  $D = 3$ , one can express the left-hand side of (3.21) in terms of the Casimir operators  $p_1^2$  and  $W$ . For this purpose we have to calculate the invariant  $W$  on the surface in the phase space determined by all the constraints and gauge conditions except for  $\phi_1^{(1)}$ . After a rather cumbersome calculation (see Appendix ) we obtain

$$1 - x = \frac{W - (\alpha \sqrt{m^2 - p_1^2} + |\beta| m \sqrt{x})^2}{\beta^2 p_1^2} \frac{x}{1+x}, \quad (3.22)$$

where  $x = \beta^2 / (p_3^2 q_3^2)$ . Thus the condition (3.21)

$$(1 - x) |\psi\rangle = 0 \quad (3.23)$$

is equivalent to

$$[W - (\alpha \sqrt{m^2 - p_1^2} + |\beta| m)^2] |\psi\rangle = 0. \quad (3.24)$$

From (3.16), (3.20) and (3.24) we obtain immediately the mass spectrum

$$M^2 j^2 = (\alpha \sqrt{m^2 - M^2} + |\beta| m)^2. \quad (3.25)$$

the value of this operation  $M'$  should be less than  $m^2$  in accordance with (3.12)

If  $|\alpha| \leq |\beta|$ , then eq (3.25) can be solved with respect to the ratio  $M/m$

$$\frac{M}{m} = \frac{|\beta| |j|}{\alpha^2 + j^2} \left( 1 + \sqrt{1 + \frac{\alpha^2 + j^2}{\beta^2 j^2} (\alpha^2 - \beta^2)} \right). \quad (3.26)$$

When

$$-|\beta| \leq \alpha \leq |\beta| \quad (3.27)$$

the ratio  $M/m$  takes real values only if the spin of the state  $j$  satisfies the condition

$$1 - \frac{\alpha^2}{\beta^2} < \frac{j^2}{\alpha^2 + j^2}. \quad (3.28)$$

It is interesting to note that under the conditions (3.27) and (3.28) the ratio  $M/m$  turns out to be a double-valued function of the model parameters  $\alpha$ ,  $\beta$  and the spin  $j$  of the state

$$\frac{M}{m} = \frac{|\beta| |j|}{\alpha^2 + j^2} \left( 1 \pm \sqrt{1 - \left( \frac{\alpha^2}{j^2} + 1 \right) \left( \frac{\alpha^2}{\beta^2} + 1 \right)} \right). \quad (3.29)$$

The same function takes place in some infinite component wave equations [18]. For  $|\alpha| = |\beta|$  it is not success to resolve eq. (3.18) with respect to  $M/m$ . Putting in (3.18) in turn  $\alpha = 0$  and  $j^2 = 0$  we obtain the mass spectra in the theory of the relativistic particle with torsion ( $\alpha = 0$ ) or with curvature ( $\beta = 0$ ) derived in [1].

Now we proceed to discuss the realization of the Lie algebra  $\mathfrak{p}_2^2$  and  $W$  as the differential operator. More easily it can be done in a rest frame, i.e. by virtue of (3.17) we have

$$W = M^2 M_{12} M^{12}, \quad (3.30)$$

and  $p_1^2$  refers to the multiplication by  $M^2$ . The operator  $M_{12}$  which describes the rotations on the plane can be taken in the form

$$M_{12} = -i \frac{\partial}{\partial \varphi} + c, \quad (3.31)$$

where  $\varphi$  is some angular variable and  $c$  is a constant to be determined below. As the wave function we shall use  $2\pi$ -periodic functions

$$\psi(\varphi) = \sum_{l \in \mathbb{Z}} e^{il\varphi} a_l. \quad (3.32)$$

Substituting of (3.29)–(3.31) into (3.20) gives

$$j^2 = (l + c)^2, \quad l \in \mathbb{Z}. \quad (3.33)$$

Without loss of generality we can regard  $l$  as an integer part of  $j$  and  $c$  as its fractional part. Thus we are dealing with the usual  $2\pi$ -periodic wave functions and nevertheless we can describe integral, half-odd-integral and continuous values of the spin in the model under consideration.

## 4 Interaction with external Abelian gauge field

As shown in the preceding Section, in the model with the action (2.1) there appears a nonvanishing intrinsic angular momentum, i.e. spin. It is worthwhile to investigate the dynamics of the spin variables in the framework of this model. However, when we are dealing with the free action (2.1), this dynamics turns out to be trivial: the spin squared ( $\sim W$ ) and its components ( $\sim M_{\mu\nu}$ ) are conserved.

Here we consider the introduction of the interaction with an external Abelian gauge field in the model (2.1). From the geometrical point of view it can be done in the following way

$$L_{\text{int}} = - \sum_{a=0}^{D-1} g_a n_a^\mu A_\mu(x) - \sum_{a \neq b} g_{ab} n_a^\mu n_b^\nu F_{\mu\nu}(x), \quad (4.1)$$

where  $n_a^\mu$ ,  $a = 0, 1, \dots, D-1$  are the unit vectors forming the moving basis on the world trajectory,  $A_\mu(x)$  is the vector potential of the external electromagnetic field and  $F_{\mu\nu}$  is its strength tensor,  $g_a$  and  $g_{ab}$  are the interaction constants. This Lagrangian obviously retains the reparametrization invariance of the whole action. In order to remove the superlight velocities we have to impose the following conditions

$$n_0^2 = \left( \frac{dx^\mu}{ds} \right)^2 = 1, \quad n_i^2 = -1, \quad i = 1, 2, \dots, \quad ds^2 = dx^\mu dx_\mu. \quad (4.2)$$

The basis vectors  $n_a^\mu$ ,  $a = 0, 1, \dots, D-1$  can be represented in terms of the derivatives of the radius-vector  $x^\mu$  [20]

$$\begin{aligned} \frac{dx^\mu}{ds} &= n_0^\mu, \\ \frac{d^2 x^\mu}{ds^2} &= k_1 n_1^\mu, \\ \frac{d^3 x^\mu}{ds^3} &= k_1 k_2 n_2^\mu - k_1^2 n_0^\mu + \frac{dk_1}{ds} n_1^\mu, \\ &\dots \dots \dots \\ \frac{d^D x^\mu}{ds^D} &= k_1 k_2 \dots k_{D-1} n_{D-1}^\mu + \dots, \end{aligned} \quad (4.3)$$

where  $k_1(s)$ ,  $k_2(s)$ ,  $\dots$ ,  $k_{D-1}(s)$  are the curvatures of the world trajectory. If  $D=3$ , then  $k_1$  is called the curvature and  $k_2$  is called the torsion.

The free action (2.1) can be generalized in the  $D$ -dimensional space-time by the formula

$$S_0 = -m \int ds - \sum_{i=1}^{D-1} \alpha_i \int k_i(s) ds. \quad (4.4)$$

If we restrict for simplicity the summation over  $i$  in eq.(4.4), then the analogous restriction should be made in (4.1) so that the coordinate derivatives of the same order enter into (4.1) and (4.4). The canonical quantization of the simplest model of this kind is accomplished in [21].

## 5 Conclusion

The results obtained in this paper show that the point like Lagrangians with higher derivatives of the form (2.1) describe in the general case the point particles with non-vanishing spin. This conclusion relies on the fact that the Casimir operator of the Poincaré group  $W$  proportional to spin squared does not vanish on the physical submanifold of the phase space defined by the constraint equations and the gauge conditions. For  $D = 3$  we have obtained the exact expression for the Regge-trajectory in the sector of the theory without tachyonic states. When the spin of the state increases, its mass decreases. There is an upper bound on the squared mass of the state. If  $D = 3$ , we have here the possibility to describe the states with integer, half-odd-integer or continuous values of the spin. It is important to note that in the framework of this model we dealing with the coordinate representation for the wave function even in the case of the half-odd-integer spin instead of the spinor one. And there remains an open question whether one can here obtain the Dirac equation for such spin values. Further investigations are required also in order to elucidate the role of the tachyonic states in the model.

## Appendix

Here we present some details of the calculation of the invariant  $W$  on the submanifold of the phase space defined by the constraint equations and gauge conditions except for  $\phi_1 = 0$ . At first we take into account eqs. (2.9), (2.10), (2.16), (2.17), (2.22), (3.3) as well as the condition  $p_1 p_2 = 0$  valid in the three dimensional space time. As a result, one obtains

$$W = p_1^2 (q_1^2 p_2^2 + q_3^2 p_3^2) - p_2^2 (p_1 q_2)^2 + 2 (p_1 p_3) (p_1 q_2) (p_2 q_3) - q_3^2 (p_1 p_3)^2 \quad (\text{A.1})$$

Now we use the rest of the constrains ( see also eqs. (3.8) and (3.9) ) from which it follows that

$$(p_1 q_2)^2 = m^2 q_2^2, \quad p_2 q_3 = \alpha \frac{q_3^2}{\sqrt{q_2^2 q_3^2}}, \quad p_1 p_3 = \frac{\beta^2}{12}. \quad (\text{A.2})$$

It is convenient to introduce the notation

$$\frac{\beta^2}{p_3^2 q_3^2} = x. \quad (\text{A.3})$$

By making use of (A.2) and (A.3) we transform eq (A.1) in the form

$$W - \left( \alpha \sqrt{m^2 - p_1^2} + |\beta| m \sqrt{x} \right)^2 = \beta^2 p_1^2 \frac{(1-x)(1+x)}{x}. \quad (\text{A.4})$$

## References

- [1] V. V. Nesterenko, *J. Phys. A* **22** (1989), 1673.
- [2] V. V. Nesterenko, *Teor. Mat. Fiz.* (1991); *JINR E2-90-268 preprint* (1990).
- [3] M. S. Plyuscha, *Mod. Phys. Lett. 3A* (1988), 1299; *4A* (1989), 337; *Int. J. Mod. Phys. 4A* (1989), 3851.
- [4] C. Battle, J. Gomez, J. M. Pons and N. Roman-Roy, *J. Phys. A* **21** (1988), 2523.
- [5] R. Pisarski, *Phys. Rev. D* **34** (1986), 670.
- [6] H. Arodz, A. Sitarz and P. Wegrzyu, *Acta Phys. Pol. B* **20** (1985), 921.
- [7] A. M. Polyakov, *Mod. Phys. Let. 3A* (1988), 825; A. L. Kholodenko, *Ann. Phys.* **202** (1990), 186.
- [8] N. Imai, K. Ishikawa and I. Tanaka, *EPHOU 89-003 preprint* (1989).
- [9] A. Alekseev and S. Shatashvili, *Mod. Phys. Let. A* **3** (1988) 1551.
- [10] M. Wagner, J. M. P. Gunn and A. B. Baumgartner, *J. Phys. A* **18** (1985), 2007.
- [11] V. V. Nesterenko and Nguyen Suat Han, *Int. J. Mod. Phys. A* **7** (1992) 2317.
- [12] G. V. Dunne, R. Jackiw and C. A. Trugenberger, *Ann. Phys.* **194** (1989), 137; *Phys. Rev. D* **41** (1990), 661.
- [13] M. V. Ostrogradski, *Mem. Ac. St. Petersburg VI*, **4** (1850), 293. See also: B. L. Whittaker, *Analytical Dynamics of Particles and Rigid Bodies*, p.265, Cambridge University Press, 1988.
- [14] P. A. M. Dirac, *Lectures on Quantum Mechanics*, Yale University, New York, 1964.

- [15] V. V. Nesterenko and A. M. Chervjakov, *JINR P2-86-323 preprint (1986)*.
- [16] S. S. Schweber, *An Introduction to Relativistic Quantum Field Theory*, Row Peterson, New York, 1961.
- [17] H. Weyl, *The Theory of Groups and Quantum Mechanics*, Dover Publications, 1931.
- [18] A. O. Barut and R. Raczka, *Theory of Group Representations and Applications*, Polish Scientific Publishers, Warszawa, 1977.
- [19] M. Pavsic, *Phys. Lett. B221 (1989) 264*.
- [20] Yu. A. Aminov, *Differential Geometry and Topology of Curves (in Russian)*, Moscow, Nauka, 1987.
- [21] V. V. Nesterenko, *Int. J. Mod. Phys. (1991)*.

Received by Publishing Department  
on January 15, 1991.



WILL YOU FILL BLANK SPACES IN YOUR LIBRARY?

You can receive by post the books listed below. Price: — in US \$, including the packing and registered postage.

D13-85-733	Proceedings of the XII International Symposium on Nuclear Electronics, Dubna, 1985.	14.00
D1,2-85-666	Proceedings of the VIII International Seminar on High Energy Physics Problems, Dubna, 1986 (2 volumes)	25.00
D3,4,17-86-747	Proceedings of the V International School on Neutron Physics, Alushta, 1986.	25.00
D9-87-105	Proceedings of the X All-Union Conference on Charged Particle Accelerators, Dubna, 1986 (2 volumes)	25.00
E7-87-63	Proceedings of the International School-Seminar on Heavy Ion Physics, Dubna, 1986.	25.00
D2-87-123	Proceedings of the Conference "Renormalization Group 86", Dubna, 1986.	12.00
D2-87-798	Proceedings of the VIII International Conference on the Problems of Quantum Field Theory, Alushta, 1987.	10.00
D14-87-799	Proceedings of the International Symposium on Muon and Pion Interactions with Matter, Dubna, 1987.	13.00
D17-88-95	Proceedings of the IV International Symposium on Selected Topics in Statistical Mechanics, Dubna, 1987.	14.00
E1,3-88-426	Proceedings of the 1987 JINR-CERN School of Physics, Varna, Bulgaria, 1987.	14.00
D11-88-833	Proceedings of the International Workshop on Modern Trends in Activation Analysis in JINR, Dubna, 1988	8.00
D13-88-938	Proceedings of the XIII International Symposium on Nuclear Electronics, Varna, 1988	13.00
D17-88-681	Proceedings of the International Meeting "Mechanisms of High-T <sub>c</sub> Superconductivity", Dubna, 1988.	20.00
D9-89-52	Proceedings of the XI All-Union Conference on Charged Particle Accelerators, Dubna, 1988 (2 volumes)	30.00
E2-89-525	Proceedings of the Seminar "Physics of e <sup>+</sup> e <sup>-</sup> -Interactions" Dubna, 1988.	10.00
D5-89-501	Proceedings of the International School-Seminar on Heavy Ion Physics, Dubna, 1989.	19.00
D13-90-457	Proceedings of the Workshop on DNA Repair or Mutagenesis Induced by Radiation, Dubna, 1990.	15.00

Orders for the above-mentioned books can be sent at the address:  
 Publishing Department, JINR  
 Mail Post Office, P.O. Box 79 101000 Moscow, USSR

# SUBJECT CATEGORIES OF THE JINR PUBLICATIONS

Index	Subject
1.	High energy experimental physics
2.	High energy theoretical physics
3.	Low energy experimental physics
4.	Low energy theoretical physics
5.	Mathematics
6.	Nuclear spectroscopy and radiochemistry
7.	Heavy ion physics
8.	Cryogenics
9.	Accelerators
10.	Automatization of data processing
11.	Computing mathematics and technique
12.	Chemistry
13.	Experimental techniques and methods
14.	Solid state physics. Liquids
15.	Experimental physics of nuclear reactions at low energies
16.	Health physics. Shieldings
17.	Theory of condensed matter
18.	Applied researches
19.	Biophysics

Нестеренко В.В.

E2-91-19

Каноническое квантование релятивистской частицы  
с кривизной и кручением

Рассматривается обобщенное действие релятивистской точечной частицы, которое содержит помимо длины мировой траектории два дополнительных слагаемых: интегралы вдоль траектории от ее кривизны и кручения. Построен гамильтонов формализм для этой модели в  $D$ -мерном пространстве времени. Найден полный набор связей, выполнено их разделение на связи первого и второго рода. Для  $D = 3$  получен массовый спектр, задаваемый уравнением, в которое входят масса, спин состояния и параметры модели. Показана возможность описания в данной модели состояний с целым, полуцелым и непрерывным спином ( $D = 3$ ). В операторной форме построено волновое уравнение и найден пропегатор частицы. Геометрическим путем вводится взаимодействие с внешним абелевым калибровочным полем.

Работа выполнена в Лаборатории теоретической физики ОИЯИ.

Сообщение Объединенного института ядерных исследований. Дубна 1991

Nesterenko V.V.

E2-91-19

Canonical Quantization of a Relativistic Particle  
with Curvature and Torsion

A generalization of the relativistic particle action is considered. It contains, in addition to the length of the world trajectory, the integrals along the world curve of its curvature and torsion. The generalized Hamiltonian formalism for this model in the  $D$ -dimensional space-time is constructed. A complete set of the constraints in the phase space is obtained and their division into the first-class and the second-class constraints is accomplished. On this basis the canonical quantization of the model is fulfilled. For  $D = 3$  the mass spectrum is obtained in the sector without tachyonic states, the mass of the state being dependent on its spin. It is shown that in the framework of this model when  $D = 3$  we have the possibility to describe the states with integral, half-odd-integral and continuous spins. Interaction with an external Abelian gauge field introduced in the geometrical way.

The investigation has been performed at the Laboratory of Theoretical Physics, JINR.

Communication of the Joint Institute for Nuclear Research, Dubna 1991

20 коп.

Корректурa и макет Т.Е.Попеко.

Подписано в печать 23.01.91.

Формат 60x90/16. Офсетная печать. Уч.-изд. листов 1,35.

Тираж 490. Заказ 44075.

Издательский отдел Объединенного института ядерных исследований.  
Дубна Московской области.