



**INTERNATIONAL CENTRE FOR
THEORETICAL PHYSICS**

**NORMAL FORMS OF INVARIANT VECTOR FIELDS
UNDER A FINITE GROUP ACTION**

Federico Sánchez-Bringas



**INTERNATIONAL
ATOMIC ENERGY
AGENCY**



**UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION**

MIRAMARE-TRIESTE



International Atomic Energy Agency
and
United Nations Educational Scientific and Cultural Organization
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**NORMAL FORMS OF INVARIANT VECTOR FIELDS
UNDER A FINITE GROUP ACTION**

Federico Sánchez-Bringas *

International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

Let Γ be a finite subgroup of $GL(n, \mathbb{C})$. This subgroup acts on the space of germs of holomorphic vector fields vanishing at the origin in \mathbb{C}^n . We prove a theorem of invariant conjugation to a normal form and linearization for the subspace of invariant elements and we give a description of these normal forms in dimension $n = 2$.

MIRAMARE – TRIESTE

July 1992

* Permanent address: Instituto de Matemáticas, Universidad Nacional Autónoma de México, Ciudad Universitaria, Mexico 04510, D.F., Mexico.

Introduction

The goal of this paper is to show that the classic theorems of Poincaré-Dulac [DU] and Siegel [SI] of conjugation to a normal form and linearization of germs of holomorphic vector fields at $0 \in \mathbb{C}^n$ hold for the quotient space \mathbb{C}^n/Γ , where Γ is a finite subgroup of $GL(n, \mathbb{C})$. In this situation we consider the germs of holomorphic vector fields and the germs of conjugating diffeomorphism of \mathbb{C}^n invariant by the action of the subgroup.

It is well known that \mathbb{C}^n/Γ has the structure of an algebraic variety and furthermore any variety which is the quotient of a finite group of local diffeomorphisms of \mathbb{C}^n is of this form (in a specific system of coordinates) [CA], then we obtain here results for conjugations to normal forms and linearizations of germs of holomorphic vector fields in this kind of algebraic varieties.

In a different context, like bifurcation theory, sometimes conjugation to a normal form of germs of holomorphic vector fields which preserves symmetries are needed, this results can also be applied.

In the first section we prove the main theorem using the algebraic approach developed in [CH]. In the second section we analyse carefully the case \mathbb{C}^2/Γ and we give a description of normal forms.

Finally we wish to thank Xavier Gómez-Mont for his helpful comments and remarks concerning this work.

§1 Invariant conjugation to a normal form and linearization in \mathbb{C}^n

Let $\mathcal{X}(\mathbb{C}^n, 0)$ be the space of germs of holomorphic vector fields at $0 \in \mathbb{C}^n$ vanishing at the origin. Let Γ be a finite subgroup of $GL(n, \mathbb{C})$ which acts naturally on $\mathcal{X}(\mathbb{C}^n, 0)$, we say that X is invariant if it is invariant by this action, namely if for all $\gamma \in \Gamma$, $d\gamma^{-1}(X(\gamma(z))) = X(z)$. Let $\mathcal{X}(\mathbb{C}^n/\Gamma, 0)$ be the subspace of invariant elements of $\mathcal{X}(\mathbb{C}^n, 0)$.

Given $X \in \mathcal{X}(\mathbb{C}^n, 0)$ denote by X_1 its linear part, $dX(0)$ and suppose it belongs to $GL(n, \mathbb{C})$. Let S be the semisimple part of X_1 , we say X is a normal form if $L_S X = 0$, where L_S is the Lie derivative of S .

X_1 is said to be resonant if its eigenvalues satisfy a relation (resonance) like this:

$$\lambda_u = \sum_1^n m_i \lambda_i, \quad u = 1, \dots, n \quad (m_1, \dots, m_n) \in \mathbb{N}^n, \quad \sum_1^n m_i \geq 2$$

The vector field $z_1^{m_1} \dots z_n^{m_n} e_u$, $u = 1, \dots, n$ is the monomial vector field associated to this resonance.

Suppose that $X \in \mathcal{X}(\mathbb{C}^n, 0)$ is a normal form and the coordinates of \mathbb{C}^n are given by a basis of eigenvectors of the semisimple part S of X_1 . Then condition $L_S X = 0$ implies that $X - X_1$ is a sum of resonant monomial vector fields.

Let $Diff(\mathbb{C}^n, 0)$ be the group of germs of holomorphic diffeomorphisms which fix the origin of \mathbb{C}^n , Γ acts by conjugation here. We say that $\phi \in Diff(\mathbb{C}^n, 0)$ is invariant if it is invariant by this action, namely if for all $\gamma \in \Gamma, \gamma^{-1}\phi\gamma = \phi$. Denote by $Diff(\mathbb{C}^n/\Gamma, 0)$ the group of invariant elements of $Diff(\mathbb{C}^n, 0)$.

For any $X, Y \in \mathcal{X}(\mathbb{C}^n, 0)$ we say X is conjugated to Y if there is a $\phi \in Diff(\mathbb{C}^n, 0)$ such that $\phi^*X = Y$, where $\phi^*X = d\phi^{-1}X\phi$. When Y is the linear part of X we say X is linearizable. If X and Y are invariant, the conjugation (linearization) is called invariant.

Theorem 1 Let Γ be a finite subgroup of $GL(n, \mathbb{C})$ and $X \in \mathcal{X}(\mathbb{C}^n/\Gamma, 0)$. Suppose the linear part X_1 of X is invertible. Then:

1.1 X is invariantly conjugated, possibly formally to a normal form. If X_1 is non-resonant then this conjugation is an invariant linearization.

1.2 If X is holomorphically conjugated to a normal form, then it can be conjugated in an invariant holomorphic way.

The proof of this theorem is a consequence of the following lemma. Let $\mathcal{O}_{\mathbb{C}^n, 0}$ be the algebra of germs of holomorphic functions at $0 \in \mathbb{C}^n$ and $m = \{f \in \mathcal{O}_{\mathbb{C}^n, 0}; f(0) = 0\}$ its maximal ideal. For each non-negative integer k denote $J_{\mathbb{C}^n, 0}^k = \mathcal{O}_{\mathbb{C}^n, 0}/m^k$ the algebra of finite dimension of k -jets of holomorphic functions. The element $X \in \mathcal{X}(\mathbb{C}^n, 0)$ defines a derivation X^* of $\mathcal{O}_{\mathbb{C}^n, 0}$, $X^*f = L_X f$ and in a natural way the k -jet of X determines a derivation X_k^* of $J_{\mathbb{C}^n, 0}^k$, then X_k^* has a canonical decomposition: $X_k^* = S_k^* + N_k^*$, where S_k^* is the semisimple part and N_k^* is the nilpotent part. A remarkable fact proved in [CH] is that S_k^* is a derivation.

In a similar way, we denote by $Diff_k(\mathbb{C}^n, 0)$ the group of k -jets of germs of holomorphic diffeomorphisms of \mathbb{C}^n . The definition of conjugation to a normal form (linearization) is extended in a natural way to the space of k -jets of germs of vector fields, $\mathcal{X}^k(\mathbb{C}^n, 0)$.

Lemma 2 Let Γ be a finite subgroup of $GL(n, \mathbb{C})$, k a non-negative integer and $X_k \in \mathcal{X}^k(\mathbb{C}^n/\Gamma, 0)$ then

2.1 The semisimple part S_k^* of X_k^* is invariant.

2.2 S_k^* is invariant linearizable if X_1 is invertible.

Proof. 2.1 On one hand we have the following fact [HU]:

Let V be a \mathbb{C} -vector space of finite dimension and T an endomorphism of V . Then the semisimple part of T has a polynomial expression in T , $p(T)$ with coefficients in \mathbb{C} . On the other hand, as X is invariant and $\gamma \in \Gamma$ is linear we have $d\gamma^{-1}X\gamma = \gamma^{-1}X\gamma = X$ then for any non-negative integer k , $\gamma^{-1}X_k^*\gamma = X_k^*$ and $\gamma^{-1}X_k^* \circ \dots \circ X_k^*\gamma =$

$X_k^* \circ \dots \circ X_k^*$ then any polynomial expression in X_k^* with coefficients in \mathbb{C} is invariant.

2.2 Let $\phi_k \in \text{Diff}_k(\mathbb{C}^n, 0)$ be the Poincaré-Dulac diffeomorphism which exists because X_1 is invertible. ϕ_k is tangent (κ -order) to the identity diffeomorphism, and linearizes the semisimple part of X_k^* . Define the average $\tilde{\phi}_k = |\Gamma|^{-1} \sum_{\gamma \in \Gamma} \gamma^{-1} \phi_k \gamma$.

$\tilde{\phi}_k$ is invariant and tangent (κ -order) to the identity diffeomorphism. Besides,

$$\begin{aligned} (|\Gamma|^{-1} \sum_{\gamma \in \Gamma} \gamma^{-1} \phi_k \gamma)^*(S_k^*) &= |\Gamma|^{-1} \sum_{\gamma \in \Gamma} (\gamma^{-1} \phi_k^{-1} \gamma) \cdot S_k^*(\gamma^{-1} \phi_k \gamma) = \\ &= |\Gamma|^{-1} \sum_{\gamma \in \Gamma} \gamma S_1^* \gamma^{-1} = S_1^* \end{aligned}$$

where S_1^* is the linear part of S_k^* which is invariant because of S_k^* .

Proof of the theorem. 1.1 $L_{S_1} X_k = 0$ if and only if $0 = (L_{S_1} X_k)^* = S_1^* X_k^* - X_k^* S_1^*$, then the canonical decomposition of X_k^* implies that conjugating to a normal form in $\mathcal{X}^*(\mathbb{C}^n, 0)$ is equivalent to linearizing the semisimple part S_k^* . Now let $\tilde{\phi}_k$ be like in lemma 2. Since composition $\tilde{\phi}_1 \circ \dots \circ \tilde{\phi}_2$ has the same l -jet than $\tilde{\phi}_{l+1} \circ \dots \circ \tilde{\phi}_2$ then the limit $\text{Lim}_{k \rightarrow \infty} (\tilde{\phi}_k \circ \dots \circ \tilde{\phi}_2)$ defines a diffeomorphism $\tilde{\phi}$, possible formal which conjugates invariantly X_k^* to a normal form.

1.2 Let ϕ be the holomorphic conjugation (in any system of coordinates) then a similar argument as in 1 implies that $|\Gamma|^{-1} \sum_{\gamma \in \Gamma} \gamma^{-1} \phi \gamma \in \text{Diff}(\mathbb{C}^n, 0)$ conjugates X holomorphically and invariantly to the respective normal form.

§2 Description in \mathbb{C}^2 : Invariant normal forms

Let X_1 be a linear vector field in \mathbb{C}^2 with eigenvalues λ_1, λ_2 . Choose a base of \mathbb{C}^2 , $\{e_1, e_2\}$ of eigenvectors of S , the semisimple part of X_1 . We say that X_1 belongs to the Poincaré domain if 0 is not in the segment $[\lambda_1, \lambda_2]$. Otherwise we say X_1 belongs to the Siegel domain. The eigenvalues λ_1, λ_2 are of type (C, ν) , $C, \nu \geq 0$ if: $|\lambda_u - m_1 \lambda_1 - m_2 \lambda_2| \geq C(|m_1| + |m_2|)^{-\nu}$ for all $(m_1, m_2) \in (\mathbb{N})^*$, $u = 1, 2$.

Before applying theorem 1 in this context we point out that conditions in order for X to be conjugated holomorphically to a normal form have been established in [DU] if X_1 is in the Poincaré domain and in [SI] if X_1 belongs to the Siegel domain.

Theorem 3 Let X_1 be the linear part of $X \in \mathcal{X}(\mathbb{C}^n/\Gamma, 0)$. Let λ_1, λ_2 be the eigenvalues of X_1 .

3.1 If X_1 is not resonant, then X is linearizable in a holomorphic invariant way in the following cases:

i) X_1 belongs to the Poincaré domain.

ii) X_1 belongs to the Siegel domain and λ_1, λ_2 are of type (C, ν) for some $C, \nu \geq 0$.

3.2 If X_1 is resonant, then X is conjugated in a holomorphic invariant way to a normal form in the following cases:

i) X_1 belongs to the Poincaré domain.

ii) X_1 belongs to the Siegel domain and the normal form is colinear to X_1 .

Remark There exist cases where the invariant conjugation to a normal form is only formal. For example if Γ is a diagonal group (i.e. each of its elements are diagonal) we are going to show there are normal forms with linear part in the the Siegel domain which do not verify condition 2; ii). In this case the conjugating diffeomorphism ϕ may be divergent because one of its coordinate functions can have coefficients which grow like the Euler function, [BR]:

$$\eta(z) = \sum_{\delta=1}^{\infty} (\delta - 1)! z^{\delta}$$

We express the invariance condition in $\mathcal{X}(\mathbb{C}^2, 0)$ with an average morphism of the group action. Let $\Pi : \mathcal{X}(\mathbb{C}^2, 0) \rightarrow \mathcal{X}(\mathbb{C}^2/\Gamma, 0)$ be the morphism of \mathbb{C} -vectorial spaces defined by $\Pi(X) = |\Gamma|^{-1} \sum_{\gamma \in \Gamma} \gamma^* X$. Then X is invariant if and only if $\Pi(X) = X$.

There are two different cases for the family of finite subgroups of $GL(2, \mathbb{C})$:

i) If Γ is diagonal, the monomial vector fields are eigenvectors of Π and $\Pi(X) = 0$ if X is not invariant.

ii) If Γ is not diagonalizable the eigenvectors of Π are not monomials and Π does not vanish monomials.

Let us regard first the case of diagonalizable groups.

Proposition 4 If Γ is not diagonalizable and X_1 is a linear vector field, then $\Pi(X_1) = X_1$ if and only if $X_1 = \lambda(z_1, z_2)$.

Proof. Suppose X_1 is given in its Jordan canonical form. If X_1 has different eigenvalues the condition $X_1 \gamma = \gamma X_1$ implies γ is diagonal, therefore Γ must be diagonal.

If

$$X_1 = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix}$$

$X_1 \gamma = \gamma X_1$ implies

$$\gamma = \begin{pmatrix} \mu & b \\ 0 & \mu \end{pmatrix}$$

but Γ is finite then $\gamma^n = Id$ and b must be 0.

Remark This proposition implies that for non diagonalizable groups our theorem is a linearizing theorem illustrated by the following example: Let Γ be the binary dihedral group generated by $\gamma = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ and $j = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

The vector field $X(z_1, z_2) = (z_1, z_2) + (z_1^5, z_2^5)$ is not a multiple of the radial vector field which belongs to $\mathcal{X}(\mathbb{C}^2/\Gamma, 0)$.

Suppose now the group is diagonal. In order to simplify the description of invariant normal forms we will suppose Γ is cyclic and generated by

$$\begin{pmatrix} e^{(2\pi l_1 \sqrt{-1})/n_1} & 0 \\ 0 & e^{(2\pi l_2 \sqrt{-1})/n_2} \end{pmatrix}$$

$n_u \in \mathbf{N}, l_u \in \mathbf{Z}, (l_u, n_u) = 1, u = 1, 2.$

If $X(z_1, z_2) = \sum_{u=1,2} (\sum_{i,j=0}^{\infty} c_{ij}^u z_1^i z_2^j) e_u$, the equivariance condition is imposed independently for monomial vector fields. The next proposition describes invariant vector fields.

Proposition 5 Let n be the least common multiple of n_1, n_2 and $\eta_u = n l_u / n_u$. The monomial vector field $z_u (z_u^i z_v^j) e_u, u = 1, 2, u \neq v, i \geq -1, j \geq 0, i+j \geq 0$ is invariant if and only if $\eta_u i = -\eta_v j \pmod{n}$.

Proof. Suppose

$$\gamma = \begin{pmatrix} \gamma_1 & 0 \\ 0 & \gamma_2 \end{pmatrix}$$

belongs to Γ then $\Pi(z_1^i z_2^j e_u) = (1/n \sum_{\gamma \in \Gamma} \gamma_u^{-1} \gamma_1^i \gamma_2^j) z_1^i z_2^j e_u$, therefore $z_1^i z_2^j e_u$ is invariant if and only if $n^{-1} \sum_{\gamma \in \Gamma} \gamma_u^{-1} \gamma_1^i \gamma_2^j = 1$. If $u = 1, \sum_{\gamma \in \Gamma} \gamma_1^{i-1} \gamma_2^j = \sum_{k=1}^n e^{2\pi \sqrt{-1} k ((l_1/n_1)(i-1) + (l_2/n_2)j)}$. This sum is n if $(n/n_1)l_1 i + (n/n_2)l_2 j \equiv (n/n_1)l_1 \pmod{n}$ and vanishes otherwise. Similarly for $u = 2$.

When X_1 belongs to the Poincaré domain and its eigenvalues are resonant, then the origin, λ_1 and λ_2 are colinear, besides $0 \notin [\lambda_1, \lambda_2]$ therefore there is only one possible type of resonance: $\lambda_u = m_v \lambda_v, u \neq v$.

When X_1 belongs to the Siegel domain the resonance $\lambda_u = m_u \lambda_u + m_v \lambda_v$ generates an infinite family of resonances of type:

$$\lambda_u = k((m_u - 1)\lambda_u + m_v \lambda_v) + \lambda_u \lambda_v = k((m_u - 1)\lambda_u + m_v \lambda_v) + \lambda_v, k \in \mathbf{N}, u \neq v.$$

Finally let us make the following classification:

Proposition 6 Let Γ be a finite diagonal cyclic subgroup of $GL(2, \mathbf{C})$. The invariant normal forms are:

1 If X_1 belongs to the Poincaré domain.

$$X(z_1, z_2) = X_1(z_1, z_2) + a_i z_v^i e_u$$

where $\eta_v i \equiv \eta_u \pmod{n}, u \neq v$ and $u, v = 1, 2$

2 If X_1 belongs to the Siegel domain.

$$X(z_1, z_2) = X_1(z_1, z_2) + (z_1 (\sum_{k=1}^{\infty} a_k z_1^{ki} a z_2^{kj}), z_2 (\sum_{k=1}^{\infty} b_k z_1^{ki} z_2^{kj}))$$

where $\eta_1 i \equiv -\eta_2 j \pmod{n}$ or $k \equiv 0 \pmod{n}$ and i, j are like in proposition 5.

For all cases if $l_1/n_1 - l_2/n_2 \notin \mathbb{Z}$, then X_1 is diagonal.

This section can be applied to obtain conjugations to normal forms in surfaces of type \mathbb{C}^2/Γ where Γ is a finite subgroup of $SU(2)$. These surfaces are embedded in \mathbb{C}^3 with an isolated singularity at the origin [KLE]. If Γ is diagonal, then it is cyclic and generated by

$$\begin{pmatrix} e^{2\pi\sqrt{-1}/n} & 0 \\ 0 & e^{-2\pi\sqrt{-1}/n} \end{pmatrix}$$

Proposition 6 applies in this case: $\eta_1 = -\eta_2 = 1$, $l_1/n_1 - l_2/n_2 = 2/n \notin \mathbb{Z}$ if $n \neq 2$

When Γ is non-diagonalizable, we have the groups which are the inverse image of the covering surjection $\rho : SU(2) \rightarrow SO(3)$ of the groups of index 2 (preserving orientation) of triangular spherical groups [MIL], namely: The binary dihedral group of order $4p$, the binary tetrahedral group of order 24, the binary octahedral group of order 48 and the binary icosahedral group of order 120.

ACKNOWLEDGMENTS

The author wishes to thank Xavier Gómez-Mont for his helpful comments and remarks concerning this work. He would also like to thank Professor Abdus Salam, the International Atomic Energy Agency and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste.

Bibliography

- [BR] Brjuno, A.D., Analytic forms of Differential Equations, Trans. Moscow Math. Soc., vol. 25, 1971, 131-282.
- [CA] Cartan, H., Quotient d'un espace analytique par un groupe d'automorphismes, Algebraic Geometry and Topology, Princeton U.P., 1957.
- [CH] Chaperon, M., Invariant Manifolds and a preparation lemma for local holomorphic flows and actions. Holomorphic Dynamics, Proceedings of Springer Verlag, 1986, Lecture Notes in Math 1345.
- [DU] Dulac, H., Solutions d'un systeme d'équations différentielles dans le voisinage des valeurs singulières, Bull. Soc. Math. France. 40, 1904.
- [HU] Humphreys, J., Introduction to Lie Algebras and representation theory Springer-Verlag.
- [KLE] Klein, F., Lectures on the Icosahedron and the solution of the equations of fifth degree, Teubner 1884, Dover 1956.
- [MIL] Milnor, J., On the 3-dimensional Brieskorn manifolds $M(p, q, r)$ Ann. of Math. studies, Princeton U. P., study 84, 1975.
- [SI] Siegel, C.L., Über die normalform analytischer Differentialgleichungen in der Nähe einer Gleichgewichtslosung, Nächr. Akad Wiss. Gottingen Math-Phys. k1. Math.-Phis.-Chem. Abt 1952, 21-30.