


FR 9201191

I.P.N. BP n°1 - 91406 ORSAY

institut de physique nucléaire
CNRS - IN2P3 - UNIVERSITÉ PARIS - SUD



IPNO-DRE-91.10

FINE STRUCTURES IN ^{14}C EMISSION OF ^{223}Ra AND ^{224}Ra

E. Hourani, L. Rosier, G. Berrier-Ronsin, A. Elayi,
A.C. Mueller, G. Rappenecker, G. Rotbard, G. Renou,
A. Lièbe and L. Stab

Institut de Physique Nucléaire
F-91406 Orsay - Cedex, France

H.L. Ravn
Experimental Physics Division, CERN
CH-1211 Geneva 23, Switzerland

Fine structures in ^{14}C emission of ^{223}Ra and ^{224}Ra

E. Hourani, L. Rosier, G. Berrier-Ronsin, A. Elayi, A.C. Mueller,
G. Rappenecker, G. Rotbard, G. Renou, A. Lièbe and L. Stab
Institut de Physique Nucléaire
F-91406 Orsay - Cedex, France.

and

H. L. Ravn
Experimental Physics Division, CERN
CH - 1211 Geneva 23, Switzerland

Abstract

The measurement of the energy spectrum of ^{14}C nuclei emitted in the spontaneous radioactivity from ^{223}Ra and ^{224}Ra has been carried out, using thin and intense sources (480 MBq for ^{223}Ra and 3550 MBq for ^{224}Ra). The sources were obtained by implanting mass-separated beams from ISOLDE (CERN) into Al and vitreous C catchers. The measurement was performed with the supraconducting solenoidal spectrometer SOLENO installed at Orsay. Our discovery, previously reported, of a fine structure in the energy spectrum of ^{14}C emission from ^{223}Ra , which is analogous to the one known for α emission, is confirmed. Only 13% of the branching ratio in ^{14}C decay leads to the ground state of the residual nucleus, while 81% to the first excited state. For ^{14}C emission of ^{224}Ra , a lower limit of 2 for the hindrance factor has been measured for the transition to the first excited state in the residual nucleus. Also, a precise identification in Z with a $E \times \Delta E$ telescope has been performed for the radiation from the ^{223}Ra source. Our measurements of fine structures in ^{14}C emissions opens this field to nuclear structure studies.

PACS numbers: 23.90.+w, 27.90.+b

1- Introduction

Recently (1984), a natural ^{14}C radioactivity has been observed¹ from Ra isotopes, in parallel to their α decay, with branching ratios of $\approx 10^{-10}$. Emissions of Ne, Mg and Si have been consecutively observed from uranium and heavier elements with branching ratios ranging from 10^{-12} to 10^{-17} . A common feature of all these emissions is that heavier nuclei emit heavier fragments in such a way that the daughter is always the double magic ^{208}Pb or a closely neighboring nucleus. This fact is easily explained by the favorable energetic balance in the decay of heavy nuclei toward the ^{208}Pb region. Two review papers of this field are given in references 2 and 3.

Various techniques have been used so far. In particular, track detectors have been found to be the most appropriate to measure emissions with the lowest branching ratios. At Orsay, we used the magnetic spectrometer SOLENO to confirm⁴ the first discovery, i.e. the ^{14}C radioactivity of ^{223}Ra , then to give evidence⁵ for the ^{14}C radioactivity of ^{226}Ra .

The logic of searching for new emitters with ever smaller branching ratios ($^{14}\text{C}/\alpha$) was the only rule until 1989 when, at Orsay, in an experiment with the magnetic spectrometer SOLENO we found evidence for a fine structure in the energy spectrum of ^{14}C ions emitted from ^{223}Ra .⁶

Concerning the theoretical aspect, fission-like models relying simply on the Q value of the decay and on a phenomenological potential barrier have been used since 1980 and improved thereafter.⁷⁻⁹ They have shown a high predictational power in the types and the branching ratios of these new exotic emissions. As to whether excited states in the residual nucleus are populated or not in these emissions, the fission-like calculations could only appreciate the dynamical aspect predicting an attenuation of two orders of magnitude for an increase of 1 MeV in excitation energy.

According to the fission-like calculations, the decay of ^{223}Ra into ^{14}C goes mainly to the ground state of ^{209}Pb and only with a fraction of 2% to its first excited state. The results of the fine structure experiment of Orsay came in a complete disagreement showing a predominant transition to the first excited state (81%) and a weaker transition (15%) to the ground state. They are rather similar to the well-known results of fine structure in α decay, in which transitions to ground state in residual nuclei are hindered for odd-A parent nuclei. Clearly, these facts are a strong signature of nuclear structure effects requiring microscopic calculations for the interpretation.

At present, a precise measurement of the energy spectrum of ^{14}C ions emitted by Ra sources, as well as the precise identification in A or Z of such ions, are only possible by means of a spectrometer like SOLENO¹⁰, because of both its high rejection for α radiation and its large entrance solid angle (≈ 200 msr).

The significant experimental development at Orsay, after our fine structure experiment of July 1989, was the use of a new type of radioactive Ra sources. These sources are produced by implantation at ISOLDE (CERN). Compared to our earlier sources which consisted of

various chemical deposits, the new sources are of higher intensities, lower thicknesses and better purity. These new sources produced at ISOLDE were used in the experiment which will be described in this paper, in order to confirm our precedent result obtained with ^{223}Ra source and to extend it to a ^{224}Ra source. In addition, a precise measurement with Z-identification of the emitted radiation has been carried out using the ^{223}Ra source.

2- Choice of ^{223}Ra and ^{224}Ra sources

Taking advantage of the possible production at ISOLDE of more than one source at one time, we decided to produce simultaneously two sources of ^{223}Ra and ^{224}Ra . Indeed, the re-measurement of ^{223}Ra with a source of a new production type was interesting in order to confirm and to improve our earlier result. The choice of ^{224}Ra , on the other hand, was based on the discussion of the Geiger-Nuttall diagram plotted for ^{14}C emission from Ra isotopes, in analogy to α emission.

A Geiger - Nuttall diagram consists of plotting $\text{LOG } T_{1/2}$ versus $Q^{-1/2}$ for each emission, $T_{1/2}$ and Q being the half-life and the energetic balance respectively. Figure 1 shows the diagrams of both α and ^{14}C emissions for Ra isotopes.

The left hand side of the figure contains the α emission data taken from reference 17. It is seen that transitions to ground state and first excited state from even-even nuclei, i.e. ^{222}Ra , ^{224}Ra and ^{226}Ra , define a straight line, while data corresponding to odd - A nuclei, i.e. ^{223}Ra , fall apart. The vertical deviation from the straight line is called hindrance factor and is related to nuclear structure properties.

The right hand side of figure 1 corresponds to the results of the known even-even ^{14}C emitters, i.e. ^{222}Ra , ^{224}Ra and ^{226}Ra .^{5,11} The Q values were calculated under the assumption that the residual nucleus has been left in its ground state. The fact that the data lie approximately on a straight line means that this assumption is almost satisfied and that the analogy with α emission is meaningful. The results of the only odd - A ^{14}C emitter, i.e. ^{223}Ra , fall outside the line, similarly to α emission. Now, conversely, we rely on the analogy with α emission to assume that representative points of the transitions to first excited state, for even - even ^{14}C emitters, come on the straight line in the diagram. This allows to deduce their branching ratios, which hence have been predicted equal to 0.01%, 2.5% and 1% of the ground state for ^{222}Ra , ^{224}Ra and ^{226}Ra respectively.

Our interest has been fixed on ^{224}Ra among the other even - even Ra isotopes for the following reasons:

- For ^{222}Ra , three measurements have been carried out.^{5,11,12} All of them had a sufficient energy resolution to resolve the first excited state of ^{208}Pb at 2.6 MeV excitation energy location. No meaningful counts have been observed at this location. An appropriate search with the spectrometer SOLENO for a fine structure (predicted above for this isotope to be of $\leq 10^{-5}$) would require an unrealistic statistics.

- For ^{226}Ra , two measurements have been performed with a same thick source (2 mg/cm²) made at Orsay.^{5,20} The thickness of the source is inherent to both the relatively high

activity (2.6 mCi) of the source and its long half-life (1600 y). The poor energy resolution expected from such a thick source together with the low branching ratio to excited states in the residual nucleus make the feasibility of the search of a fine structure for ^{226}Ra quite questionable.

- For ^{224}Ra , the above predicted branching ratio (2.5%) to the first excited state is the highest, compared to those of ^{222}Ra and ^{226}Ra . In addition, the only preceding measurement¹¹ for this emitter was performed in track detectors with a poor statistics (22 events) and an energy resolution of about 2 MeV which is insufficient to resolve the first excited state of ^{210}Pb located at 0.795 MeV excitation energy. Thus, a search for a fine structure of this isotope is promising.

3 - Production of ^{223}Ra and ^{224}Ra sources

Radium and Francium isotopes were produced by means of the ISOLDE-2 mass-separator¹³ at the Synchrocyclotron of CERN. A 2.8 μA proton beam of 600 MeV and a thick thorium carbide target¹⁴ of 55 g/cm^2 were used during a 3 day irradiation. The ^{223}Ra and ^{224}Ra sources were produced by implantation of the A=223 ($^{223}\text{Ra} + ^{223}\text{Fr}$) and A=224 ($^{224}\text{Ra} + ^{224}\text{Fr}$) singly-charged ions of 60 keV. The ^{223}Fr and ^{224}Fr isotopes are short-lived β -emitters which transform into ^{223}Ra and ^{224}Ra respectively, contributing to the production of the Ra sources. The intensities of the collected beams are displayed in figure 2 which shows a mass spectrum of (Ra + Fr) isotopes scanned by means of a wire at the focal chamber. The A=223 ions were implanted into an Al catcher mounted in an external beam line. A uniform electrostatic scanning over an area of 15 \times 6 mm was made. The implantation of A=224 ions was performed into a vitreous carbon plate (4 mm wide and 8 mm high) placed in the focal chamber of ISOLDE. The vitreous carbon was chosen as implantation material because of both its low coefficient of sputtering and its mechanical hardness useful for subsequent handling. Because ^{224}Ra is the next neighbor to ^{223}Ra , it was not possible to produce it simultaneously in a second beam line where a scanning would have helped to decrease the consequences of the sputtering.

Indeed, sputtering of atoms from the implantation matrix is the main problem that one encounters when producing an intense source of a small area. This problem was particularly serious for ^{224}Ra source which was produced with a rate of 8×10^9 of A=224 ions per second. We estimated a sputtering of a total number of 10^{16} atoms from the carbon matrix. Assuming that the ions are implanted within a depth of 200-400 \AA , an implantation area larger than 7 mm^2 was required to avoid reaching the saturation into the implantation.¹⁵ Therefore, to increase the implantation area, the beam spot was regularly moved over the carbon plate during the irradiation: horizontally, by moving the carriage on which the plate was fixed and vertically by moving the beam itself.

4 - Control measurement of the sources

The aim of the control measurement is to determine the intensity of the sources and to check their isotopic purity and the energy resolution of the emitted α rays. The measurement was performed, after transporting the sources rapidly to Orsay, by arranging a source and a Si detector in direct view with each other inside a vacuum chamber. The source to detector distance was taken large ($\approx 2\text{m}$) and a diaphragm of a small diameter ($\phi=5.5\text{ mm}$) was placed in front of the detector, so that an α -counting rate of only a few hundreds per second was obtained.

Let us say, here, by anticipation, that the use of these sources with the spectrometer SOLENO has required to deposit on the sources by vaporization an Al layer of 2000 \AA . So, in the present section, after a general information on ^{223}Ra and ^{224}Ra decay chains, we describe the α spectra of the sources measured before then after the vaporization of Al.

Figures 3-a and 4-a give the decay chains of ^{223}Ra and ^{224}Ra and the most intense α rays of their chain members. The rays were numbered up to 12 for ^{223}Ra and up to 7 for ^{224}Ra . The plot displays the intensity of these rays versus their energy, the intensity having been normalized to the total Ra parent one. One has to note the presence of a Rn isotope in each of the two decay chains. Indeed, the Rn, as a gas, will emanate from the source and will then cause various troubles.

Not only Rn is an α -emitter, but also it generates daughters which are themselves α -emitters and which, in addition, involve one member of a relatively long half-life ($T_{1/2}=36.1\text{ m}$ for ^{211}Pb and 10.64 h for ^{212}Pb). Consequently, when a source is handled in air, a special health protection against radon is required and when a source is measured inside a vacuum chamber a serious α radiation background from the daughters deposited on the wall of the chamber is encountered.

Figures 3-b and 4-b show the α radiation energy spectra of ^{223}Ra and ^{224}Ra sources respectively, these spectra having been established before the vaporization of Al on the sources.

In figure 3-b, peaks labelled 11 and 12 which correspond to ^{219}Rn and its next daughter ^{215}Po , present two anomalies: (i) their intensities compared to the ones of peaks 1-6 corresponding to ^{223}Ra are much higher than expected from figure 3-a and (ii) they have large low energy bumped tails. These anomalies are due to the Rn escaped from the source and to its next daughter which both have decayed in the neighborhood of the detector under a large solid angle. Apart from these apparent anomalies, all of the expected α rays are normal. In particular, the α rays corresponding to ^{223}Ra are well resolved and show an energy resolution of 23 keV . If we take into account the resolution of 20 keV achieved by the same detector with the 5.480 MeV α -ray of an ^{241}Am standard source, we can say that the energy degradation due to the ^{223}Ra source is small. This means that the type of the sources obtained by implantation at ISOLDE is of high quality and appropriate to α and ^{14}C emission spectroscopic studies. As to the activity of the source, it was deduced from the area of the ^{223}Ra peaks and was found equal to 480 MBq (13 mCi) in ^{223}Ra at the end of the irradiation.

In figure 4-b corresponding to ^{224}Ra source, the peak with a low energy tailing lying between peaks 5 and 6 is spurious. It is due to ^{211}Bi traces left in the vacuum chamber after the preceding measurement with ^{223}Ra source (it corresponds to peak n° 10 in figure 3-b). The figure displays a good quality spectrum characterized by an α ray energy resolution of 28 keV and an activity of the source of 3550 MBq (96 mCi) in ^{224}Ra normalized to the end of the irradiation.

Finally, looking for common rays in the spectra of the two sources due to isotopic mixing in the production, peak n° 12 of the decay chain of ^{223}Ra was revealed in the spectrum of ^{224}Ra source and peak n° 7 of the decay chain of ^{224}Ra was revealed in the spectrum of ^{223}Ra . The cross contaminations, i.e. the activity ratio of ^{223}Ra to ^{224}Ra in the ^{224}Ra source and of ^{224}Ra to ^{223}Ra in the ^{223}Ra source, were determined to be $\leq 1/1000$ and $\leq 5/1000$, respectively, at the end of the irradiation.

Figures 3-c and 4-c display the α -ray energy spectra of ^{223}Ra and ^{224}Ra sources respectively, these spectra having been measured after the vaporization of Al on the sources. In both figures, the relative height of the various peaks is in agreement with the expectation from figures 3-a and 4-a. This means that most of Rn and its daughters decay inside the source itself and consequently that the emanation of Rn is much lower than without coating with Al. This analysis is reinforced by the fact that a small peak appears at the right side of the radon peak (peak n° 11 in figure 3-c and peak n° 5 in figure 4-c). The small peak is due to Rn nuclei having decayed after emanation from the source, because its shift in energy of 40 keV with respect to the main peak of Rn corresponds to the α energy loss in the Al deposit on the source. The spurious peak at the left side of peak n° 6 in figure 4-c is due to ^{211}Bi traces from ^{223}Ra source measured this time also before ^{224}Ra source. As to the energy resolution measured for the α rays of Ra, the two sources gave a value of 40 keV, still quite acceptable for the success of the experiment. Unfortunately, there is a low-energy tailing for α - rays of Ra which is more pronounced in the case of ^{223}Ra source. This may indicate that the Al vaporization on the Al matrix was not as successful as on the carbon one.

5 - Setup with SOLENO

Let us recall here that SOLENO is a superconducting coil surrounding a vacuum chamber ($\phi=360$ mm) on the axis of which the source is placed at one side and the detector at the other side. When the electric current is set to focus on the detector $^{14}\text{C}^{6+}$ or $^{14}\text{C}^{5+}$ ions emitted by the source, the high flux of α radiation emitted by the same source are focussed outside the detector. A scheme of the setup is shown in figure 5.

5.1 - Transmission curve

The focussing of SOLENO is characterized by the effective solid angle Ω at its entrance. For a fixed source - detector geometry, Ω depends on a single parameter y such that:

$$y = B\rho/I \tag{1}$$

where, $B\rho$ is the magnetic rigidity of the ions emitted by the source and I is the electric current in SOLENO. The curve obtained by the plot of Ω in terms of y is called the transmission curve of SOLENO. For each source - detector geometry used in this work a transmission curve has been measured, using an α source of calibrated intensity and known $B\rho$, and varying the current I of SOLENO. The transmission curve corresponding to a quasi punctual source and to a single Si detector with a $\phi= 22$ mm diaphragm in front is shown in figure 6. The same figure indicates the current values used in the various measurements and the locations (indicated with arrows) corresponding to the ^{14}C ions which populate the low-lying excited states in the residual nucleus.

5.2 Detectors

In the search for a fine structure in the energy spectrum of ^{14}C ions from ^{223}Ra and ^{224}Ra , a single Si detector was used in view to reach the best energy resolution. It was a passivated implanted planar detector (made by Canberra), partially depleted, with a thickness of $100\ \mu\text{m}$ and an area of $450\ \text{mm}^2$. Its energy resolution was of $18\ \text{keV}$ for α particles and $110\ \text{keV}$ for ^{12}C particles of $30\ \text{MeV}$.

In the measurement with a telescope for an identification in Z , surface barrier Si detectors manufactured at the IPN of Orsay were used. The E detector was partially depleted, $300\ \mu\text{m}$ thick and with an area of $450\ \text{mm}^2$. The ΔE detector was of epitaxial type, totally depleted, $9\ \mu\text{m}$ thick and with an area of $250\ \text{mm}^2$. A detailed description for the ΔE detector is given in Fig. 6 of reference 16.

5.3 The α background

A serious problem expected to occur in the measurement of such intense sources with SOLENO is a too high background from α particles. This may cause strong multiple pileup and even damage rapidly the detector.

One origin of α particles is radon emitted from the source and traveling as a neutral gas inside the vacuum chamber of SOLENO. The radon and its daughters decay in the neighborhood of the detector producing an overall α background which is independent of the current setting of SOLENO but proportional to the detector area. In order to reduce this type of background, an efficient pumping with two turbomolecular pumps in addition to the standard cryogenic pump was used. Also, a diaphragm ($\phi=16\ \text{mm}$) destined to cover half of the detector area was prepared.

The other origin of α particles falling on the detector is the source itself which emits a tremendous flux of energetic (5-8 MeV) singly- or doubly-charged α radiation. In the previous uses of SOLENO, the current was set for a magnetic rigidity $B\rho$ of $^{14}\text{C}^{j+}$ at the maximum of the transmission curve. This was not possible for the source of ^{224}Ra , because of the proximity of the $B\rho$ of $8.784\ \text{MeV}\ \alpha^{++}$ emitted by ^{216}Po which is a member of the decay chain of ^{224}Ra . Therefore, the $^{14}\text{C}^{5+}$ state was rather focussed in our measurement.

5.4 - Charge state distribution

We needed to know the percentages of $^{14}\text{C}^{5+}$ and $^{14}\text{C}^{6+}$ charge states in an interval of a few MeV around a ^{14}C ion energy of $30\ \text{MeV}$. Under the assumption that ^{12}C and ^{14}C

atoms of equal velocity have approximately the same charge state distributions, we measured such distributions for a ^{12}C beam of about 25 MeV delivered by the Tandem at Orsay. To simulate the Al deposit on ^{223}Ra and ^{224}Ra sources, the ^{12}C beam had to cross an Al foil of $100\mu\text{g}/\text{cm}^2$.

Two methods of measurement were performed. In one simple method, the analyzed $^{12}\text{C}^{4+}$ beam of the Tandem was let to cross an Al foil of $100\mu\text{g}/\text{cm}^2$. At the exit of the foil the charge equilibrated $^{12}\text{C}^{6+}$, $^{12}\text{C}^{5+}$ and $^{12}\text{C}^{4+}$ beams were consecutively switched towards a Faraday cup and their intensity measured. In the second method, the $^{12}\text{C}^{4+}$ beam hit a $100\mu\text{g}/\text{cm}^2$ Al target and the elastically scattered $^{12}\text{C}^{6+}$ and $^{12}\text{C}^{5+}$ ions were focussed with a Split-Pole spectrometer placed at 9° and measured with a multiwire proportional counter. The two methods were in agreement.

The results of the two measurements were plotted in figure 7 together with the results of a ^{12}C beam incident on a C foil taken from the compilation of K. Shima et al.¹⁸ and which are connected by continuous lines. We read, for our use, the percentages of 5^+ and 6^+ charge states on the dashed curves which are parallel to the continuous lines and also fit our two results. The data measured by W. Kutschera et al.¹⁹ at Argonne are also shown on the same figure. They are in agreement with our results.

6- Fine structure in ^{14}C radioactivity of ^{224}Ra

In a preliminary measurement, the current of SOLENO was set to focus $^{14}\text{C}^{5+}$ ions. A too high counting rate of α particles originating from radon was recorded and, in addition, noticeable traces of ^{224}Ra sputtered from the source were detected. Having realized that both troubles were originating from the superficial lying of Ra atoms in the source (due to the removal of the 200 \AA matrix layer by sputtering during the irradiation), we decided to deposit on the source by vaporization a 2000 \AA layer of Al, in order to reduce the emanation of radon and to prevent the sputtering of Ra. The deposit was achieved successfully. The emanation of radon was reduced by a factor 5. The sputtering of Ra was completely stopped and the energy resolution in the α spectrum measured in direct view of the source, was only slightly degraded from 28 to 40 keV.

The measurement with SOLENO consisted of a 6.3 d run, the current of SOLENO being set at the value of 330 A which focus $^{14}\text{C}^{5+}$ ions emitted from ^{224}Ra . Having started the measurement with a diaphragm of $\phi=16\text{ mm}$ placed in front of the detector, this diaphragm was exchanged for another one of $\phi=22\text{ mm}$, 2.2 d later when the activity of the source had decreased by a factor of 1.5.

All along the running of the measurement, the multiple pileup has been closely controlled. Indeed, taking into account the energy range (5.449-8.784 MeV) of the α rays generated by ^{224}Ra and the members of its decay chain, the triple pileup has been expected to extend in energy until a higher edge of 26.353 MeV, which is due to the triple pileup of the 8.784 MeV α ray. On the other hand, as the energy of ^{14}C ions feeding the ground state

of the residual nucleus ^{210}Pb is 28.631 MeV, therefore, an interval of 2.3 MeV, between the ground state position and the upper limit of the triple pileup, was left to observe the four low-lying excited states in ^{210}Pb . In fact, the measured α counting rate has not exceeded 3000 counts/sec during the experiment. The quadruple pileup which could disturb our measurement was 9 orders of magnitude lower. So, only a few counts, i.e. not more than 3 or 4, could have occurred during the whole measurement and, in addition, these should have fallen on a wide range of energy.

Transmission curves of SOLENO have been established before and after the measurement for the two diaphragms of $\phi = 16$ and 22 mm used consecutively in front of the detector. The curve corresponding to $\phi = 22$ mm diaphragm has been found to be 1.41 higher than the other. This curve is presented in figure 6, where the arrows indicate the locations of the ground state (label a) and the first and the second excited states (labels b and c) of ^{210}Pb . The current setting of SOLENO was chosen to place the excited states at the maximum of the transmission curve.

The results of the measurement are reported in figure 8, where a very sharp peak is seen. The peak contains 149 events. It shows an energy resolution of 150 keV. It is located exactly at the position expected for the ground state of ^{210}Pb , as deduced from the energy calibration with ^{14}C emission of ^{223}Ra source. We conclude that this peak corresponds to the transition of ^{224}Ra by ^{14}C radioactivity towards the ground state of the residual nucleus ^{210}Pb . At the left of the figure a few triple pileup events are seen and the upper edge of the triple pileup is indicated. In between the ^{14}C ground-state peak and the α triple pileup region, the locations of the four low-lying excited states of ^{210}Pb are indicated with arrows. No event at all is seen at these locations. This fact means that within the sensitivity of our experiment no transition of ^{224}Ra by ^{14}C emission towards the low-lying excited states of ^{210}Pb is observed.

We calculated for ^{224}Ra the branching ratio BR of ^{14}C emission with respect to α decay according to the following formula

$$n = BR \times N \times \Omega \times \epsilon / 4\pi \quad (2)$$

where, n is the number of ^{14}C nuclei which have been detected (149), N is the total number of ^{224}Ra nuclei having decayed during the measurement (5.01×10^{14}), Ω is the effective solid angle of SOLENO (161 msr for a $\phi = 22$ mm diaphragm) and ϵ is the 5^+ charge state percentage of the ^{14}C ions emitted by ^{224}Ra (at 28 MeV a value of 32.5% has been used). Taking into account the numerical values and the change in transmission for two different diaphragms, a branching ratio ($^{14}\text{C}/\alpha$) of $(6.5 \pm 1.0) \times 10^{-11}$ has been deduced. It is slightly higher than the only one previously reported¹¹ equal to $(4.3 \pm 1.2) \times 10^{-11}$. Nevertheless both results are in agreement within their respective uncertainties. On the other hand, as no event was detected for the first excited state of ^{210}Pb , a lower limit of 2 for the hindrance factor (confidence level of 90%) has been deduced. Two final conclusions can be drawn:

(i) a sensitivity in ^{14}C emission measurement to a branching ratio of 4×10^{-13} with respect to α decay has been reached, which is a record for the setup with SOLENO, and (ii) the measured hindrance factor limit being of nuclear structure origin appeals for a spectroscopic interpretation.

7- Confirmation of the fine structure in ^{14}C emission from ^{223}Ra

Our main goal with the ^{223}Ra source was a confirmation of our previous result⁶ which gave evidence for a fine structure. Naturally, we had in view to achieve a better statistics and a better energy resolution.

On the other hand, the present measurement with ^{223}Ra source was immediately consecutive to the one with ^{224}Ra source. The same detection conditions for the two sources were maintained in order to have an identical energy calibration.

Five successive runs have been carried out for different values of the current in SOLENO. The current values have been chosen such that the transitions towards the ground state (label 0) and the excited states (labels 1, 2 and 3) in ^{209}Pb (figure 6) had various locations on the transmission curve.

A total number of 700 events has been detected, twice the statistics of our previous measurement.⁶ The energy spectrum is given in figure 9. Two prominent peaks are clearly present. They correspond to the ground state and to the first excited state in ^{209}Pb . This result definitely confirms the fine structure in the energy spectrum of ^{14}C radioactivity from ^{223}Ra .

The main part of the peaks has a width of 150 keV which is better than our previous value of 240 keV. Unfortunately, the shape of the peaks is not gaussian and there is a low energy tailing, which is more important than in the case of ^{224}Ra source, as expected from the α - ray control spectra (see section 4).

As no peak of the characteristic width (150 keV) is seen over the location of the second and higher excited states, we can conclude that none of these has been detected.

Nevertheless, it should be interesting to repeat this experiment in the near future with improved sensitivity. This should be possible after the installation of ISOLDE at CERN's PS-booster. We plan to do it benefitting from our experience gained in the choice of source implantation matrices and in the control of Al deposition.

For the calculation of the branching ratio ($^{14}\text{C}/\alpha$) we label a given run with an index i specifying the current of SOLENO and call N_i the total number of ^{223}Ra nuclei which have decayed.

Furthermore, the excitation energies in ^{209}Pb is divided into intervals labelled j . With these notations, the branching ratio $(BR)_j$ is given by

$$n_{ij} = (BR)_j \times N_i \times \Omega_{ij} \times \epsilon_j / 4\pi \quad (3)$$

where, the quantity n_{ij} is the number of ^{14}C nuclei which has been measured in the interval j during the run i , Ω_{ij} is the effective solid angle of SOLENO corresponding to the interval j in the run i and ϵ_j is the 6^+ charge state percentage of the $^{14}\text{C}^{6+}$ ions populating the interval j .

The total ^{14}C emission branching ratio with respect to α emission was found equal to $(7.0 \pm 0.4) \times 10^{-10}$, whereas the branching ratio to the ground state of ^{209}Pb was equal to $(0.91 \pm 0.20) \times 10^{-10}$ instead of the value 6×10^{-10} previously used in predictional models. The partial branching ratios to the ground state and to the first excited state of the residual nucleus have been deduced equal to $(13 \pm 3)\%$ and $(81 \pm 6)\%$ respectively. All these numbers concerning the various calculated branching ratios are in agreement with those obtained in our first measurement (table 1) and do not appreciably modify figure 1 which is made from our previous results. In figure 10, we report for comparison, our old result together with the present. We note that both results exhibit a moderate sensitivity for the detection of higher excited states with small branching ratios, due to low energy tailing.

8 - Identification with a telescope

In the previous measurements of ^{14}C emission, either with track detectors or with Si telescopes, none was focussed to a search for a fine structure in the atomic number Z of the emitted light nuclei. The measurement, described hereafter, is the first to have been optimized, in statistics and in detection quality, in order to be sensitive to a few percent fine structure in Z .

The source of ^{223}Ra was used and the single detector was replaced by the Si $E \times \Delta E$ telescope. The current of SOLENO was set to the value of 284 A already used to focus $^{14}\text{C}^{6+}$ ions emitted by ^{223}Ra . The focussing is illustrated by the positions of the arrows on the transmission curve in figure 6.

A measurement of 6 days was performed. A number of 210 events were recorded and are given in a bidimensional $E \times \Delta E$ plot (fig. 11). One has to notice that in this diagram the quantity E is the response of the back detector, i.e. the residual energy, instead of the total energy more conventionally used in the literature. A small size dot represents a single event whereas a big size dot represents a set of 10 grouped events.

One can see that more than one half of the total number of events are concentrated over a small region, while the rest are mainly spread along two lines, one line being horizontal and the other with a slope such that $E \times \Delta E = \text{const}$.

In the same figure, there are reported the results of a calibration with ^{12}C and ^{16}O beams of different energies delivered by the Tandem of Orsay. Contour limits at 1/5 and 1/50 of the maximum of the telescope response to each beam have been drawn.

In an overall comparison, it clearly appears that C and O beam locations fall on two distinct horizontal lines and that the ^{223}Ra source results lie very close to the 29 MeV C beam position (the shift in the location was caused by electronic signal attenuation due to

longer cables for the measurement with beams). This fact illustrates the conclusion that at least a large part of ^{223}Ra source results corresponds to C and not to O or other ions. Now, analyzing the spread of the source results, one can explain the horizontal extension by the natural width in energy of ^{14}C nuclei emitted by ^{223}Ra . As to the few events distributed over $E \times \Delta E = \text{const.}$ line, they are compatible with C ions having undergone a channeling effect in ΔE detector.

Thus, at the present level of statistics, no fine structure in Z was visible.

9 - Discussion and conclusions

The present work has associated two unique facilities, ISOLDE as a source production system and SOLENO as a large solid angle (200 msr) spectrometer.

The quality criteria of the sources are the energy resolution for α -rays and the isotopic purity. The α energy resolution measured in this work was excellent and very close to the limiting value (≈ 20 keV) obtained with standard α -sources. As to the cross contamination by neighboring Ra isotopes, it has not exceeded a few parts per thousand.

On the other hand, one had to suspect that the sputtering during the implantation sets a saturation limit on the activity, in the production of sources with small areas (a few units to a few tens of mm^2). In fact, the saturation should be reached beyond several days of irradiation, which is a too long period in a normal irradiation request at ISOLDE.

Indeed, at the first measurement of the α spectrum of ^{224}Ra source, we realized that the intensity of this source was exceptional and hence appropriate to overcome the weak ^{14}C emission from ^{224}Ra (one order of magnitude lower than that from ^{223}Ra). So, despite a difficult handling due to the presence of a long half - life daughter (^{212}Pb , $T_{1/2}=10.64$ h) and a high energy γ emitter (^{208}Tl , $E_\gamma=2.6$ MeV) in the decay chain of the emanating radon, a very precise result has been obtained, clearly improving the previously measured value for the branching ratio of ^{14}C emission from ^{224}Ra and giving evidence for the existence of a hindrance factor in ^{14}C emission to the first excited state in the residual nucleus.

The measurement of ^{223}Ra source served three purposes. The first one was to confirm the fine structure in the energy spectrum of ^{14}C emission from ^{223}Ra that we have discovered using a chemically deposited ^{227}Th source. The fine structure result was unambiguously reproduced with the new type of ^{223}Ra source. The second purpose was to carry out a precise identification in Z. This also was successfully achieved and showed that all the events fall at the C location as determined with a C beam. As to the third purpose, it was the use of the ground state location in the energy spectrum in order to calibrate the energy scale in the ^{224}Ra measurement. Indeed, with this calibration, the observed peak in the ^{224}Ra measurement came at the ground state location expected for ^{14}C ions emitted from ^{224}Ra nuclei.

During this work, we have experienced the production, the handling and the measurement with SOLENO of Ra sources produced at ISOLDE. In particular, we have experienced

two different implantation matrices, i.e. Al and vitreous carbon, and the Al vaporization on the sources. After our present success, we plan to perform a next experiment with again a ^{223}Ra source under optimized conditions to attain an ultimate energy resolution. Such a resolution together with a good statistics are required to resolve the second and higher excited states in the residual nucleus. In addition, a precise measurement would tell us if there are events occurring in the energy spectrum at the location of ^{13}C emitted from ^{223}Ra . Such an occurrence was suggested by the mass identification results of Kutschera et al.¹⁹ which yielded one event out of 24 at the ^{13}C location.

Already in 1986, the plot of the branching ratio of ^{14}C emissions versus the mass number of the parent nucleus showed an odd-even effect analogous to the one of the α emissions.²⁰ At present, we explain this odd-even effect as a manifestation of a large hindrance for transitions from odd-A parent nuclei to the ground state of their daughter nuclei. In particular, we expect ^{231}Pa and ^{233}U which are odd-A Ne emitters to have in their decays transitions toward the excited states of their daughters.

In fact, only the outlines of this type of radioactivity have been explained so far.^{8,9} It has been explained as a superasymmetric fission toward the ^{208}Pb region governed by the Q value of the decay and by an appropriate potential barrier. This fission reproduces approximately the absolute branching ratios but fails in the explanation of the observed hindrance factors which are connected to the overlapping of microscopic wavefunctions of the parent nucleus and the emitted fragments. A microscopic calculation similar to the one initiated by R. Blendowske et al.^{21,22} should be developed and has to explain not only the transition observed from ^{223}Ra to the first excited state of ^{209}Pb , but also the hindrance which we have observed in this work from ^{224}Ra to the first excited state in ^{210}Pb . It is probably a hindrance having a nuclear structure origin which has also lowered the ^{14}C emission from ^{221}Fr and ^{221}Ra in such a way that they have not been observed yet.²⁰

Acknowledgements

We are obliged to H. Sergolle, the Director of the IPN of Orsay, for his constant attention for the success of this experiment. We thank Mrs A. Traverse for valuable discussions on the sputtering. We thank, also, D. Sznajderman who successfully deposited by vaporization the Al layer on the sources. We acknowledge the help of J. Geurtz in the furniture of liquid helium and of Mrs R. Bzyl in the preparation of the Si detectors. We are grateful to the crew of the Synchrocyclotron and of ISOLDE at CERN for the excellent running of these facilities that allowed the obtention of high quality sources. We thank R. Deltenre at CERN for his help in handling the sources and the health service staff at the IPN of Orsay for their constant presence during the experiment. We also thank the crew of the Tandem of Orsay.

REFERENCES

1. H. J. Rose and G. A. Jones, *Nature (London)* **307**, 245 (1984).
2. E. Hourani, M. Hussonnois and D. N. Poenaru, *Ann. Phys. Fr.* **14**, 311 (1989).
3. P.B. Price, *Ann. Rev. Nucl. Part. Sc.* **39**, 19 (1989).
4. S. Gales, E. Hourani, M. Hussonnois, J. P. Schapira, L. Stab and M. Vergnes, *Phys. Rev. Lett.* **53**, 759 (1984).
5. E. Hourani, M. Hussonnois, L. Stab, L. Brillard, G. Gales and J. P. Schapira, *Phys. Lett.* **160 B**, 375 (1985).
6. L. Brillard, A. G. Elayi, E. Hourani, M. Hussonnois, J. F. Le Du, L. H. Rosier et L. Stab, *C. R. Ac. Sci. Paris t.* **309 II**, 1105 (1989).
7. A. Sandulescu, D. N. Poenaru and W. Greiner, *Sov. J. Part. Nucl.* **11**, 528 (1980).
8. D. N. Poenaru, M. Ivascu, A. Sandulescu and W. Greiner, *Phys. Rev.* **C32**, 572 (1985).
9. Y. J. Shi and W. J. Swiatecki, *Phys. Rev. Lett.* **54**, 300 (1985).
10. J. P. Schapira, F. Azaiez, S. Fortier, S. Gales, E. Hourani, J. Kumpulainen and J. M. Maison, *Nucl. Instr. Meth.* **224**, 337 (1984).
11. P. B. Price, J. D. Stevenson, S. W. Barwick and H. L. Ravn, *Phys. Rev. Lett.* **54**, 297 (1985).
12. M. Hussonnois, J. F. Le Du, L. Brillard, J. Dalmaso and G. Ardisson, IPNO-DRE 9107, IPN, 91406 Orsay Cedex.
13. H. L. Ravn, *Phys. Rep.* **54**, 201 (1979).
14. T. Bjornstad, L. C. Carraz, H. A. Gustafsson, J. Heinemeier, B. Jonson, O. C. Jonsson, V. Lindfors, S. Mattsson and H. L. Ravn, *Nucl. Instr. Meth.* **186**, 391 (1981).
15. The implantation was simulated using code TRIM. Reference: J. F. Ziegler, J. P. Bier-sack and U. Littmark, *the stopping and range of ions in solids*, Pergamon Press, 1985.
16. L. Stab, *Nucl. Instr. Meth.* **A288**, 24 (1990).
17. *Table of Isotopes*, edited by C. M. Lederer and V. S. Shirley (Wiley, New York, 1978), 7th ed.
18. K. Shima, T. Mikumo and H. Tawara, *At. Dat. Nucl. Dat. Tables*, **34**, 357 (1986).
19. W. Kutschera et al., *Phys. Rev.* **C32**, 2036 (1985).
20. S. W. Barwick, P. B. Price, H. L. Ravn, E. Hourani and M. Hussonnois, *Phys. Rev.* **C34**, 362 (1986).
21. R. Blendowske, T. Fliessbach and H. Walliser, *Nucl. Phys.* **A464**, 75 (1987).
22. R. Blendowske and H. Walliser, *Phys. Rev. Lett.* **61**, 1930 (1988).

Figure Captions

Figure 1

Geiger - Nuttal diagram for α and ^{14}C emissions of Ra isotopes. The decimal logarithm of the half-lives expressed in seconds is plotted versus $Q^{-1/2}$, the energetic balance Q of the reaction being expressed in MeV.

Figure 2

The intensity of mass production at the focal chamber of ISOLDE as scanned by a wire and displayed on a oscilloscope. The peaks correspond to integer mass number A and they are composed of Fr and Ra isobars.

Figure 3

The α particle spectra of ^{223}Ra source produced at ISOLDE. In (a), the decay chain of ^{223}Ra , with the plot of the α -ray intensities versus the α energy, taken from reference 17. In (b), the α -ray spectrum of the ^{223}Ra source produced at ISOLDE, as given by a Si surface barrier detector placed in direct view of the source. In (c), as in (b) but after the deposit of an Al layer on the source by vaporization.

Figure 4

As in figure 3 but for the ^{224}Ra source produced at ISOLDE.

Figure 5

The setup with SOLENO. The drawing shows the mechanical structure of SOLENO: the coil, the iron shield and the vacuum chamber. The vacuum pumping and the source detector configuration with ion trajectories are shown.

Figure 6

The transmission curve of SOLENO, with a $\phi = 22$ mm diaphragm in front of the detector. The solid angle Ω of SOLENO is given as function of the parameter $y = B\rho/I$, where $B\rho$ is the magnetic rigidity of the ions emitted by the source and I is the current in SOLENO. The locations of the ground state and some low lying excited states in the residual nucleus of the decay of ^{223}Ra and ^{224}Ra are labelled with numbers and letters respectively.

Figure 7

The charge state distribution against the kinetic energy, for ^{12}C . The solid line connects three points taken from reference 18 and given for a ^{12}C beam incident on a C foil. The symbols (o and +) represent two measurements we have performed for ^{12}C beam incident on a $100\mu\text{g}/\text{cm}^2$ Al foil. The symbols (solid triangle) represent the results of Kutschera et al.¹⁹ for ^{14}C beam incident on a $100\mu\text{g}/\text{cm}^2$ Al foil.

Figure 8

The energy spectrum of ^{14}C ions measured with ^{224}Ra source. The arrows indicate the locations of the ground state and some low lying excited states in the residual nucleus ^{210}Pb . No events are seen at the locations of the excited states. The energy calibration has been deduced from the peak positions for ^{223}Ra source, these latter having been calibrated in a previous experiment with ^{14}C beam from the Tandem of Orsay. The energy resolution is 150 keV.

Figure 9

The energy spectrum of ^{14}C ions measured with ^{223}Ra source. The arrows indicate the locations of the ground state and some low lying excited states in the residual nucleus ^{209}Pb . The presence of the peak at 0.779 MeV is the unambiguous confirmation of the evidence for a fine structure. The main parts of the ^{14}C rays show a resolution of energy of about 150 keV.

Figure 10

Comparison of our previous result which gave evidence for the fine structure and of the present result which has confirmed the discovery. The statistics in the present result is twice higher and the energy resolution is better. Low energy tailing in both results is seen.

Figure 11

The $E \times \Delta E$ bidimensional spectrum obtained with ^{223}Ra source. The dots represent the detected events. The contour plots represent the response of the telescope to ^{12}C and ^{16}O beams of various energies delivered by the Tandem of Orsay. As expected, the event location is very close to 29 MeV C beam. The small shift in energy has an electronic attenuation origin due to longer cables used in the measurement with beam. The $E \times \Delta E = \text{const.}$ slope, either in the measurement with the source or in the one with the beams, is due to channeling effect in ΔE detector.

parent nucleus	Daughter nucleus	B.R.($^{14}\text{C}/\alpha$)	B.R.($^{14}\text{C}/\alpha$)	Ref.
		This work	Other works	
^{224}Ra	^{210}Pb G.S.	$(6.5\pm 1.0)\times 10^{-11}$	$(4.3\pm 1.2)\times 10^{-11}$	11
^{224}Ra	^{210}Pb $E_x=0.795$ MeV	$\leq 4\times 10^{-13}$		
^{223}Ra	^{209}Pb G.S.	$(0.9\pm 0.2)\times 10^{-10}$	$(1.0\pm 0.2)\times 10^{-10}$	6
^{223}Ra	^{209}Pb $E_x=0.779$ MeV	$(5.7\pm 0.4)\times 10^{-10}$	$(5.2\pm 0.4)\times 10^{-10}$	6

Table 1. Comparison of the branching ratio results of this work and of previous works.




Fig. 1




Fig. 2




Fig. 3





Fig. 4



1-iron shield 2-solenoidal coil 3-vacuum chamber
 4-source 5-iris 6-obturator 7-detector
 - - - trajectory of a focussed $^{14}\text{C}^{6+}$ ion.

Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10




Fig. 11