

Theory of Anomalous Transport in Toroidal Helical Plasmas

K. Itoh, S.-I. Itoh and A. Fukuyama

(Received - Feb. 24, 1992)

NIFS-140

Mar. 1992

This report was prepared as a preprint of work performed as a collaboration research of the National Institute for Fusion Science (NIFS) of Japan. This document is intended for information only and for future publication in a journal after some rearrangements of its contents.

Inquiries about copyright and reproduction should be addressed to the Research Information Center, National Institute for Fusion Science, Nagoya 464-01, Japan.

Theory of Anomalous Transport in Toroidal Helical Plasmas

K. Itoh, S.-I. Itoh and A. Fukuyama*

National Institute for Fusion Science, Nagoya, 464-01, Japan

*Faculty of Engineering, Okayama University, Okayama 700, Japan

Abstract

Theoretical model of the anomalous transport in Torsatron/Heliotron plasmas is developed, based on the current-diffusive interchange instability which is destabilized due to the averaged magnetic hill near edge. Analytic formula of transport coefficient is derived. This model explains the high edge transport, the power degradation and energy confinement scaling law and the enhanced heat-pulse thermal conduction.

keywords: anomalous transport, interchange instability, magnetic hill, current diffusivity, torsatron/heliotron.

1. Introduction

Much work has recently been done on the plasma transport across the magnetic surface in toroidal plasmas. This is the case for stellarators with the shear and magnetic hill such as Torsatron/Heliotron devices¹⁻³⁾. This plasma, in which the pressure gradient rather than the toroidal current plays the dominant role in instabilities, may yield a knowledge on anomalous transport complementary to tokamaks. Experiments have shown that (I) the energy confinement time τ_E degrades with power⁴⁻⁷⁾. By the comparison within different devices, a scaling law of τ_E has been proposed⁵⁾. Detailed studies have shown that (II) the effective thermal conductivity χ_{eff} increases with temperature for given minor radius, r ⁸⁾, (III) the profile of χ_{eff} for a given heating power, however, is an increasing function of r ^{7,8)}, and a simple form like $\chi_{eff} \propto T_e^{1.5}/B^2$ is not valid. It is also known that (IV) the thermal transport coefficient that is determined by the heat pulse propagation, χ_{HP} , is larger than χ_{eff} ⁹⁾. These results have similarities and differences with tokamaks, and the explanation of these will provide a key to understand the anomalous transport in toroidal plasmas.

The interchange mode¹⁰⁾ has been thought to be a candidate to explain the anomalous transport in Torsatron/Heliotron. This mode can be destabilized, for instance in the presence of finite resistivity, η . Much efforts has been done on the anomalous transport driven by the resistive interchange mode¹¹⁻¹⁴⁾. Theoretical method is either mixing length model¹⁵⁾, scale invariance method¹⁶⁾ or one/two point renormalization technique¹⁷⁾. The

different methods have given the same result on χ_{eff} from the physics point of view (the difference appears only for a numerical constant)¹⁸⁾. In spite of these intensive studies, the anomalous transport, characterized by (I)-(IV), remains unexplained.

We have recently investigated the effects of transport coefficients on the interchange instability, such as thermal diffusivity, χ , viscosity, ν , and especially, the current diffusivity, λ ¹⁹⁾. The important role of the current diffusivity was found: Below the critical beta value against global MHD mode, the microscopic interchange mode is destabilized through current-diffusivity, not by the resistivity.

In this article we derive the anomalous transport coefficient based on the microscopic current diffusive interchange mode (abbreviated by ' λ -mode'). We found $\chi_{\text{eff}} = (d\beta/d\rho)^{3/2} \delta_S^2 v_A / R$ (β : ratio of plasma pressure to magnetic pressure, ρ : normalized minor radius r/a , a : plasma minor radius, δ_S : collisionless skin depth, v_A : Alfvén velocity, R : major radius). This result is consistent with the experimental knowledge (I)-(IV).

We use a model equation based on the reduced set of equations for stellarators²⁰⁾. The cylindrical model [coordinates (r, θ, z)] is employed in order to look for the analytical insight. We also consider the case of zero equilibrium current. The model equations consist of the Ohm's law, equation of motion and the energy balance equation, as is explicitly given in Refs. [19, 21]. (The term $\lambda \nabla^2 j$ is added to the Ohm's law as $E \cdot \nabla \times B = \eta j + \lambda \nabla^2 j$.) We use the picture of the mean field theory: we analyze the growth

rate and mode structure of the λ mode by keeping x, ν, λ, η . The instability driven transport coefficient (derived by the mixing length theory) is equal to the one which is used for stability analysis.

We solve the eigen value equation by the Fourier transform. The parallel derivative is approximated by $k_{\parallel} = k_{\theta} s x$ ($k_{\theta} = m/\rho_1$, $x = \rho - \rho_1$, m poloidal mode number, ρ_1 : rational surface, $s = \rho_1 \chi'(\rho_1)$, χ : rotational transform), since we study the microscopic mode which is localized to the rational surface. The variable is changed from x to k as $u(x) = \exp(i\tau - im\theta - inz/R) \int u(k) \exp(ikx) dk$, where u is the perturbing stream function, τ is the growth rate, and n is the toroidal mode number. Eliminating the current and pressure perturbations from set equations, and assuming the electrostatic perturbation, we have the eigenvalue equation for τ in the k space as¹⁹⁾

$$\frac{1}{L^2} \frac{\partial}{\partial k} \frac{1}{\bar{\eta} + \lambda k^2} \frac{\partial}{\partial k} u + \frac{D_0 k_{\theta}^2}{\bar{\nu} + \bar{\lambda} k^2} u - (\bar{\gamma} k^2 - \bar{D} k^4) u = 0, \quad (1)$$

where $k_{\perp}^2 = k_{\theta}^2 + k^2$, $1/L = k_{\theta} s$, $D_0 = \beta_0 \Omega' p_{eq}' / 2 \epsilon^2$, $p_{eq} = p_0(\rho) / \rho_0$ (p_0 being the equilibrium pressure), $\epsilon = a/R$ and Ω' is approximately given as $\Omega' = \epsilon^2 (N/l) \rho^{-2} (\chi \rho^4)'$, (N is the toroidal pitch number and l is the multiporarity.) The term D_0 denotes the drive by the pressure gradient with the bad curvature. The transport coefficients are normalized as $\bar{\lambda} = \lambda \tau_{Ap} / a^2$, $\bar{\eta} = \eta \tau_{Ap} / \mu_0 a^2$, $\bar{\nu} = \nu \tau_{Ap} / a^2$, $\bar{\lambda} = \lambda \tau_{Ap} / \mu_0 a^4$, and the time is normalized as $\bar{\tau} = \tau \tau_{Ap}$, where $\tau_{Ap} = R \sqrt{\mu_0 m_i n_i} / B_0$, n_i is the ion density, m_i is the

ion mass and B_0 is the equilibrium magnetic field.

Equation (1) is solved by Rayleigh Ritz method. Writing Eq. (1) as $\mathcal{L}u=0$, and the functional $\mathcal{R}[u]$ is defined as $\mathcal{R}[u]=\int_{-\infty}^{\infty} u \mathcal{L}u dk / \int_{-\infty}^{\infty} u^2 dk$. Test function of $u=\exp(-\alpha^2 k^2/2)$ is employed. Equations $\mathcal{R}[u]=0$ and $\partial \mathcal{R}[u]/\partial \alpha=0$ determines the growth rate τ and α . The value α is the typical radial extent of the mode. Apart from a numerical coefficient of order unity, the transport coefficient χ is given as¹⁵⁾

$$\chi = \frac{\tau}{\alpha^2}. \quad (2)$$

The Rayleigh quotient \mathcal{R} is obtained as

$$\begin{aligned} \mathcal{R} = & -s^2 \bar{\chi}^{-1} \alpha^2 \{y^2 - 2y^3 \exp(y^2) \operatorname{Erfc}(y)\} - 2\nu_0 k_\theta^2 \zeta \exp(\zeta^2) \operatorname{Erfc}(\zeta) \\ & \bar{\tau} k_\theta^2 \{1 + 1/2y^2\} - \bar{\nu} k_\theta^4 \{1 - 1/y^2 + 3/4y^4\} \end{aligned} \quad (3)$$

where $y = \alpha^2 k_\theta^2$, $\zeta = \alpha^2 (\bar{\tau}/\bar{\chi} + k_\theta^2)$, and $\operatorname{Erfc}(y) = \int_r^\infty \exp(-v^2) dv$. [Resistivity contribution is small] if $\bar{\eta} c \bar{\chi} k_\theta^2$. This condition is satisfied, as shown a posteriori, and η is neglected.] In order to obtain the physics insight, we obtain the analytic expression. In the following, we assume that $\bar{\chi} \gg \bar{\nu}$, since the electrostatic ExB transport is studied. (It is straightforward to study the general case of arbitrary ratio of $\bar{\chi}/\bar{\nu}$, but this does not change the result qualitatively.)

In the large αk_θ limit, the asymptotic limit of the function Erfc is used. Taking the leading term in αk_θ (note $\bar{\chi} \gg \bar{\nu} \gg \bar{\tau} \alpha^2$), the

eigenvalue equation $R(u) = \partial R(u) / \partial \alpha = 0$ gives the growth rate and the radial extent α of the fast interchange mode²²⁾ as $\bar{r} = \sqrt{D_0} / (\alpha k_\theta)^2$ and $\alpha^2 = \sqrt{\lambda(\bar{r} + 2\bar{v}k_\theta^2)} / s$. For the small αk_θ limit, the Taylor expansion of R is used. The first order term is written as $\bar{r} = (\sqrt{8/5}) \alpha k_\theta \sqrt{D_0}$. From these results, the largest growth rate is given for the poloidal mode number satisfying $k_\theta \alpha \sim 1$. For such mode, we have the estimate

$$\bar{r} \approx \sqrt{D_0} \quad (4-1)$$

$$\alpha^2 \approx s^{-1} \sqrt{3\lambda\sqrt{D_0}}. \quad (4-2)$$

Substituting Eq.(4) into Eq.(2), we have

$$\bar{x} = \frac{3}{s^2} \frac{\bar{\lambda}}{\bar{x}} D_0^{3/2} \quad (5)$$

We use the relation²³⁾ $\bar{\lambda} / \bar{x} \sim (\delta_S / a)^2$. Noting the normalization, the explicit form of \bar{x} is finally given as,

$$\bar{x} = F(\rho) \left[\frac{d\beta}{d\rho} \right]^{3/2} \delta_S^2 v_{\perp R}^{-1} \quad (6-1)$$

where $F(\rho)$ is the geometry-dependent numerical coefficient

$$F(\rho) = \frac{3}{s^2} \left\{ \frac{N}{2L} \frac{1}{\rho^2} \frac{d}{d\rho} (\mathcal{L}\rho^4) \right\}^{3/2}. \quad (6-2)$$

The ratio between the relative amplitudes of density and potential fluctuations, \tilde{n}/n and $e\tilde{\phi}/T$, can be derived from Eq.(4). Since the convective change dominates in n , we have the relation $\tilde{n}/n = (\omega_e/\tau)e\tilde{\phi}/T$, where ω_e is the drift frequency, $Tk_e\kappa/eB$ ($\kappa = |\nabla n/n|$ and we assume that $T_e = T_i$). Using the condition $k_e\alpha \gg 1$ and the expressions for τ and α , we have

$$\tilde{n}/n \approx [3.15D_0^{-1} \kappa R \beta(a)] e\tilde{\phi}/T. \quad (7)$$

This result gives that the density fluctuation is usually smaller than the potential fluctuation.

We study what is predicted from this model, comparing to the experimental results (I)-(IV).

Firstly, the dimensional dependence of χ is such that $[\chi] \propto [T]^{1.5}/RB^2$ and is independent of that of density. Equation(6) predicts χ of the experimental range (see Fig.1). Second, the point model analysis gives the energy transport scaling law as

$$\tau_E = A^{0.2} B^{0.8} n^{0.6} a^2 R P^{-0.6} \langle F \rangle^{0.4} \quad (8)$$

where A is the ion mass ratio, P is the heating power and $\langle F \rangle$ is the average of F near the boundary²⁴). The weak but positive dependence on the mass ratio is obtained. We also find that the improvement of the confinement by increasing the shear (s^{-2} term in F) is almost offset by the increment of the magnetic hill ($(N(\rho^4 \chi)')^{3/2} \rho^{-3}$ term in F). This result explains the fact that, from the comparison between different devices, τ_E seems to weakly

depend on the rotational transform/shear. $\langle F \rangle^{0.4}$ weakly depends on geometrical parameters. The predicted indices to B, n, a, R, and P, as a whole, agrees to the scaling law⁵⁾.

Third, the formula χ includes the radial dependence $(\rho'/n)^{3/2}$, not $T^{3/2}$, and predicts a large transport near edge. Since the pressure gradient is substantial near the edge (even though the pressure itself must be small) and $n(r)/n(0)$ is decreasing towards the edge, the anomalous transport can be large near edge. With this radial dependence and that of $F(\rho)$, χ is larger near the edge, as is shown in Fig. 1.

Forth, the heat pulse propagation time is faster than τ_E . For simplicity, we assume that $|\nabla T/T| \gg |\nabla n/n|$ in the region where heat pulse propagation is studied. Writing the heat flux $q = q_0 + \tilde{q}$ and $\nabla T = \nabla T_0 + \tilde{\nabla T}$, we have $\tilde{q} \approx 2.5 \chi_{eff} \tilde{\nabla T}$, where χ_{eff} is the ratio between q_0 and $|\nabla T_0|$ (i.e., χ in this article). From this relation, we see that the heat transport coefficient, which is derived from the heat pulse propagation, χ_{HP} , satisfies the relation

$$\chi_{HP} \approx 2.5 \chi_{eff}. \quad (9)$$

Fifth, the relative perturbation of density is smaller than that of the potential. We have $D_0 \approx 60 \kappa_p \beta(0)$ and $s \approx 4$ for Helio tron-E plasma⁴⁾, which gives $\tilde{n}/n \approx 2(\kappa/\kappa_p)^2 \beta(a)/\beta(0) |e\tilde{\phi}/T| (\kappa_p - |\nabla p/p|)$. This number is order of one tenth. Fluctuation measurements in high power heating experiments have shown that \tilde{n}/n is smaller than $e\phi/T$ ^{26, 27)}, confirming our model. This relation

also suggests that $(\tilde{n}/n)/(e\tilde{\phi}/T)$ increases as the pressure profile becomes broader.

These results are consistent with experiments including (I)-(IV).

In summary, we have developed a new model for the anomalous transport in the toroidal helical plasma with magnetic hill and magnetic shear and presented an analytic formula. The microscopic current diffusive interchange mode (λ mode) is analyzed by keeping the transport coefficients χ , ν , and λ . Mixing length estimate is used to derive the transport coefficient from the mode growth rate and structure. The mean field theory is employed so that the obtained transport coefficient is equated with the given value of χ . By this theoretical analysis we derived the formula of the anomalous transport coefficient. The mode analysis gives that the normalized density perturbation is usually smaller than the normalized potential fluctuation. The comparison with experimental result shows that the derived formula recovers the scaling law, radial shape of χ and difference between $\chi_{||p}$ and χ_{eff} , and that the relation between the amplitudes of the density and potential perturbations is also reproduced. This formula also explains the weak effects of ion mass and magnetic shear/rotational transform on χ_E .

We would like to note that Eq.(6) is derived apart from a numerical coefficient of the order of unity. The previous analyses on resistive interchange mode by two point renormalization method have shown the factor 5 enhancement over the mixing length

estimate^{11,13)}. Equation (6) will be changed by a factor like that, but the physical dependences of χ is not altered.

In this work the importance of the current diffusivity is shown. The resistivity is negligible if $\bar{\eta} < \bar{\lambda} k_{\theta}^2$ holds. This condition is rewritten by using the result of α ($\approx 1/k_{\theta}$) as $\bar{\chi}/\bar{\eta} > (a/\delta_s)^2 \bar{\eta} s^{-2} \sqrt{D_0}$, which is usually satisfied for experimental plasmas for which transport analysis is made.

We compare Eq. (6) to the formula derived by Onkawa for magnetic turbulence²⁸⁾. Compared to the Onkawa formula, $\chi \sim \delta_s^2 v_e/R$. Eq. (6) has an additional dependence on β . In our model, the current diffusion is proportional to δ_s^2 , so that the similar dependence on δ_s^2 is obtained, though the perturbation is assumed to be electrostatic.

Authors heartily wish to acknowledge useful discussions with Drs. H. Zushi, M. Yagi and K. Ichiguchi. They are grateful to Heliotron-E Group, CHS Group, and Dr. H. Sanuki for discussion on experimental data. Thanks are also due to discussion with Prof. A. J. Lichtenberg, through which the importance of the current diffusivity for the MHD mode was highlighted. This work is partly supported by the Grant-in-Aid for Scientific Research of Ministry of Education, Japan.

References


- [1] C. Gouldon, et al.: Plasma Physics and Controlled Nuclear Fusion Research (IAEA, Vienna, 1968) Vol.1, p847.
- [2] A. Mohri: J. Phys. Soc. Jpn. 28 (1970) 1549;
A. Mohri and M. Azumi: J. Phys. Soc. Jpn. 29 (1970) 1580.
- [3] K. Uo: Plasma Physics 13 (1971) 243.
- [4] K. Uo, et al.: Plasma Physics and Controlled Nuclear Fusion Research 1986 (Proc. 11th Int. Conf. Kyoto, 1986) Vol.2, IAEA, Vienna (1987) 355.
- [5] S. Sudo, et al.: Nucl. Fusion 30 (1990) 11.
- [6] M. Murakami, et al.: Plasma Physics and Controlled Nuclear Fusion Research 1990 (Proc. 13th Int. Conf. Washington, 1990) Vol.2, IAEA, Vienna (1987) 455.
- [7] H. Iguchi, et al.: in Controlled Fusion and Plasma Physics (Proc. 16th Eur. Conf. Venice, 1989) Vol.14B, European Physical Society, (1990) 451.
- [8] F. Sano, et al.: Nucl. Fusion 30 (1990) 81.
- [9] H. Zushi: private communications. (1991)
- [10] M. N. Rosenbluth and C. L. Longmire: Ann. Phys. (1957) 1120.
- [11] B. Carreras, L. Garcia, P. H. Diamond: Phys. Fluids 30 (1987) 1388.
- [12] K. C. Shaing, B. A. Carreras: Phys. Fluids 28 (1985) 2027.
- [13] H. Sugama, M. Wakatani: J. Phys. Soc. Jpn. 57 (1988) 2010.
- [14] M. Yagi, M. Wakatani, K. C. Shaing: J. Phys. Soc. Jpn. 57 (1988) 117.
- [15] B. B. Kadomtsev: in Plasma Turbulence (Academic Press, New

- York, 1965).
- [16] J. W. Connor and J. B. Taylor: Nucl. Fusion 17 (1977) 1047.
- [17] T. H. Dupree: Phys. Fluids 9 (1966) 1773, 15 (1972) 334.
- [18] M. Yagi: PhD Thesis, Kyoto University (1989).
- [19] K. Itoh, K. Ichiguchi, S.-I. Itoh: 'Beta Limit of Resistive Plasma in Torsatron/Heliotron Devices', Research Report NIFS-131 (National Inst. Fusion Science, 1992).
- [20] H. Strauss: Plasma Phys. 22 (1980) 733.
- [21] K. Ichiguchi, Y. Nakamura, M. Wakatani: Nucl. Fusion 29 (1991) 2073.
- [22] J. Greene, J. L. Johnson: Phys. Fluids 4 (1961) 875.
- [23] P. J. Kaw, E. J. Valeo, P. H. Rutherford: Phys. Rev. Lett. 43 (1979) 1398. See also A. J. Lichtenberg, et al.: 'Role of Stochasticity in Sawtooth Oscillation', Nucl. Fusion 31 (1992) in press.
- [24] The reason to choose the edge value of F is discussed in K. Itoh and S.-I. Itoh: Comments Plasma Phys. Cont. Fusion 14 (1991) 1.
- [25] M. Sato, et al.: in Controlled Fusion and Plasma Heating (Proc. 15th Eur. Conf. Dubrovnik, 1988) Vol. 13B, European Physical Society, (1988) 470.
- [26] H. Zushi, et al.: Nucl. Fusion 28 (1988) 433.
- [27] C. P. Ritz, et al. Plasma Physics and Controlled Nuclear Fusion Research 1990 (Proc. 13th Int. Conf. Washington, 1990) Vol. 2, IAEA, Vienna (1987) 589.
- [28] T. Ohkawa: Phys. Letters 67A (1978) 35.

Figure Caption

Fig.1 Example of the prediction of Eq.(6) in the model of Heliotron E ($\chi(\rho)=\chi_0+1.6\chi_1\rho^2$, $\varepsilon=0.1$, $B=2T$, $T(0)=500eV$, $n(0)=5\times 10^{19}m^{-3}$). Profiles are chosen such that $p_{eq}(\rho)=1+\Delta\rho^2$ and $n(\rho)/n(0)=(1+\Delta\rho^2)^{1/2}$ ($\Delta=0.05$). Thick dashed line indicates 4 times of the formula (6). Shaded region shows the range of experimental data, which is quoted from Ref.[18,25].

Fig. 1



Recent Issues of NIFS Series

- NIFS-90 H. Hojo and T.Hatori, *Radial Transport Induced by Rotating RF Fields and Breakdown of Intrinsic Ambipolarity in a Magnetic Mirror*; May 1991
- NIFS-91 M. Tanaka, S. Murakami, H. Takamaru and T.Sato, *Macroscale Implicit, Electromagnetic Particle Simulation of Inhomogeneous and Magnetized Plasmas in Multi-Dimensions*; May 1991
- NIFS-92 S. - I. Itoh, *H-mode Physics, -Experimental Observations and Model Theories-. Lecture Notes, Spring College on Plasma Physics, May 27 - June 21 1991 at International Centre for Theoretical Physics (IAEA UNESCO) Trieste, Italy* ; Jun. 1991
- NIFS-93 Y. Miura, K. Itoh, S. - I. Itoh, T. Takizuka, H. Tamai, T. Matsuda, N. Suzuki, M. Mori, H. Maeda and O. Kardaun, *Geometric Dependence of the Scaling Law on the Energy Confinement Time in H-mode Discharges*; Jun. 1991
- NIFS-94 H. Sanuki, K. Itoh, K. Ida and S. - I. Itoh, *On Radial Electric Field Structure in CHS Torsatron / Heliotron*; Jun. 1991
- NIFS-95 K. Itoh, H. Sanuki and S. - I. Itoh, *Influence of Fast Ion Loss on Radial Electric Field in Wendelstein VII-A Stellarator*; Jun. 1991
- NIFS-96 S. - I. Itoh, K. Itoh, A. Fukuyama, *ELMy-H mode as Limit Cycle and Chaotic Oscillations in Tokamak Plasmas*; Jun. 1991
- NIFS-97 K. Itoh, S. - I. Itoh, H. Sanuki, A. Fukuyama, *An H-mode-Like Bifurcation in Core Plasma of Stellarators*; Jun. 1991
- NIFS-98 H. Hojo, T. Watanabe, M. Inutake, M. Ichimura and S. Miyoshi, *Axial Pressure Profile Effects on Flute Interchange Stability in the Tandem Mirror GAMMA 10*; Jun. 1991
- NIFS-99 A. Usadi, A. Kageyama, K. Watanabe and T. Sato, *A Global Simulation of the Magnetosphere with a Long Tail : Southward and Northward IMF*; Jun. 1991
- NIFS-100 H. Hojo, T. Ogawa and M. Kono, *Fluid Description of Ponderomotive Force Compatible with the Kinetic One in a Warm Plasma* ; July 1991
- NIFS-101 H. Momota, A. Ishida, Y. Kohzaki, G. H. Miley, S. Ohi, M. Ohnishi, K. Yoshikawa, K. Sato, L. C. Steinhauer, Y. Tomita and M. Tuszewski *Conceptual Design of D-³He FRC Reactor "ARTEMIS"* ; July

- NIFS-102 N. Nakajima and M. Okamoto, *Rotations of Bulk Ions and Impurities in Non-Axisymmetric Toroidal Systems* ; July 1991
- NIFS-103 A. J. Lichtenberg, K. Itoh, S. - I. Itoh and A. Fukuyama, *The Role of Stochasticity in Sawtooth Oscillation* ; Aug. 1991
- NIFS-104 K. Yamazaki and T. Amano, *Plasma Transport Simulation Modeling for Helical Confinement Systems*; Aug. 1991
- NIFS-105 T. Sato, T. Hayashi, K. Watanabe, R. Horiuchi, M. Tanaka, N. Sawairi and K. Kusano, *Role of Compressibility on Driven Magnetic Reconnection* ; Aug. 1991
- NIFS-106 Qian Wen - Jia, Duan Yun - Bo, Wang Rong - Long and H. Narumi, *Electron Impact Excitation of Positive Ions - Partial Wave Approach in Coulomb - Eikonal Approximation* ; Sep. 1991
- NIFS-107 S. Murakami and T. Sato, *Macroscale Particle Simulation of Externally Driven Magnetic Reconnection*; Sep. 1991
- NIFS-108 Y. Ogawa, T. Amano, N. Nakajima, Y. Ohyabu, K. Yamazaki, S. P. Hirshman, W. I. van Rij and K. C. Shaing, *Neoclassical Transport Analysis in the Banana Regime on Large Helical Device (LHD) with the DKES Code*; Sep. 1991
- NIFS-109 Y. Kondoh, *Thought Analysis on Relaxation and General Principle to Find Relaxed State*; Sep. 1991
- NIFS-110 H. Yamada, K. Ida, H. Iguchi, K. Hanatani, S. Morita, O. Kaneko, H. C. Howe, S. P. Hirshman, D. K. Lee, H. Arimoto, M. Hosokawa, H. Idei, S. Kubo, K. Matsuoka, K. Nishimura, S. Okamura, Y. Takeiri, Y. Takita and C. Takahashi, *Shafranov Shift in Low-Aspect-Ratio Heliotron / Torsatron CHS* ; Sep 1991
- NIFS-111 R. Horiuchi, M. Uchida and T. Sato, *Simulation Study of Stepwise Relaxation in a Spheromak Plasma* ; Oct. 1991
- NIFS-112 M. Sasao, Y. Okabe, A. Fujisawa, H. Iguchi, J. Fujita, H. Yamaoka and M. Wada, *Development of Negative Heavy Ion Sources for Plasma Potential Measurement* ; Oct. 1991
- NIFS-113 S. Kawata and H. Nakashima, *Tritium Content of a DT Pellet in Inertial Confinement Fusion* ; Oct. 1991
- NIFS-114 M. Okamoto, N. Nakajima and H. Sugama, *Plasma Parameter Estimations for the Large Helical Device Based on the Gyro-*

Reduced Bohm Scaling ; Oct. 1991

- NIFS-115 Y. Okabe, *Study of Au⁺ Production in a Plasma-Sputter Type Negative Ion Source* ; Oct. 1991
- NIFS-116 M. Sakamoto, K. N. Sato, Y. Ogawa, K. Kawahata, S. Hirokura, S. Okajima, K. Adati, Y. Hamada, S. Hidekuma, K. Ida, Y. Kawasumi, M. Kojima, K. Masai, S. Morita, H. Takahashi, Y. Taniguchi, K. Toi and T. Tsuzuki, *Fast Cooling Phenomena with Ice Pellet Injection in the JIPP T-IIU Tokamak*; Oct. 1991
- NIFS-117 K. Itoh, H. Sanuki and S. -I. Itoh, *Fast Ion Loss and Radial Electric Field in Wendelstein VII-A Stellarator*; Oct. 1991
- NIFS-118 Y. Kondoh and Y. Hosaka, *Kernel Optimum Nearly-analytical Discretization (KOND) Method Applied to Parabolic Equations <<KOND-P Scheme>>*; Nov. 1991
- NIFS-119 T. Yabe and T. Ishikawa, *Two- and Three-Dimensional Simulation Code for Radiation-Hydrodynamics in ICF*; Nov. 1991
- NIFS-120 S. Kawata, M. Shiromoto and T. Teramoto, *Density-Carrying Particle Method for Fluid* ; Nov. 1991
- NIFS-121 T. Ishikawa, P. Y. Wang, K. Wakui and T. Yabe, *A Method for the High-speed Generation of Random Numbers with Arbitrary Distributions*; Nov. 1991
- NIFS-122 K. Yamazaki, H. Kaneko, Y. Taniguchi, O. Motojima and LHD Design Group, *Status of LHD Control System Design* ; Dec. 1991
- NIFS-123 Y. Kondoh, *Relaxed State of Energy in Incompressible Fluid and Incompressible MHD Fluid* ; Dec. 1991
- NIFS-124 K. Ida, S. Hidekuma, M. Kojima, Y. Miura, S. Tsuji, K. Hoshino, M. Mori, N. Suzuki, T. Yamauchi and JFT-2M Group, *Edge Poloidal Rotation Profiles of H-Mode Plasmas in the JFT-2M Tokamak* ; Dec. 1991
- NIFS-125 H. Sugama and M. Wakatani, *Statistical Analysis of Anomalous Transport in Resistive Interchange Turbulence* ;Dec. 1991
- NIFS-126 K. Narihara, *A Steady State Tokamak Operation by Use of Magnetic Monopoles* ; Dec. 1991
- NIFS-127 K. Itoh, S. -I. Itoh and A. Fukuyama, *Energy Transport in the Steady*

State Plasma Sustained by DC Helicity Current Drive ;Jan. 1992

- NIFS-128 Y. Hamada, Y. Kawasumi, K. Masai, H. Iguchi, A. Fujisawa, JIPP T-IIU Group and Y. Abe, *New High Voltage Parallel Plate Analyzer* ; Jan. 1992
- NIFS-129 K. Ida and T. Kato, *Line-Emission Cross Sections for the Charge-exchange Reaction between Fully Stripped Carbon and Atomic Hydrogen in Tokamak Plasma*; Jan. 1992
- NIFS-130 T. Hayashi, A. Takei and T. Sato, *Magnetic Surface Breaking in 3D MHD Equilibria of $l=2$ Heliotron* ; Jan. 1992
- NIFS-131 K. Itoh, K. Iguchi and S. -I. Itoh, *Beta Limit of Resistive Plasma in Toratron/Heliotron* ; Feb. 1992
- NIFS-132 K. Sato and F. Miyawaki, *Formation of Presheath and Current-Free Double Layer in a Two-Electron-Temperature Plasma* ; Feb. 1992
- NIFS-133 T. Maruyama and S. Kawata, *Superposed-Laser Electron Acceleration* Feb. 1992
- NIFS-134 Y. Miura, F. Okano, N. Suzuki, M. Mori, K. Hoshino, H. Maeda, T. Takizuka, JFT-2M Group, S.-I. Itoh and K. Itoh, *Rapid Change of Hydrogen Neutral Energy Distribution at L/H-Transition in JFT-2M H-mode* ; Feb. 1992
- NIFS-135 H. Ji, H. Toyama, A. Fujisawa, S. Shinohara and K. Miyamoto *Fluctuation and Edge Current Sustainment in a Reversed-Field-Pinch*; Feb. 1992
- NIFS-136 K. Sato and F. Miyawaki, *Heat Flow of a Two-Electron-Temperature Plasma through the Sheath in the Presence of Electron Emission*; Mar. 1992
- NIFS-137 T. Hayashi, U. Schwenn and E. Strumberger, *Field Line Diversion Properties of Finite β Helias Equilibria*; Mar. 1992
- NIFS-138 T. Yamagishi, *Kinetic Approach to Long Wave Length Modes in Rotating Plasmas*; Mar. 1992
- NIFS-139 K. Watanabe, N. Nakajima, M. Okamoto, Y. Nakamura and M. Wakatani *Three-dimensional MHD Equilibrium in the Presence of Bootstrap Current for Large Helical Device (LHD)*; Mar. 1992