






AIM

Association for Information and Image Management




1100 Wayne Avenue, Suite 1100
Silver Spring, Maryland 20910
301/587-8202



Centimeter



Inches



MANUFACTURED TO AIM STANDARDS
BY APPLIED IMAGE, INC.

1 of 1

LA-UR- 94-1907

Conf-9405188--1

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36.

TITLE: Progress in applying the FKK multistep reaction theory to intermediate-energy data evaluation

AUTHOR(S): Chadwick, M. B.*
Young, P. G., T-2

*University of California, Nuclear Data Group, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

SUBMITTED TO: For presentation at the Organization For Economic Co-Operation and Development Conference Specialist Meeting on Intermediate Energy Nuclear Data

Issy-les-Moulineaux, France
May 30, 1994 - June 1, 1994

By acceptance of this article, the publisher recognizes that the U. S. Government retains a non-exclusive, royalty-free license to publish or reproduce the published form this contribution, or to allow others to do so, for U. S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U. S. Department of Energy.

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

MASTER

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

875

Progress in applying the FKK multistep reaction theory to intermediate-energy data evaluation

M.B. Chadwick¹ and P.G. Young²

¹ *University of California, Nuclear Data Group,
Lawrence Livermore National Laboratory, Livermore, CA 94550, USA*

⁽²⁾ *University of California, Theoretical Division,
Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

Recent developments to the physics modeling in the FKK-GNASH code system are reviewed. We describe modifications to include a linking of multistep direct and multistep compound processes, which are important when the incident energy is less than about 30 MeV. A model for multiple preequilibrium emission is given, and compared with experimental measurements of proton reactions on ⁹⁰Zr at 160 MeV. We also give some preliminary observations concerning FKK calculations which use both normal and non-normal DWBA matrix elements.

We describe the application of the FKK-GNASH code to a range of nuclear data applications, including intermediate energy reactions of importance in the accelerator transmutation of waste, and fast neutron and proton cancer radiation treatment. We outline areas where further work is needed for the accurate modeling of nuclear reactions using the FKK theory.

I. INTRODUCTION

The FKK-GNASH code is a version of GNASH [1] which uses the Feshbach-Kerman-Koonin (FKK) theory [2] for preequilibrium emission of nucleons. The Hauser-Feshbach theory is used for equilibrium decay, in an open-ended sequence of reaction chains with full conservation of angular momentum. Transmission coefficients for particles are obtained from an optical model, and for gamma rays from a generalized Lorentzian giant resonance model.

The FKK-GNASH code was developed with a view towards improved modeling capabilities up to 200 MeV. It has been used in a number of analyses in the 14-26

MeV range [3–6], and was extended to analyze higher energy reactions up to the pion threshold [7–9]. In Ref. [10] we described recent developments to the GNASH and FKK-GNASH code systems. In considering reactions at higher energies it became clear that certain extensions to the FKK theory had to be made. In particular, inclusion of multiple preequilibrium processes was found to be essential when calculating emission spectra while simultaneously satisfying flux conservation [7, 8]. We had already observed [11] the need for including multiple preequilibrium in analyses of $(n, xn\gamma)$ excitation functions up to 200 MeV, and analyses of emission spectra provided further evidence for their importance. Other improvements that we have made in higher-energy modeling include a description of preequilibrium spin effects [4], and a theory linking the direct and compound multistep chains [3, 5]. In this paper we give an account of some of these developments.

According to the FKK theory, a nuclear reaction takes place in a series of stages corresponding to the interaction of the incident particle with nucleons in the target nucleus. At each stage, nucleons are excited to higher states and may be emitted; these are the preequilibrium reactions. The theory distinguishes between two types of interaction, the P -chain in which the projectile remains always in the continuum and the Q -chain where all the nucleons are bound after the initial interaction. The P -chain dominates at higher energies and gives cross sections peaked in the forward direction, while the Q -chain becomes important at low energies and gives cross sections symmetric about 90 degrees characteristic of compound nucleus emission. Emission from the P -chain is referred to as multistep direct (MSD) and emission from the Q -chain as multistep compound (MSC).

At incident energies above about 10-20 MeV the MSD process dominates, especially for the higher outgoing energies. The cross sections for such reactions have been extensively compared with the FKK theory, generally with good results. For a recent review see Ref. [12]. At lower energies both the MSC and MSD process contribute. The MSC formalism is given in the review of Bonetti *et al.* [13]. These analyses show the usefulness of the FKK theory in understanding the cross sections of nuclear reactions over a wide range of energies. The FKK-GNASH code includes both MSD and MSC calculations and calculates the whole emission spectrum in a consistent manner. Details of the theoretical formalism used can be found in Refs. [3, 8].

In Sec. II we describe recent developments in FKK analyses that can be important for modeling nuclear reactions up to 250 MeV: the development of a multistep scattering theory that links MSD and MSC processes; the inclusion of multiple preequilibrium emission; and investigations into the appropriate boundary conditions for matrix elements in multistep processes. In Sec. III we discuss the FKK-GNASH code's use in a number of applications. A summary of some of the outstanding questions that still need to be solved in FKK analyses is given in Sec. IV.

II. RECENT THEORY DEVELOPMENTS

A. Linking of MSD and MSC processes

Recent FKK analyses by Chadwick *et al.* [14, 3] and Marcinkowski *et al.* [15] have pointed to the importance of transitions from the MSD P -chain to the MSC Q -chain. These mechanisms were ignored in the original FKK paper, which assumed no crossover transitions after the initial separation into P and Q space. Evidence for $P \rightarrow Q$ transitions comes from two considerations: (1) Analyses of the partitioning between MSD and MSC emission, as described in Sec. IIA; and (2) Unitarity. The angular shape of the emission spectra in the preequilibrium emission regime determines the emission from the early MSC stages, and consequently the production of the early MSC stages. However, unitarity dictates that flux not lost to MSC or MSD preequilibrium emission must decay by compound nucleus emission. From such considerations it was concluded that a certain amount of reaction flux must bypass the initial Q -chain stages, entering the Q -chain after a number of collisions in P -space. It is not surprising that the Q -chain should be populated in this manner, since the probability of all particle excitations being bound will increase after more nucleon-nucleon collisions degrade the incident particle energy.

These works have postulated that the probability of flux entering a given Q -stage, with p particles and h holes and energy E , is given by the phase space ratio $\omega^B(p, h, E)/\omega(p, h, E)$, where the superscript “ B ” implies that only bound excitations are included. This prescription was checked by Chadwick and Young [3] for the initial flux entering the Q -chain, and was found to account for experimental data reasonably well. Also, independent investigations by Sato and Yoshida [17], on the imaginary optical potential in preequilibrium analyses partly confirmed this partitioning.

A full theoretical description of the coupling of MSD and MSC chains has recently been developed by Arbanas *et al.* [5], who removed the FKK assumptions which prevented $P \rightarrow Q$ transitions. They showed that in general there can be a number of P -space scatterings before, or after, scatterings in Q -space, and presented expressions for determining these processes. The attractive convolution structure of the MSD theory also appeared in the expressions for linked multistep scatterings in P and Q space.

The main physical consequence of these processes is that flux passing through preequilibrium stages can bypass the initial MSC Q -chain stages. Since it is predominantly these initial stages that give the high-energy MSC emission, this results in a reduced MSC component, and increased compound nucleus emission.

B. Multiple Preequilibrium Emission

Recent work [8] has shown that FKK analyses at incident energies above about 50 MeV should include multiple preequilibrium mechanisms, where more than one particle can be emitted from a preequilibrium stage. These processes were accounted

for using a model which is very easy to implement computationally, and which does not require any additional DWBA matrix elements to be determined. It was assumed that any particles left excited after primary preequilibrium can also immediately escape with transmission-coefficient probability.

It should be noted that the FKK-GNASH calculations [7] submitted to the NEA intermediate energy code intercomparison [18] used an earlier exciton-model algorithm for the multiple preequilibrium [1, 7]. Our new model, described in detail in Ref. [8], provides a more consistent way to describe these processes in FKK analyses.




Fig. 1. Primary and multiple MSD, and Hauser-Feshbach, compared with data for 160 MeV $^{90}\text{Zr}(p, xn)$.




Fig. 1. Primary and multiple MSD, and Hauser-Feshbach, compared with data for 160 MeV $^{90}\text{Zr}(p, xp)$.

In Figs. 1 and 2 we show comparisons between theoretical predictions and experimental measurements [25, 26] of angle-integrated neutron and proton emission spectra for proton reactions on zirconium at 160 MeV. The full curves represent calculations including all reaction mechanisms contributing to the inclusive spectra. The contribution from primary MSD is shown by the dashed line, multiple MSD by the dashed-dot line, and Hauser-Feshbach equilibrium decay by the dotted line. It is evident that the calculations account for the measurements well, for both neutron and proton emission. The importance of multiple preequilibrium emission can be clearly seen for all emission energies except the very highest, and at the lower emission energies this mechanism accounts for much of the data. Furthermore, it was shown in Ref. [8] that if the FKK theory is used to account for emission spectra without including multiple preequilibrium, it is likely that unitarity is violated.

In Ref. [11] we also demonstrated the importance of including multiple preequilibrium emission when describing excitation functions in $^{208}\text{Pb}(n, xn\gamma)$ reactions, for incident neutron energies up to 200 MeV, and for $x=1-9$. Including multiple preequilibrium greatly enhances the production of the residual nuclei $^{207,206}\text{Pb}$ (i.e. $x=2,3$), since the fast preequilibrium particles carry off much of the available incident energy. Likewise, production of residuals such as ^{200}Pb (i.e. $x=9$) is then strongly reduced since there is less energy available for many sequential compound-nucleus decays. Without including multiple preequilibrium emission the calculations would underpredict the data by orders of magnitude in some cases.

C. Non-normal DWBA?

Virtually all FKK analyses to date have applied the version of the MSD theory advocated by Feshbach [19], i.e. one uses normal DWBA matrix elements for the multistep transitions, allowing MSD to be obtained by convoluting 1-step cross sections. Yet there is little consensus among FKK aficionados as to whether this approach is correct. As pointed out by Kawai, and later in Koning and Akkerman's derivation [20], there are reasons to believe that non-normal DWBA matrix elements for the multistep transitions should be used. Physically, this accounts for cascading nucleons being located within the nucleus and thus should not be described by a wave coming in from infinity. In order to investigate this issue, we performed FKK calculations using a version of DWUCK4 modified by F.S. Dietrich to calculate non-normal DWBA. We find that the multistep contributions are greatly enhanced (an order of magnitude enhancement for the 2-step MSD is typical). We have the following (preliminary) observations concerning calculations based on these two approaches:-

- In the 10-30 MeV incident energy range, both theories describe data well. While the multistep contributions are bigger using non-normal DWBA, they are still small compared to 1-step MSD. But at 14 MeV, the magnitude of multistep contributions from the exciton model are similar to those from the non-normal DWBA theory [4]

- At higher energies we find non-DWBA multistep contributions which are too large, giving spectra that are of the wrong shape. However, it is possible that this is due to inadequacies in other assumptions, e.g. in the model used for single-particle level densities at high excitations
- The theory of Arbanas *et al.* [5] only yields significant $P \rightarrow Q$ flux when the non-normal DWBA theory is used. As discussed in Sec. IIA, there are reasons to believe such processes significant

In short, we have contradictory results which still need to be understood. In practice the normal DWBA theory advocated by Feshbach generally works very well and will continue to be used in analyses until the high-energy problems with the non-normal DWBA theory are resolved.

III. NUCLEAR DATA EVALUATION

Over the last couple of years there have been considerable developments in nuclear data evaluation work using the FKK theory, and here we briefly summarize this progress.

In order to use the FKK theory in applications, it is essential that the sensitivity of the calculations to different input parameters is understood. Watanabe [21] has shown that the value of the extracted residual interaction strength V_0 (the only free parameter in the theory) is rather sensitive to the choice of optical model, spin cut-off, and level density parameters, though the shape of the calculated spectra is not. Since it is desirable to find the systematical energy dependence of V_0 for predictive purposes, it is clear that a "standard" set of input parameters should be established for the analysis of different data sets. In this way the systematic behavior of V_0 can then be used with confidence to evaluate reactions where no measurements exist.

Accurate modeling of reactions at intermediate energy is needed for accelerator-base transmutation of waste (ATW) applications, where secondary neutrons from a proton-induced reaction are used to transmute long-lived radionuclides to stable or short-lived nuclei. In order to assess the modeling capabilities available, Blann *et al* established an international code intercomparison for intermediate energy nuclear data, organized by the Nuclear Energy Agency of the OECD [18]. Participants calculated neutron and proton emission spectra from proton-induced reactions on lead and zirconium, for a range of incident energies. The majority of the codes used semi-classical preequilibrium or intranuclear cascade models. However, FKK calculations were submitted by Chadwick and Young using the FKK-GNASH code system [7], and by Koning using the KAPSIES code system, and performed favorably in the intercomparison. There was evidence, though, that the FKK-GNASH calculations underpredicted backward-angle emission at the higher energies (see Fig. 3.), a problem

which still needs to be solved.




Fig. 3. FKK-GNASH calculation of the 80 MeV $^{90}\text{Zr}(p, xp)$ angular distributions at proton emission energies of 20, 40, and 60 MeV, compared with experimental measurements of Cowley *et al.* [24].

The FKK-GNASH code system has also been applied to evaluate neutron and proton induced reactions on biologically-important elements. The Lawrence Livermore National Laboratory Medical Applications Program is using Monte Carlo transport to calculate dose deposition in cancer radiation treatment. Certain types of tumor respond well to neutron or proton radiation, and to simulate dose deposition, libraries of nuclear cross sections are needed. A range of modeling codes have been applied to this problem (FKK-GNASH, GNASH, ALICE, and IDA), and their predictions compared with experimental measurements where available. As an example, in Figs. 4 and 5 we show the calculated inclusive proton and alpha spectrum from the $n+^{12}\text{C}$ reaction at 60 MeV, compared with measurements from UC Davis [27]. The calculations are seen to describe the data fairly well, though the highest proton energies are underpredicted since we have not included direct mechanisms to the discrete levels. In the case of alpha emission, much of the inclusive cross section comes from the 3α breakup of ^{12}C , which we treat in a sequential way using Hauser-Feshbach theory.

The total kerma (*kinetic energy released in matter*) in this reaction represents an integral check on our calculations. This quantity is crucial for obtaining an accurate prediction of dose deposition in neutron therapy. In Fig. 6 we see that the FKK-GNASH and GNASH calculations describe the data well, though the omission of preequilibrium clusters in ALICE leads to an underprediction.




Fig. 4. Various model calculations compared with data for the 60 MeV $^{12}\text{C}(n,p)$ reaction.




Fig. 5. Various model calculations compared with data for the 60 MeV $^{12}\text{C}(n,\alpha)$ reaction.




Fig. 6. Calculated total kerma factor for $n+^{12}\text{C}$ compared with data [28].

Finally, we mention that the FKK theory has also been used to determine radioactive isomer production in neutron-induced reactions of importance in fusion technology [4]. For this, particular attention was paid to spin effects in the theory. Certain reactions were selected and studied by an IAEA coordinated research programme, and the evaluated cross sections at 14 MeV and at lower energies will be taken into account during the engineering design of the ITER fusion reactor.

IV. FUTURE CHALLENGES

We have given an overview of some of the recent extensions and applications of the FKK-GNASH code. Here we itemize areas which still need to be addressed:

- When a 1-component model is used, different V_0 systematics result for (p, p') and (p, n) reactions [8]. Hopefully a 2-component theory which follows neutron and proton excitations would unify these systematics.
- There is still a need for a rigorous multiple preequilibrium theory which uses basic DWBA cross sections involving more than one final continuum particle.
- Present FKK-GNASH calculations at high energies underpredict back-angle emission. Hopefully this inadequacy can be resolved. Since Koning's FKK calculations [22] do not seem to suffer from this drawback, we shall investigate the use of collective (rather than microscopic) form factors in our calculations.
- The preequilibrium emission of composite particles such as alphas and deuterons has yet to be formulated in the FKK theory. At present we use the semiclassical approach of Kalbach [23].

- Further challenges in applying the FKK theory up to 200 MeV include: use of appropriate optical potentials and level densities at high energies; relativistic effects; and inclusion of quasifree scattering physics.

We wish to thank G. Arbanas, M. Blann, F. Dietrich, P. Hodgson, A. Kerman, A. Koning, Y. Watanabe, for useful discussions. This work was performed in part under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract no. W-7405-ENG-48.

REFERENCES

- [1] P.G. Young, E.D. Arthur and M.B. Chadwick, Los Alamos National Laboratory Report LA-12343-MS (1992).
- [2] H. Feshbach, A. Kerman, and S. Koonin, *Ann. Phys. (N.Y.)* **125**, 429 (1980).
Phys. Rev. C **47**, 2255 (1993).
- [3] M.B. Chadwick and P.G. Young, *Phys. Rev. C* **47** 2255 (1993).
- [4] M.B. Chadwick, P.G. Young, P. Oblozinsky, and A. Marcinkowski, *Phys. Rev. C* (in print, June 1994).
- [5] G. Arbanas, M.B. Chadwick, F.S. Dietrich, and A. Kerman, submitted to *Phys. Rev. C* (1994); Lawrence Livermore National Laboratory document UCRL-JC-117172 (1994).
- [6] Y. Watanabe *et al.*, in *Proceedings of the International Conference on Nuclear data for Science and Technology*, Gatlinburg, Tennessee, May 9-13 1994, ed. K. Dickens (to be published by the American Nuclear Society).
- [7] M.B. Chadwick and P.G. Young, Los Alamos National Laboratory document LA-UR-93-104 (1993); M.B. Chadwick and P.G. Young, in *Proceedings of the Workshop on Simulating Accelerator Radiation Environments*, 11-15 Jan. 1993, ed.? (World Scientific, in press).
- [8] M.B. Chadwick, P.G. Young, D.C. George, and Y. Watanabe, submitted to *Phys. Rev. C* (1994).
- [9] M.B. Chadwick, M. Blann, P.G. Young, G. Reffo, in *Proceedings of the International Conference on Nuclear data for Science and Technology*, Gatlinburg, Tennessee, May 9-13 1994, ed. K. Dickens (to be published by the American Nuclear Society).
- [10] P.G. Young and M.B. Chadwick, in *Proceedings of the International Conference on Nuclear data for Science and Technology*, Gatlinburg, Tennessee, May 9-13 1994, ed. K. Dickens (to be published by the American Nuclear Society).
- [11] H. Vonach, A. Pavlik, M.B. Chadwick, R.C. Haight, R.O. Nelson, and P.G. Young, submitted to *Physical Review C* (1994); P.G. Young, M.B. Chadwick and M. Bozoian, in *Proceedings of the Symposium on Nuclear Data Evaluation Methodology*, Brookhaven National Laboratory, Upton, New York, 1992, edited by C. Dunford (World Scientific, Singapore) p. 480.
- [12] P.E. Hodgson and M.B. Chadwick, in *Proceedings of the International Conference on Nuclear Data for Science and Technology*, Gatlinburg, Tennessee, May 9-13 1994, ed. K. Dickens (to be published by the American Nuclear Society).
- [13] R. Bonetti, M.B. Chadwick, P.E. Hodgson, B.V. Carlson, and M.S. Hussein, *Physics Reports* **202(4)**, 171 (1991).
- [14] M.B. Chadwick, D.Phil thesis, Oxford (1989).
- [15] A. Marcinkowski, J. Rapaport, R.W. Finlay, C. Brient, M. Herman and M.B.

- Chadwick, Nuclear Physics A561, 387 (1993)
- [16] H. Feshbach, Phys. Rev. C48, R2553 (1993); H. Feshbach, Ann. Phys. (N.Y.) 159, 150 (1985).
- [17] Sato and Yoshida, (1993).
- [18] M. Blann and P. Nagel, International Code Comparison (1994).
- [19] H. Feshbach, Phys. Rev. C48, R2553 (1993); H. Feshbach, Ann. Phys. (N.Y.) 159, 150 (1985).
- [20] A. Koning and J.M. Akkermans, Ann. Phys. 208, 216 (1991).
- [21] Y. Watanabe *et al.* to be published (1994).
- [22] A. Koning, O. Bersillon, and J.P. Delaroche, in *Proceedings of the International Conference on Nuclear data for Science and Technology*, Gatlinburg, Tennessee, May 9-13 1994, ed. K. Dickens (to be published by the American Nuclear Society).
- [23] C. Kalbach, Z. Phys. ?
- [24] A.A. Cowley, A. van Kent, J.J. Lawrie, S.V. Fortsch, D.M. Whittal, J.V. Pilcher, F.D. Smit, W.A. Richter, R. Lindsay, I.J. van Heerden, R. Bonetti, and P.E. Hodgson, Phys. Rev. C43, 678 (1991).
- [25] W. Scobel, M. Trabandt, M. Blann, B.A. Pohl, B.R. Remington, R.C. Byrd, C.C. Foster, R. Bonetti, C. Chiesa, and S.M. Grimes, Phys. Rev. C41, 2010 (1990).
- [26] W.A. Richter, A.A. Cowley, G.C. Hillhouse, J.A. Stander, J.W. Koen, S.W. Steyn, R. Lindsay, R.E. Julies, J.J. Lawrie, J.V. Pilcher, and P.E. Hodgson. Phys. Rev. C 49, 1001 (1994).
- [27] T.S. Subramanian, J.L. Romero, F.P. Brady, J.W. Watson, D.H. Fitzgerald, R. Garrett, G.A. Needham, J.L. Ullmann, C.I. Zanelli, D.J. Brenner, and R.E. Prael, Phys. Rev. C 28, 521 (1983).
- [28] R.M. White, J.J. Broerse, P.M. DeLuca Jr., G. Dietze, R.C. Haight, K. Kawashima, H.G. Menzel, N. Olsson, and A. Wambersie, Rad. Prot. Dos. 44, 11 (1992).

**DATE
FILMED**

8/29/94

END

