



DEMANDE D'AUTORISATION N° (AN) 1-5 (3)
EN VUE D'UNE PUBLICATION OU D'UNE COMMUNICATION

Direction : DSM
 Centre : Saclay
 Réf. émetteur : SPAM 94/362

34002545
 NTG n° 316
 J

Titre original du document : Propagation of intense laser pulses in an underdense plasma

Titre traduit en anglais :

A13573 IN. FN 910 2 71

Titre traduit en français : Propagation d'une impulsion laser intense dans un plasma sous dense

AUTEURS	AFFILIATION ¹	DEPT./SERV./SECT.	VISA (d'un des auteurs)	DATE
P. MONOT	CEA	DRECAM / SPAM	<i>100972</i> <i>J. Auguste</i> <i>1251</i>	20/07/94
T. AUGUSTE	CEA	DRECAM / SPAM		
P. GIBBON	CEA	DRECAM / SPAM		
F. JAKOBER	CEA	DRECAM / SPAM		
G. MAINFRAY	CEA	DRECAM / SPAM		
J.L. MIQUEL	CEA	DAM/DLPP/EPL/IL		
M. LOUIS-JACQUET	CEA	DAM/DLPP/EPL/IL		

Nature du document² :

PÉRIODIQUE
 CONF/CONGRÈS
 POSTER
 RAPPORT
 THÈSE
 COURS
 MÉMOIRE DE STAGE
 Chapitre d'OUVRAGE
 Pièces jointes : RÉSUMÉ
 TEXTE

CONGRES
CONFERENCE

Nom : High field interactions and shortwavelength generation
 Ville : ST MALO Pays : FRANCE Date : du 21 AOUT au 25 AOUT 1994

Organisateur :

PERIODIQUE

Titre :
 Comité de lecture : *ARRIVEE*

OUVRAGE

Titre :
 Éditeur : *14. MAR. 1995*

THESE
MEMOIRE DE STAGE
COURS

Université / Établissement d'enseignement : *S B D S / S P P I*

DOMAINES :

LANGUE : Anglais

N° EPAC : 3100

SUPPORT : Disquette Papier

MOTS-CLES :

Les visas portés ci-dessous attestent que la qualité scientifique et technique de la publication proposée a été vérifiée et que la présente publication ne divulgue pas d'information brevetable, commercialement utilisable ou classée.

	SIGLE	NOM	DATE	VISA	OBSERVATIONS	REF
CHEF DE SERVICE	SPAM	F. GOUNAND	20/07/94	<i>Tu</i> <i>[Signature]</i>	<i>Avis favorable</i>	
CHEF DE DEPARTEMENT	DRECAM	P. BERGE	-3-	<i>[Signature]</i>		

Date limite d'envoi du résumé

Date limite d'envoi du texte :

Date limite d'envoi du poster

Destinataires:

Les correspondants publication des départements se chargent de transmettre à l'INSTN/MIST/CIRST (Saclay) copies des demandes d'autorisation de publication, du résumé et du texte définitif.

¹ Entité d'appartenance de l'auteur. Ex. : CEA, CNRS, INSERM ...
² Cocher la case correspondante

ARRIVEE - CIRST

11 AOUT 94 004312

Circ. : Cit. :

VOL 2 N° 05

Propagation of intense laser pulses in an underdense plasma.

P. Monot, T. Auguste, P. Gibbon, F. Jakober and G. Mainfray

C.E.A. Saclay, Service des Photons, Atomes et Molécules, Bât 522, 91191 Gif-sur Yvette cedex, France. Tel: 33-1-69-08-91-49. Fax: 33-1-69-08-90-63. E-mail: monot@penduick.saclay.cea.fr.

J. L. Miquel, M. Louis-Jacquet.

C.E.A.-C.E.L.-V., 94195 Villeneuve Saint-Georges cedex, France

The new generation of short duration lasers provides pulses in excess of the terawatt level, that can be focused up to 10^{18} W/cm² [1]. For such an intensity, the quiver motion of a free electron becomes relativistic and numerous new physical effects are expected, such as harmonic generation [2], particle acceleration [3] and relativistic self-focusing [4,5]. In order to observe these effects resulting from laser-electron interaction, a high electron density (N_e) is required. In fact, with regard to the small laser-electron interaction cross-section, a large number of electrons is needed for any significant field emission. Furthermore, using a high density, a collective response of electrons is driven that induces intense longitudinal fields required to accelerate particles. A significant change of the refractive index should also occur that will influence beam propagation if the electron density is large enough.

Relativistic self-focusing results from an increase of electron mass ($m = \gamma m_0$) induced by the motion in the intense laser field. The refractive index n of the medium, $n = \sqrt{1 - \frac{N_e}{\gamma N_c}}$, has then a maximum on the beam axis and the effect is similar to the one produced by a converging lens. Calculations show that a critical power is required in order to overcome classical diffraction ($P_c = 2 \times 10^{10} N_c / N_e$) [5-7]. We present here experimental study of propagation for laser powers close to this critical power.

The laser beam is focused into a vacuum chamber onto a 3-mm long, pulsed hydrogen jet ($N_e \approx 10^{19}$ cm⁻³). At the output of the chamber the laser beam is collimated onto a CCD camera. A pair of lenses image the perpendicular Thomson emission onto a second camera.

Far field patterns are shown in fig. 1 for a laser power of 0.5 TW, which is under the critical power required to self-focus (2 TW).




Fig. 1. Laser beam profile at the output of the experimental chamber. Solid line: beam profile without interaction. Dashed line: The beam is focused into the hydrogen gas jet. The beam radius is significantly reduced after propagation in hydrogen for laser power (0.5 TW) below the critical power required for relativistic self focusing (2 TW).

The beam radius is reduced by a factor of three in the presence of hydrogen. This result shows clearly that even below the critical power required for self focusing a significant reduction of the output beam divergence is obtained.

Fig. 2 shows the evolution of the Thomson emission profile with the input laser power.




Fig. 2. Profile of the Thomson emission detected perpendicularly to the laser propagation direction. Solid line: $P = 1.5 \times P_c$. Dashed line: $P = 0.1 \times P_c$. The beam is guided when the laser power exceed the critical power.

A large increase of the emitting length is visible for power above the critical power. This length is limited by the jet size and corresponds to five times the Rayleigh length. This guiding of the pulse is in good agreement with numerical simulations and other experiments [8].

In conclusion, we have shown that an underdense plasma is able to significantly reduce the divergence of an intense laser pulse. The propagation mode demonstrated is in good agreement with theoretical predictions of relativistic self focusing.

- [1] P. Maine, D. Strickland, P. Badot, m. Pessot and G. Mourou, IEEE J. Quantum Electron. QE-24, 398(1988)
- [2] E. Sarachik and G. Schappert, Phys. Rev. D 1, 2738(1970).
- [3] P. Sprangle, E. Esarey, A. Ting and G. Joyce, Appl. Phys. Lett. 53, 2146(1988).
- [4] C. Max, J. Arons and A. Langdon, Phys. Rev. Lett. 33, 209(1974).
- [5] A. Borisov, A. Borovsky, V. Korobkin, A. Prokhorov, C. Rhodes and O. Shiryayev, Phys. Rev. Lett. 65, 1753(1990).
- [6] G-Z Sun, E. Ott, Y.C. Lee, and P. Guzdar, Phys. Fluids 30, 526(1987).
- [7] P. Sprangle, C. Tang and E. Esarey, IEEE PS-15, 145(1987).
- [8] R. W. Falcone, Int. Colloquium on X-ray Lasers, Schliersee Germany 1992, Inst. Phys. Conf. Ser. no. 125, section 4, p 213.

Propagation d'une impulsion laser intense dans un plasma sous-dense.

Nous présentons les résultats expérimentaux obtenus lors de l'étude de la propagation d'un faisceau laser intense dans un plasma sous-dense. Deux séries d'expériences, l'une menée à Saclay et l'autre à Limeil, montrent que la propagation du faisceau est altérée par des effets relativistes.