

BR 96 35080

UNICAMP - IMECC - RP -- 65/95

RELATÓRIO DE PESQUISA

I

ABOUT THE VELOCITY OPERATOR.
FOR SPINNING PARTICLES
IN QUANTUM MECHANICS

Giovanni Salsi, Erasmo Recami

and

Waldyr A. Rodrigues Jr.

Dezembro

RP 65/95

**INSTITUTO DE MATEMÁTICA
ESTATÍSTICA E CIÊNCIA DA COMPUTAÇÃO**



UNICAMP

UNIVERSIDADE ESTADUAL DE CAMPINAS

1995 12 10

ABSTRACT — Starting from the formal expressions of the hydrodynamical (or “local”) quantities employed in the applications of Clifford Algebras to quantum mechanics, we introduce —in terms of the ordinary tensorial framework— a new definition for the *field* of a generic quantity. By translating from Clifford into tensor algebra, we also propose a new (non-relativistic) velocity operator for a spin $\frac{1}{2}$ particle. This operator is the sum of the ordinary part \vec{p}/m describing the *mean* motion (the motion of the center-of-mass), and of a second part associated with the so-called *zitterbewegung*, which is the spin “internal” motion observed in the center-of-mass frame. This spin component of the velocity operator is non-zero not only in the *Pauli* theoretical framework, i.e. in presence of external magnetic fields and spin precession, but also in the *Schrödinger* case, when the wave-function is a spin eigenstate. In the latter case, one gets a *decomposition* of the velocity field for the *Madclung fluid* into two distinct parts: which constitutes the non-relativistic analogue of the Gordon decomposition for the Dirac current. We find furthermore that the *zitterbewegung* motion involves a velocity field which is solenoidal, and that the local angular velocity is parallel to the spin vector. In presence of a non-constant spin vector (Pauli case) we have, besides the component normal to spin present even in the Schrödinger theory, also a component of the local velocity which is parallel to the rotor of the spin vector.

IMECC – UNICAMP
Universidade Estadual de Campinas
CP 6065
13081-970 Campinas SP
Brasil

O conteúdo do presente Relatório de Pesquisa é de única responsabilidade do(s) autor(es).

Dezembro – 1995

ABOUT THE VELOCITY OPERATOR FOR SPINNING PARTICLES IN QUANTUM MECHANICS*

GIOVANNI SALESÌ

*Dipartimento di Fisica, Università Statale di Catania, 95129-Catania, Italy;
and INFN-Sezione di Catania, Catania, Italy: Salesi@ct.infn.it .*

and

ERASMO RECAMI[†] and WALDYR A. RODRIGUES JR.

*Dept. of Applied Math., State University at Campinas, Campinas, S.P.,
Brazil: Recami@turing.unicamp.br ; walrod@ime.unicamp.br*

Submitted 27 November 1995

Abstract. Starting from the formal expressions of the hydrodynamical (or “local”) quantities employed in the applications of Clifford Algebras to quantum mechanics, we introduce—in terms of the ordinary tensorial framework—a new definition for the *field* of a generic quantity. By translating from Clifford into tensor algebra, we also propose a new (non-relativistic) velocity operator for a spin $\frac{1}{2}$ particle. This operator is the sum of the ordinary part \vec{p}/m describing the mean motion (the motion of the center-of-mass), and of a second part associated with the so-called *zitterbewegung*, which is the spin “internal” motion observed in the center-of-mass frame. This spin component of the velocity operator is non-zero not only in the *Pauli* theoretical framework, i.e. in presence of external magnetic fields and spin precession, but also in the *Schrödinger* case, when the wave-function is a spin eigenstate. In the latter case, one gets a *decomposition* of the velocity field for the *Madelung fluid* into two distinct parts: which constitutes the non-relativistic analogue of the Gordon decomposition for the Dirac current. We find furthermore that the *zitterbewegung* motion involves a velocity field which is solenoidal, and that the local angular velocity is parallel to the spin vector. In presence of a non-constant spin vector (Pauli case) we have, besides the component normal to spin present even in the Schrödinger theory, also a component of the local velocity which is parallel to the rotor of the spin vector.

Key words: ICTE-1995, L^AT_EX, spin, *zitterbewegung*, velocity field, velocity operator, bilinear operators, Real Pauli Algebra, tensor algebra, Gordon decomposition, Madelung fluid, hydrodynamical picture, Dirac current, Pauli current, Schroedinger theory, Asim O. Barut.

1. HYDRODYNAMICAL OBSERVABLES IN QUANTUM THEORY

The *Multivector* or *Geometric* Algebras are essentially due to the work of great mathematicians of the nineteenth century as Hamilton (1805–1863), Grassmann (1809–1877) and mainly Clifford (1845–1879). More recently, starting from the sixties, some authors, and in particular Hestenes,[1-3] did systematically study various interesting physical applications of such algebras, and especially of the Real Dirac Algebra $\mathbb{R}_{1,3}$, often renamed *Space-Time Algebra* (STA).[4-6] Rather interesting

* Work partially supported by UNAM, by FAPESP, CNPq, and by INFN, MURST, CNR.

[†] On leave from Facoltà di Ingegneria, Università Statale di Bergamo, 24044-Dalmine (BG), Italy; and INFN-Sezione di Milano, Milan, Italy: Recami@mi.infn.it .

appear, in microphysics, the *applications* to space-time [O(3), Lorentz] transformations, to gauge [SU(2), SU(5), strong and electroweak isospin] transformations, to chiral [SU(2)_L] transformations, to the Maxwell equations, magnetic monopoles,[7] and so on. But the most rich and rigorous application is probably the formal and conceptual analysis of the geometrical, kinematical and hydrodynamical content of the Pauli and Dirac equations, performed by means of the Real Pauli and Real Dirac Algebras, respectively. We shall refer ourselves here to non-relativistic physics, and therefore adopt the Real Pauli Algebra, which is known to be *isomorphic to the ordinary tensorial algebra of the so-called Pauli matrices* [SU(2)]. In this paper, when ambiguities arise, the operators will be distinguished by a cap.

As well-known, in the usual *hydrodynamical picture of fluids*, every physical quantity depends not only on time, but also on the considered space point. In other words, every quantity is a *local* or *field* quantity:

$$G \equiv G(x) \quad x = (t; \mathbf{x}) . \quad (1)$$

In the Pauli Algebra the *local* value of G may be expressed in the following form:

$$G(x) = \rho^{-1} \langle \Psi \widehat{G} \Psi^\dagger \rangle_0 , \quad (2)$$

where $\langle \rangle_0$ indicates the *scalar part* of the *Clifford product* of the quantities appearing within the brackets. Let us translate eq.(2) into the ordinary tensorial language:

$$G(x) = \rho^{-1} \text{Re} [\psi^\dagger \widehat{G} \psi] . \quad (3)$$

It is easy to see, and remarkable, that this operator definition for $G(x)$ is equivalent to the *real part* of the so-called dual representation for *bilinear* operators, sometimes utilized in the literature.[8-10] In the ordinary approaches, such operators are commonly, even if *implicitly* only, employed for obtaining the probability densities of various quantities entering the Schrödinger, Klein-Gordon or Dirac wave-equations. In this sense, one can say that definition (3) agrees with the theoretical apparatus of ordinary wave-mechanics.

In connection with the first two mentioned wave-equations (and in the Dirac case, when confining ourselves to the translational-convective part of the well-known Gordon decomposition[11]), the energy density may be put into the following form:

$$\frac{i\hbar}{2} [\psi^*(\partial_t \psi) - (\partial_t \psi^*)\psi] \equiv \frac{1}{2} \psi^* i\hbar \overleftarrow{\partial}_t \psi , \quad (4)$$

as easily obtained from eq.(3) for the hamiltonian $\widehat{G} \equiv H \equiv i\hbar \partial_t$. Analogously, for the current density one can write:

$$-\frac{i\hbar}{2m} [\psi^*(\nabla \psi) - (\nabla \psi^*)\psi] \equiv \frac{1}{2m} \psi^* (-i\hbar \overleftarrow{\nabla}) \psi , \quad (5)$$

as required by eq.(3) if $G \equiv p/m$; $\widehat{G} \equiv -i\hbar \nabla/m$. Therefore, the use of the bilinear operators $\overleftarrow{\partial}_t$ and $\overleftarrow{\nabla}$ does allow us to write the above densities in the form expected for quantum-mechanical densities:[9] namely, in the form $\psi^* \widehat{X} \psi$.

Let us notice that, even if $\hat{G} \neq \hat{G}^\dagger$ [non-hermiticity], quantity $G(x)$ computed by means of eq.(3) will be always *real*. The only difference with respect to the case of hermitian operators is that the mean value $\langle G \rangle$ and the eigenvalues G_i will not be real; but this does *not* necessarily mean that G is unobservable. From the very definition of eigenstate in quantum mechanics, in fact, an eigenstate of \hat{G} from a "local point of view" is characterized by a function $G(x)$ *uniformly distributed (spatially homogeneous and constant in time)*: $G(x) = G_i$ for any x . Then, one can conclude[1] that necessary condition for the hermiticity of G , and the consequent existence of real eigenvalues G_i , is the possibility of creating and observing a *uniform distribution for quantity G* in correspondence with the chosen eigenstates. The inverse does not hold: it is possible to have real and locally uniform quantities not corresponding to hermitian operators. A noticeable example of this occurrence is given by the non-hermitian *velocity operator* proposed below. In spite of its non-hermiticity, we shall see for *plane waves* [$p = \text{constant}$] that its non-hermitian part will give no contribution, so that the velocity field will be real, locally uniform and equal to p/m .

2. THE NEW NON-RELATIVISTIC VELOCITY OPERATOR ENDOWED WITH ZITTERBEWEGUNG

In the framework of the Pauli geometric algebra, the local velocity is obtained from the usual operator \hat{p}/m , once it is "translated" into the new algebraic language. Thus we shall have, following the standard rules for that translation,

$$i\hbar \longrightarrow i2s, \quad (6)$$

where s represents the *spin vector* and i the Pauli algebra *pseudoscalar unity* (which corresponds to the matrix $\sigma_x \sigma_y \sigma_z$, so that $i^2 = -\mathbb{1}$). Therefore, we can write for the velocity field:

$$v(x) = -\frac{1}{m} \langle \psi \nabla i \sigma \psi^\dagger \rangle_0. \quad (7)$$

The "tensorial version" of this expression is the velocity operator: #1

$$\hat{v} = \frac{1}{m} \sigma (\hat{p} \cdot \sigma), \quad (8)$$

where σ indicates the usual 2×2 Pauli matrices (hereafter we shall work in the tensorial formalism). Due to the mathematical identity

$$\sigma (\mathbf{a} \cdot \sigma) \equiv \mathbf{a} + i \mathbf{a} \wedge \sigma, \quad (9)$$

\mathbf{a} being a generic 3-vector, we shall finally get:

$$\hat{v} = \frac{\hat{p}}{m} + \frac{i}{m} (\hat{p} \wedge \sigma) \equiv -\frac{i\hbar}{m} \nabla + \frac{\hbar}{m} (\nabla \wedge \sigma), \quad (10)$$

where \hat{p} and σ commute, making influential the order in which they appear in the product.

#1 Let us recall that, with regard to the *vectorial* basis $\vec{\sigma}^1, \vec{\sigma}^2, \vec{\sigma}^3$ of the Pauli multivector algebra, we have by definition: $\vec{p} \equiv \vec{\sigma}^i p_i$, indicating by p_i the i -th component of vector \vec{p} .

The above operator results to be composed by a hermitian part, $\widehat{\mathbf{p}}/m$, and by a non-hermitian part, $i(\widehat{\mathbf{p}} \wedge \boldsymbol{\sigma})/m$. The hermitian part reduces to the ordinary (but “incomplete”: see below) non-relativistic operator for wave-mechanics, usually written as $i[\widehat{H}, \widehat{\mathbf{x}}]/\hbar \equiv i[\widehat{\mathbf{p}}^2/2m, \widehat{\mathbf{x}}]/\hbar$. The non-hermitian part is strictly related to the so-called *zitterbewegung* (zbw), [12-14] which is the spin motion, or “internal” motion—since it is observed in the CMF,—expected to exist for spinning particles. Such a motion of an internal “constituent” \mathcal{Q} appears only for particles endowed with spin, [13, 14] and is to be added to the drift-translational, or “external”, motion of the CM, \mathbf{p}/m (which is the only one occurring for scalar particles). In the Dirac theory, indeed, the operators $\widehat{\mathbf{v}}$ and $\widehat{\mathbf{p}}$ are in general not parallel:

$$\widehat{\mathbf{v}} \neq \widehat{\mathbf{p}}/m. \quad (11)$$

Moreover, while $[\widehat{\mathbf{p}}, \widehat{H}] = 0$ so that \mathbf{p} is a conserved quantity, \mathbf{v} is *not* a constant of the motion: $[\widehat{\mathbf{v}}, \widehat{H}] \neq 0$ (quantity $\widehat{\mathbf{v}} \equiv \boldsymbol{\alpha} \equiv \boldsymbol{\gamma}^0 \boldsymbol{\gamma}$ being the usual vector matrix of the Dirac theory). Let us notice that in case of zbw it is highly convenient [12-14] to split the motion variables as follows (the dot meaning derivation with respect to time):

$$\widehat{\mathbf{x}} \equiv \widehat{\boldsymbol{\xi}} + \widehat{\mathbf{X}}; \quad \widehat{\dot{\mathbf{x}}} \equiv \widehat{\mathbf{v}} = \widehat{\mathbf{w}} + \widehat{\mathbf{V}}, \quad (12)$$

where $\widehat{\boldsymbol{\xi}}$ and $\widehat{\mathbf{w}} \equiv \widehat{\dot{\boldsymbol{\xi}}}$ describe the motion of the CM, whilst $\widehat{\mathbf{X}}$ and $\widehat{\mathbf{V}} \equiv \widehat{\dot{\mathbf{X}}}$ describe the zbw motion. From an electrodynamical point of view, the conserved electric current is associated with the trajectories of \mathcal{Q} (i.e., with $\widehat{\mathbf{x}}$), whilst the center of the particle Coulomb field—obtained [15] via a time average over the field produced by the quickly oscillating charge—coincides with the particle CM (i.e., with $\widehat{\mathbf{w}}$) and therefore, for free particles, with the geometrical center of the helical trajectory. #2

As a consequence, it is \mathcal{Q} which performs the *total motion*, while the CM undergoes the *mean motion* only. The resulting electron can be regarded as “extended-like” [13, 14] because of the existence in the CMF of an internal spin motion.

As required by eq.(11), one has to assume the existence of zbw also in the standard Dirac theory. In fact, the above decomposition for the total motion comes out in two well-known relativistic quantum-mechanical procedures: namely, in the *Gordon decomposition* of the Dirac current, and in the *decomposition of the Dirac velocity operator and Dirac position operator* proposed by Schrödinger in his pioneering works. [16]

The Gordon decomposition of the Dirac current reads (hereafter we shall choose units such that $c = 1$):

$$\bar{\psi} \boldsymbol{\gamma}^\mu \psi = \frac{1}{2m} [\bar{\psi} \widehat{\mathbf{p}}^\mu \psi - (\widehat{\mathbf{p}}^\mu \bar{\psi}) \psi] - \frac{i}{m} \widehat{\mathbf{p}}_\nu (\bar{\psi} S^{\mu\nu} \psi), \quad (13)$$

$\bar{\psi}$ being the “adjoint” spinor of ψ ; quantity $\widehat{\mathbf{p}}^\mu \equiv i\partial^\mu$ the 4-dimensional impulse operator; and $S^{\mu\nu} \equiv \frac{i}{4}(\boldsymbol{\gamma}^\mu \boldsymbol{\gamma}^\nu - \boldsymbol{\gamma}^\nu \boldsymbol{\gamma}^\mu)$ the spin-tensor operator. The ordinary

#2 As shown in refs.[14], solutions of the Dirac equations exist for which the zbw disappears while spin is still present! Such solutions correspond however to the limiting cases of plane waves ($\widehat{\mathbf{p}}$ -eigenfunctions). Moreover, even when the helical motion goes into a straight motion, the Frenet tetrad still rotates, denouncing an internal motion even in this limiting situation.

interpretation of eq.(13) is in total analogy with the decomposition given in eq.(10). The first term in the r.h.s. of eq.(13) results to be associated with the translational motion of the CM (the *scalar* part of the current, corresponding to the traditional Klein-Gordon current). By contrast, the second term in the r.h.s. results related to the existence of spin, and describes the zbw motion.

In the above quoted papers, Schrödinger started from the Heisenberg equation for the time evolution of the acceleration operator in Dirac theory

$$\hat{a} \equiv \frac{d\hat{v}}{dt} = \frac{i}{\hbar} [\hat{H}, \hat{v}] = \frac{i}{\hbar} 2(\hat{H}\hat{v} - \hat{p}); \quad (14)$$

where \hat{H} is equal as usual to $\hat{v} \cdot \hat{p} + \beta m$ (where $\hat{v} \equiv \alpha$). By integrating once this operator equation over time, he obtained:

$$\hat{v} = \hat{H}^{-1} \hat{p} + \hat{\eta}(0) e^{-2i\hat{H}t/\hbar} \quad (\hat{\eta} \equiv \hat{v} - \hat{H}^{-1} \hat{p}). \quad (15)$$

After a little algebra, we may get a more interesting form for the velocity decomposition:

$$\hat{v} = \hat{H}^{-1} \hat{p} - \frac{1}{2} i\hbar \hat{H}^{-1} \hat{a}. \quad (16)$$

By integrating a second time, Schrödinger ended up also with the spatial part of the decomposition:

$$\hat{x} \equiv \hat{\xi} + \hat{X}, \quad (17)$$

where we have

$$\hat{\xi} = \hat{r} + \hat{H}^{-1} \hat{p}t, \quad (18)$$

linked to the motion of the CM, and

$$\hat{X} = \frac{1}{2} i\hbar \hat{\eta} \hat{H}^{-1}, \quad (19)$$

linked to the zbw motion.

We can therefore consider decomposition (10) of our velocity operator as the *non-relativistic analogue of decomposition (16) of the relativistic velocity operator*. It is not at all surprising that (besides spin and the related intrinsic magnetic moment) also another “spin effect”, the zbw, does not vanish in the non-relativistic limit, i.e., for small velocities of the CM [$\beta \rightarrow 0$]. Therefore also the Schrödinger electron, being endowed with a zbw motion, does actually show its spinning nature, and is not a “scalar” particle (as often assumed)! As a matter of fact, when constructing atoms, we have necessarily to introduce “by hand” the Pauli exclusion principle; and in the Schrödinger equation the Planck constant \hbar implicitly denounces the presence of spin. In ref.[17] we have proved that the non-hermitian (zbw) part of our velocity field gives origin to the *quantum potential* of the Madelung fluid, as well as to the related *zero-point energy* of the Schrödinger theory.

3. ON THE VELOCITY FIELD OF THE MADELÜNG FLUID

Even if the above operator \hat{v} is not hermitian, the local velocity $v(x)$, as said above, will result to be always a real quantity. Let us see it by inserting eq.(10) into definition (3).

A spinning non-relativistic particle can be represented by means of a Pauli 2-components spinor:

$$\psi \equiv \sqrt{\rho} \Phi, \quad (20)$$

where, if we require $|\psi|^2 = \rho$, quantity Φ must obey the normalization constraint

$$\Phi^\dagger \Phi = 1. \quad (21)$$

Inserting the factorization (20) into definition (3), we have:

$$\rho s \equiv \text{Re} \left\{ \psi^\dagger \frac{\hbar}{2} \sigma \psi \right\} \equiv \rho \Phi^\dagger \frac{\hbar}{2} \sigma \Phi, \quad (22)$$

if G is the spin vector; and

$$\rho p \equiv \text{Re} \left\{ \psi^\dagger (-i\hbar \nabla) \psi \right\} = \frac{i\hbar\rho}{2m} [(\nabla\Phi^\dagger)\Phi - \Phi^\dagger\nabla\Phi], \quad (23)$$

if G is the impulse.

In the most general case (*Pauli* generalization of Schrödinger theory), when it is present an external potential $\mathbf{A} \neq 0$, we have to replace in the translational term of expression (10), as a "minimal prescription", the canonic impulse \hat{p} by the kinetic impulse $\hat{p} - e\mathbf{A}$; where e is the particle electric charge. Let us substitute in eq.(3) this "generalized" velocity operator for G , and factorization (20) for ψ : we finally get the following decomposition for the velocity field of the non-relativistic quantum fluid:

$$\mathbf{v} = \frac{\mathbf{p} - e\mathbf{A}}{m} + \frac{\nabla \wedge (\rho s)}{m\rho}. \quad (24)$$

This expression may be considered as the non-relativistic analogue of the Gordon decomposition (13) of the Dirac current.

We want to stress that decomposition (24), just now derived by means of definition (3), may be also obtained within standard wave-mechanics: and this will be a further test of the validity of our operator. It is sufficient, in fact, to take the familiar expression of the Pauli current (i.e., the non-relativistic limit of the Gordon decomposition[18])

$$\mathbf{j} \equiv \rho \mathbf{v} = \frac{i\hbar}{2m} [(\nabla\psi^\dagger)\psi - \psi^\dagger\nabla\psi] - \frac{e\mathbf{A}}{m}\psi^\dagger\psi + \frac{\hbar}{2m}\nabla \wedge (\psi^\dagger\sigma\psi), \quad (25)$$

and to insert into it factorization (20) in place of ψ , for obtaining the velocity distribution given by eq.(24).

Let us single out in the total velocity field the zbw-component:

$$\mathbf{V} \equiv \frac{\nabla \wedge (\rho s)}{m\rho}. \quad (26)$$

It is furtherly decomposable in two distinct parts:

$$A) \quad \mathbf{V}_1 \equiv (\nabla\rho \wedge s)/m\rho$$

due to the presence of the gradient of ρ , this term refers to local motions, in which

the constant density surfaces $[\nabla\rho = 0]$ do rotate around the spin axis; and it vanishes identically for the plane waves $[p = \text{constant}]$, for which $\nabla\rho = 0$;

$$\text{B) } \mathbf{V}_2 \equiv (\text{rot } \mathbf{s})/m$$

such a term does not depend on the density ρ and is different from 0 only in presence of external magnetic fields (the spin vector precedes, and the wave-functions are *not* spin eigenstates).

The *Schrödinger theory* is of course a particular case in the present Pauli framework, corresponding to $\mathbf{A} = 0$ (and then to zero magnetic field) and to a *uniform local spin vector* with no precession. The wave-function is then a *spin eigenstate* and may be factorized as the product of a "scalar" part $\sqrt{\rho}e^{i\varphi}$ and of a "spin" part χ (a 2-components spinor):

$$\psi \equiv \sqrt{\rho} e^{i\frac{\varphi}{\hbar}} \chi, \quad (27)$$

quantity χ being *constant* in space and time. Let us underline that, even if $\mathbf{s} = \chi^\dagger \frac{\hbar}{2} \boldsymbol{\sigma} \chi = \text{constant}$, in the *Schrödinger case the zbw does not vanish*—except for the unrealistic case of plane waves—, while the velocity \mathbf{V} reduces to

$$\mathbf{V} = \mathbf{V}_1 \equiv \frac{(\nabla\rho \wedge \mathbf{s})}{m\rho}. \quad (28)$$

All this does actually contribute to remove some difficulties remaining in the classical representation of the particle motion, and in the interpretation of the particle energies, for some stationary solutions of the Schrödinger equation. For instance, let us refer ourselves to the stationary states of a particle inside a box or of the harmonic oscillator, or the $l = 0$ eigenstates of the hydrogen atom. In all these cases the wave-function results to be *real* (φ is uniform and equal to a constant which for "global gauge invariance" may be assumed equal to zero), and therefore *the velocity obtained from standard quantum mechanics $p/m \equiv \nabla\varphi/m$ is zero everywhere, at any time.* As it was first remarked by Einstein and Perrin and by de Broglie,[19] such a result *seems to be really in contrast* from a classical point of view *with the non-vanishing of the energy eigenvalue for those stationary states.* But we now know, from our previous analysis, that p/m is the *mean* velocity, describing only the motion of the CM, whilst the zbw-component \mathbf{V} —which depends on the ρ -gradient, and not on the phase-gradient of ψ — does not identically vanish, thus implying an internal motion around the spin axis.

Let us finally analyse the velocity distribution (24) for $\mathbf{A} = 0$, and stress some interesting properties of its.

Since the rotor of a gradient is identically zero, we shall have $\text{rot } p = 0$; and since \mathbf{s} is furthermore uniform, it will also be $\nabla\mathbf{s} = 0$. As a consequence, by employing the known property of the double vectorial product

$$\mathbf{a} \wedge (\mathbf{b} \wedge \mathbf{c}) = (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c}, \quad (29)$$

we can show that the local rotational properties of the Madelung fluid are actually given by the following expression:

$$\text{rot } \mathbf{v} = \frac{1}{m} \left[\left(\frac{\nabla\rho}{\rho} \right)^2 - \frac{\Delta\rho}{\rho} \right] \mathbf{s}. \quad (30)$$

Moreover, the z bw field \mathbf{V} results to be *solenoidal* (and this happens also in the most general case of the Pauli fluid with a non-uniform \mathfrak{s}):

$$\operatorname{div} \mathbf{V} \equiv \operatorname{div} [\operatorname{rot}(\rho \mathfrak{s})] = 0. \quad (31)$$

The flux stream-lines will be *closed lines*, as in the magnetic field case: therefore, we expect *the zbw to be a limited, finite, periodical motion*. The explicit calculations performed by us for the known Barut–Zanghi model[13, 14] did really lead in the CMF of the electron to closed periodical motions. The total motion of the system will be helical around the p -direction. We can see eventually that the *local angular velocity ω is parallel to the spin vector \mathfrak{s}* :

$$\omega = \frac{1}{2} \operatorname{rot} v = \frac{1}{2m} \left[\left(\frac{\nabla \rho}{\rho} \right)^2 - \frac{\Delta \rho}{\rho} \right] \mathfrak{s}. \quad (32)$$

ACKNOWLEDGEMENTS

This work is dedicated to the memory of Asim O. Barut. The authors wish to acknowledge stimulating discussions with J. Keller, Z. Oziewicz and J. Vaz. For the kind cooperation, thanks are also due to G. Andronico, M. Baldo, A. Bonasera, M. Borromeli, A. Bugini, F. Catara, L. D'Amico, G. Dimartino, M. Di Toro, G. Giuffrida, C. Kiihl, L. Lo Monaco, G. Marchesini, R.L. Monaco, E.C. Oliveira, M. Pignanelli, G.M. Prosperi, R.M. Salesi, M. Sambataro, S. Sambataro, M. Scivoletto, R. Sgarlata, R. Turrisi, M.T. Vasconcelos, J.R. Zeni, and particularly I. Aragón, C. Dipietro and J.P. dos Santos. One of the authors (ER) wishes to thank Prof. J. Keller and all the Organizers for generous hospitality during this International Conference; and W.A Rodrigues and FAPESP for a very helpful research grant.

REFERENCES

1. Hestenes D.: "Space-time algebra" (Gordon & Breach; New York, 1966); "New foundations for classical mechanics" (Kluwer; Dordrecht, 1986); *Found. Phys.* **20** (1990) 1213; **23** (1993) 365; **15** (1985) 63; **12** (1981) 153; *Am. J. Phys.* **47** (1979) 399; **39** (1971) 1028; **39** (1971) 1013; *J. Math. Phys.* **14** (1973) 893; **16** (1975) 573; **16** (1975) 556; **8** (1979) 798; **8** (1967) 809; **8** (1967) 1046.
2. Hestenes D. and G. Sobczyk: "Clifford algebra to geometric calculus" (Reidel; Dordrecht, 1984).
3. Gurtler R. and D. Hestenes: *J. Math. Phys.* **16** (1975) 573.
4. Gueret Ph.: "Lectures at the Bari University" (Bari, 1989).
5. Lounesto P.: Clifford algebras, relativity and quantum mechanics, in "Gravitation: the space-time structure (Proceedings of SILARG VIII)", ed. by W.A. Rodrigues et al. (World Scientific; Singapore, 1994).
6. Benn I.M. and R.W. Tucker: "An introduction to spinors and geometry with applications in physics" (Hilger; Bristol, 1987); Crumeyrolle A.: "Orthogonal and symplectic Clifford algebras: Spinor structures" (Kluwer; Dordrecht, 1990).
7. Faria-Rosa M., E. Recami and W.A. Rodrigues: A satisfactory formalism for magnetic monopoles by Clifford algebras, *Phys. Lett.* **B173** (1986) 233; Maia A., E. Recami, W.A. Rodrigues and M.A.F. Rosa: Magnetic monopoles without strings: A geometrical interpretation, *J. Math. Phys.* **31** (1990) 502; Rodrigues W.A., E. Recami, A. Maia and M.A.F. Rosa: The classical problem of the charge and pole motion by Clifford algebra, *Phys. Lett.* **B220** (1989) 195.
8. Recami E., W.A. Rodrigues and P. Smrz: Some applications of non-hermitian operators in QM and QFT, *Nucl. Phys. B* **220** (1983) 1773.

9. Olkhovsky V.S. and E. Recami: About a Space and Time Operator in collision descriptions, *Lett. Nuovo Cim. (first series)* **4** (1970) 1165; Space-Time Shifts and cross-sections in collisions between relativistic wave-packets, *Nuovo Cimento* **A53** (1968) 610; Time operator in quantum mechanics, *Nuovo Cimento* **A22** (1974) 263; Recami E.: A time operator and the time-energy uncertainty relation, in "The uncertainty principle and foundations of quantum mechanics", ed. by W.C. Price and S.S. Chissick (J. Wiley; London, 1977), p.21; Descrizione spazio-temporale di processi elementari, *Rendic. Acc. Naz. Lincei, Cl. Sc.* **49** (1970) 77; An operator for Time, in "Recent development in relativistic quantum field theory and its application", ed. by W. Karwowski (WUW; Wroclaw, 1976), vol.2, p.251. Cf. also Kálnay A.J. et al.: *Phys. Rev.* **188** (1967) 1484; *D1* (1970) 1092; *D3* (1971) 2357, 2977.
10. Agodi A., M. Baldo and E. Recami: Approach to the theory of unstable states, *Ann. of Phys.* **77** (1973) 157. See also Agodi A., F. Catara and M. Di Toro: *Ann. of Phys.* **49** (1968) 445; Baldo M. and E. Recami: *Lett. Nuovo Cim.* **2** (1969) 643.
11. Bjorken J.D. and S.D. Drell: "Relativistic Quantum Mechanics", p.36 (McGraw-Hill; New York, 1964).
12. Barut A.O. and N. Zanghi: *Phys. Rev. Lett.* **52** (1984) 2009; Barut A.O. and A.J. Bracken: *Phys. Rev. D* **23** (1981) 2454; *D24* (1981) 3333; Barut A.O. and I.H. Duru: *Phys. Rev. Lett.* **53** (1984) 2355; Barut A.O.: *Phys. Lett. B* **237** (1990) 436; *Phys. Lett. A* **189** (1994) 277; Pavšič M.: *Phys. Lett. B* **205** (1988) 231; *B221* (1989) 264; *Class. Quant. Grav.* **7** (1990) L187; Barut A.O. and M. Pavšič: *Class. Quant. Grav.* **4** (1987) L131; *Phys. Lett. B* **216** (1989) 297; and refs. therein.
13. Pavšič M., E. Recami, W.A. Rodrigues, G.D. Maccarrone, F. Raciti and G. Salesi: Spin and electron structure, *Phys. Lett. B* **318** (1993) 481; Rodrigues W.A., J. Vaz, E. Recami and G. Salesi: Zitterbewegung and electron structure, *Phys. Lett. B* **318** (1993) 623; Salesi G. and E. Recami: Field theory of the extended-like electron, preprint IC/94/185 (ICTP; Trieste, 1994), to appear in the volume in memory of D.D. Ivanenko, ed. by P.I. Pronin & G.A. Sardanashevily (Moscow); About the kinematics of spinning particles, submitted for pub.; Recami E. and G. Salesi: Field theory of the spinning electron: The new, non-linear field equations, Preprint INFN/AE-94/13 (INFN; Frascati, 1994), submitted to *Adv. Appl. Cliff. Alg.*; About kinematics and hydrodynamics of spinning particles: Some simple considerations, in these Proceedings (preceding article); and refs. therein. Cf. also Corben H.C.: *Int. J. Theor. Phys.* **34** (1995) 19, and refs. therein.
14. G. Salesi and E. Recami: Field theory of the spinning electron: Internal motions, *Phys. Lett. A* **190** (1994) 137, E389; W.A. Rodrigues, J. Vaz and M. Pavšič: The Clifford bundle and the dynamics of the superparticle, to appear in a *Janach Center Publications Volume* (Polish Acad. Sci., Warsaw).
15. Vaz J. and W. A. Rodrigues: Zitterbewegung and the electromagnetic field of the electron, *Phys. Lett. B* **319** (1993) 203.
16. Schrödinger E.: *Sitzunger. Preuss. Akad. Wiss. Phys. Math. Kl.* **24** (1930) 418; **3** (1931) 1.
17. Salesi G.: Spin and Madelung fluid, submitted to *Phys. Rev. A*; see also Rylov Y.A.: Dirac equation in terms of hydrodynamical variables, *Adv. Appl. Cliff. Alg.* **5** (1995) 1.
18. Landau L.D. and E.M. Lifshitz: "Teoria quantistica relativistica" (MIR; Moscow, 1978), p.153.
19. de Broglie L.: *C. R. Acad. Sc. (Paris)* **183** (1926) 447; "Non-Linear Wave Mechanics" (Elsevier; Amsterdam, 1960); "Théorie Générale des Particules à Spin" (Gauthier-Villars; Paris, 1954).

RELATÓRIOS DE PESQUISA — 1995

- 01/95 Modelo de Regressão Weibull Dependente para Testes Acelerados em Riscos Competitivos - *Silvia Emiko Shimakura and Círcilia Yuko Wada.*
- 02/95 Semilinear Elliptic Equations with Exponential Nonlinearities - *João Marcos Bezerra do Ó.*
- 03/95 Some inequalities for immersed surfaces - *Valery Marenich and Irwen Valle Guadalupe.*
- 04/95 Geometria e Topologia de Fluxos de Anosov em 3-variedades - *Sérgio R. Fenley.*
- 05/95 Tangent Cones at Infinity Under Quadratic Sectional Curvature Decay - *G. Pacelli Bessa and Valery Marenich.*
- 06/95 The Holonomy in Open Manifolds of Nonnegative Curvature - *Valery Marenich.*
- 07/95 The Log Gamma Model and the Choice of a Parametric Lifetime Model - *Dione Maria Valença and Jonathan Biele.*
- 08/95 Shock Indicator for Adaptive Schemes for Conservation Laws - *Cristina Cunha and Sônia M. Gomes.*
- 09/95 Singularities of Reversible Vector Fields - *Marco Antonio Teixeira.*
- 10/95 On the Homology of Manifolds — *Ricardo N. Cruz.*
- 11/95 Submersions of Open Manifolds of Nonnegative Curvature — *Valery Marenich.*
- 12/95 Topological Gap-Phenomenon — *Valery Marenich.*
- 13/95 Cohomogeneity one Manifolds and Hypersurfaces of Revolution — *Antonio Carlos Asperti, Francesco Mercuri and Maria Helena Noronha.*
- 14/95 Low Codimensional Submanifolds of Euclidean Space with Nonnegative Isotropic Curvature - *Francesco Mercuri and Maria Helena Noronha.*
- 15/95 Creation of Particles in the Early Friedmann Universe — *A. A. Grib.*
- 16/95 EPR Paradox, Bell's Inequalities and Telepathic Communication — *A. A. Grib.*
- 17/95 O Grupo das Rotações com Dez Parâmetros — *G.C. Ducati, E. Capelas de Oliveira and G. Arcidiacono.*
- 18/95 Again On The Mass And Energy In General Relativity — *Yuri Bozhkov and Waldyr A. Rodrigues Jr.*
- 19/95 On The Question of Coincidence Between Spaces of Multilinear Absolutely Summing Mappings and Hilbert-Schmidt Mappings — *Mário C. Matos.*

- 20/95 Existence of Primitives for Analytic Mappings Between Banach Spaces — *Mário C. Matos.*
- 21/95 The Exact Bohr-Sommerfeld Quantization Rule — *Rodolfo L. Monaco and Waldyr A. Rodrigues, Jr.*
- 22/95 On the Cohomological Characterization of Real Free Pro-2-Groups — *Antonio José Engler.*
- 23/95 Eficiência Relativa Assintótica de Estimadores do Tamanho de uma População em Captura-Recaptura — *Luz Jeanet Bullon Camarena, Círcia Yuko Wada and Clarice Azevedo de Luna Freire.*
- 24/95 Non-symmetric Truncated Sequential and Repeated Significance Tests for the B-value — *Ronaldo Dias and Nancy Lopes Garcia.*
- 25/95 On the Stable Size Distribution in a Mathematical Model for Tumor Cells Growth Reproducing by Fission and the Cellular Resistance Problem — *H. F. Leckar and L. L. Vendite.*
- 26/95 Petrov-Galerkin and Collocation Methods Using Biorthogonal Spline Wavelets — *Sônia M. Gomes and Cristina Cunha.*
- 27/95 Chemotherapeutic Treatments: A Study of the Interplay Among Drug Resistance, Toxicity and Recuperation from Side Effects — *M. I. S. Costa and J. L. Boldrini.*
- 28/95 Conflicting Objectives in Chemotherapy with Drug Resistance — *M. I. S. Costa and J. L. Boldrini.*
- 29/95 A Refined Estimate of the Size of Concentration Sets for 2D Incompressible Inviscid Flow — *Helena J. Nussenzweig Lopes.*
- 30/95 Some Remarks on Non-Reconstructable Tournaments — *Cosimo Guido and José Carlos S. Kiihl.*
- 31/95 Second Order Connections and Stochastic Calculus — *Pedro José Catuogno.*
- 32/95 Tunneling Times and the “Hartman Effect” Revisited — *Vladislav S. Olkhovsky, Erasmo Recami, Fabio Raciti and Aleksandr K. Zaichenko.*
- 33/95 Semilinear Dirichlet Problems for the N -Laplacian in \mathbb{R}^N with Non-Linearities in the Critical Growth Range — *João Marcos B. do Ó.*
- 34/95 Patterns in Post-Feeding Larval Dispersion of Blowflies — *Jose Luiz Boldrini and Antonio Carlos Moretti.*
- 35/95 A Utilização das Simetrias na Integração das Equações Diferenciais — *Antonio C. Gilli Martins.*
- 36/95 Nonlinear Programming Algorithms using Trust Regions and Augmented Lagrangians with Nonmonotone Penalty Parameters — *Francisco A.M. Gomes, Maria Cristina Maciel and José Mario Martínez.*

- 37/95 Solving Nonlinear Systems of Equations by Means of Quasi-Newton Methods with a Nonmonotone Strategy — *Ana Friedlander, Marcia A. Gomes-Ruggiero, Daniel N. Kozakevich, José Mario Martínez and Sandra Augusta Santos.*
- 38/95 Convergence Properties of the Inverse Column-Updating Method — *Vera L. R. Lopes and José Mario Martínez.*
- 39/95 Quasi-Newton Methods with Derivatives — *José Mario Martínez.*
- 40/95 On Nonquadratic Hamiltonian Elliptic Systems — *D. G. de Figueiredo and C. A. Magalhães.*
- 41/95 From Perplex Numbers to Perplex Calculus — *Adilson Enio Motter and Márcio A. Faria Rosa.*
- 42/95 Magneto-Micropolar Fluid Motion: Existence of Weak Solutions — *Marko A. Rojas-Medar and José Luiz Boldrini.*
- 43/95 Magneto-Micropolar Fluid Motion: Global Existence of Strong Solutions — *Elva E. Ortega-Torres and Marko A. Rojas-Medar.*
- 44/95 Subluminal and Superluminal Solutions in Vacuum of the Maxwell Equations and the Massless Dirac Equation — *Waldyr A. Rodrigues, Jr. and Jayme Vaz, Jr.*
- 45/95 Electromagnetism and Quantum Mechanics as an Already Unified Theory — *Jayme Vaz, Jr. and Waldyr A. Rodrigues, Jr.*
- 46/95 Quasi-Normable Preduals of Spaces of Holomorphic Functions — *Domingo García and Jorge Mujica.*
- 47/95 On Fadeev's Operator Anomaly for the Gauss Law and the Generalization of Kac Moody Algebras — *Alex I. Shimabukuro and Márcio A. F. Rosa.*
- 48/95 Spaces of Holomorphic Mappings on Banach Spaces with a Schauder Basis — *Jorge Mujica.*
- 49/95 Teste de Hipóteses Não-paramétricos para Experimentos em Parcelas subdivididas (Split-Plot) — *Belmer García Negrillo.*
- 50/95 Regularization of Discontinuous Vector Fields on a 2-Dimensional Manifold — *J. Sotomayor and M.A. Teixeira.*
- 51/95 On the Convergence Rate of Semi-Galerkin Approximation for the Equations of Viscous Fluids in the Presence of Diffusion — *Pedro D. Damázio and Marko A. Rojas-Medar.*
- 52/95 Chemotherapeutic treatments with time increasing mutation rate to drug resistance — *J. L. Boldrini and M. I. S. Costa.*
- 53/95 Geodesics in Heisenberg groups — *Valery Marenich.*
- 54/95 Seismic Image Waves — *Peter Hubral, Jörg Schleicher and Martin Tygel.*
- 55/95 Compact Homogeneous Einstein Manifolds in Codimension Two — *Antonio Carlos Asperti, Helvecio Pereira de Castro and Maria Helena Noronha.*

- 56/95 **Witten's Monopole Equations as a Spinorial Representation of the Free Maxwell Equations** — *Jayme Vaz Jr.*
- 57/95 **True-Amplitude Migration to Zero Offset (MZO) by Diffraction Stack** — *Martin Tygel, Jörg Schleicher and Peter Hubral.*
- 58/95 **Restricted alternatives tests in bivariate exponential models with covariates** — *Cicilia Yuko Wada, Luiz Koodi Hotta and Luciana Jia Lin Hsieh.*
- 59/95 **Successive Projection Methods for the Solution of Overdetermined Nonlinear Systems** — *Maria A. Diniz-Ehrhardt and José Mario Martínez.*
- 60/95 **Abelian Subgroups of pro- p Galois Groups** — *Antonio José Engler and Jochen Koehnigsmann.*
- 61/95 **Tamanho da Amostra para Testes Não Paramétricos, com $C > 2$ Tratamentos** — *Belmer Garcúa Negrillo.*
- 62/95 **Teoria de Resposta de Item para Testes de Itens Dicotômicos** — *Clarice Azevedo de Luna Freire, Eugênia Maria Reginato Charnet, Márcia Milena Pivatto and Cherubino de Santana Colares.*
- 63/95 **Control Óptimo Fuzzy** — *Marko A. Rojas-Medar.*
- 64/95 **About Kinematics and Hydrodynamics of Spinning Particles: Some Simple Considerations** — *Erasmio Recami, Waldyr A. Rodrigues Jr. and Giovanni Salesi.*