




XJ9700039



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

E1-96-308

MEASUREMENT OF xF_3 , F_2 STRUCTURE
FUNCTIONS AND GROSS—LLEWELLYN SMITH
SUM RULE WITH IHEP—JINR
NEUTRINO DETECTOR

Submitted to «Physics Letters B»

VOL 28 № 11 

1996

L.S.Barabash, S.A.Baranov, Y.A.Batusov, S.A.Bunyatov, O.Y.Denisov,
M.Y.Kazarinov, O.L.Klimov, A.V.Krasnoperov, V.E.Kuznetsov,
V.V.Lyukov, Y.A.Nefedov, B.A.Popov, S.N.Prakhov, A.V.Sidorov,
V.I.Snyatkov, V.V.Tereshchenko, V.Y.Valuev

Joint Institute for Nuclear Research, Dubna, Russia¹

V.B.Anykeyev, A.A.Borisov, N.I.Bozhko, S.K.Chernishenko,
G.L.Čukin, R.M.Fachrutdinov, V.N.Goryachev, M.M.Kirsanov,
A.I.Kononov, A.S.Kozhin, V.I.Kravtsov, V.V.Lipajev, A.I.Mukhin,
S.A.Mukhin, Y.I.Salomatina, Y.M.Sapunov, K.E.Shestermanov,
A.A.Spiridonov, Y.M.Sviridov, V.V.Sytnik, V.L.Tumakov,
A.S.Vovenko

Institute for High Energy Physics, Protvino, Russia²

¹Supported by the Russian Foundation for Basic Research, Grant No 96-02-18562, by INTAS under Contract No 93-1180 and by the Russian Foundation for Basic Research, Grant No 96-02-18897.

²Supported by the Russian Foundation for Basic Research, Grant No 96-02-17608.

The data on deep-inelastic $\nu(\bar{\nu})$ -scattering in a wide region of momentum transfer provide a reliable basis for precise verification of QCD predictions [1]. In this paper the data on the $x F_3$ and F_2 structure functions (SF) are presented for the kinematic region of relatively small momentum transfer $0.55 < Q^2 < 4.0 \text{ GeV}^2$. The value of the Gross-Llewellyn Smith (GLS) sum rule [2] and $\alpha_S(M_Z)$ are evaluated.

The data samples were obtained from three independent exposures of the IHEP-JINR Neutrino detector [3] to the wide band neutrino and antineutrino beams [4] of the Serpukhov U70 accelerator. The exposure to the antineutrino beam ($\bar{\nu}_\mu$ -exposure) was performed at the proton beam energy $E_p = 70 \text{ GeV}$, whereas the two ν_μ -exposures were carried out one at $E_p = 70 \text{ GeV}$ and the other at $E_p = 67 \text{ GeV}$. The experimental set-up and selection criteria for CC events are discussed in [5]. We restricted the range of the measurements in W^2 to $W^2 > 1.7 \text{ GeV}^2$ and in $E_{\nu(\bar{\nu})}$ to $6 < E_{\nu(\bar{\nu})} < 28 \text{ GeV}$. The final number of events and the mean values of Q^2 , $\langle Q^2 \rangle$ for the three samples are given in Table 1.

Table 1: Summary of the exposures.

Beam	$\bar{\nu}_\mu$	ν_μ	ν_μ
$E_p \text{ (GeV)}$	70	70	67
Final statistics	741	2139	3848
$\langle Q^2 \rangle \text{ (GeV}^2\text{)}$	1.2	2.3	

The SF were measured as functions of x averaged over all Q^2 permissible for the energy range $6 < E_{\nu(\bar{\nu})} < 28 \text{ GeV}$. Events were binned in intervals of x , and values of $x F_3$ and F_2 were calculated in these intervals.

The number of ν_μ interactions, n^ν , and $\bar{\nu}_\mu$ interactions, $n^{\bar{\nu}}$, in a given bin of x is a linear combination of the 'average' values $\{F_2\}$ and $\{x F_3\}$ of the respective SF in this bin (we assume invariance under the charge conjugation):

$$n^{\bar{\nu}} = a^{\bar{\nu}} \cdot \{F_2\} - b^{\bar{\nu}} \cdot \{x F_3\}$$

$$n_{1,2}^\nu = a_{1,2}^\nu \cdot \{F_2\} + b_{1,2}^\nu \cdot \{xF_3\}.$$

The subscripts 1 and 2 correspond to the ν_μ -exposures at $E_p = 70 \text{ GeV}$ and $E_p = 67 \text{ GeV}$ respectively. The quantities $a^{\nu, \bar{\nu}}$ and $b^{\nu, \bar{\nu}}$ are integrals ('flux integrals') of products of the differential neutrino (antineutrino) flux $\phi^{\nu(\bar{\nu})}(E)$ and known factors depending on the scaling variables x, y as foreseen by the standard form of the differential cross-section for deep-inelastic $\nu_\mu (\bar{\nu}_\mu)$ -scattering off an isoscalar target

$$a^{\bar{\nu}} = N \frac{G^2 M}{\pi} \int \left(1 - y - \frac{Mxy}{2E} + \frac{1}{2(R+1)} y^2\right) E \phi^{\bar{\nu}}(E) dx dy dE,$$

$$b^{\bar{\nu}} = N \frac{G^2 M}{\pi} \int y \left(1 - \frac{y}{2}\right) E \phi^{\bar{\nu}}(E) dx dy dE$$

etc. Here N is the number of nucleons in the fiducial volume of the detector and the parameter $R = (F_2 - 2xF_1)/2xF_1$ measures the violation of Callan-Gross relation [6].

The number $n_{\nu(\bar{\nu})}$ of neutrino (antineutrino) interactions in a given x -bin was obtained from the measured number of neutrino (antineutrino) events in this bin by correcting the former one for acceptance, for smearing effects arising from Fermi motion and measurement uncertainties, for radiative effects (following the prescription given by De Rújula et al. [7]) and for target non-isoscalarity (assuming $d_v/u_v = 0.5$ [8]). To determine the appropriate correcting factors the Monte Carlo simulation of the experimental set up has been carried out using the program CATAS [9]. We used the Buras and Gaemers (BEBC) parametrization [10] for quark distributions. The charmed quark content of the nucleon was assumed to be zero. The kinematic suppression of $d \rightarrow c$ and $s \rightarrow c$ transitions was taken into account assuming slow rescaling [11] and the following charmed and strange quark masses: $m_c = 1.25 \text{ GeV}$, $m_s = 0.25 \text{ GeV}$. Fermi motion of nucleons was simulated according to [12]. The details of the Monte Carlo simulation of the known features of the experimental set up are discussed in [5] and [13].

The number of interactions in a given bin of x is subject to kinematic constraints imposed by the cuts in the muon momentum ($P_\mu > 1 \text{ GeV}/c$ [5]), in the neutrino energy ($6 < E_{\nu\bar{\nu}} < 28 \text{ GeV}$) and in the invariant mass square of the hadronic system ($W^2 > 1.7 \text{ GeV}^2$). These were taken into account in the calculation of the flux integrals by appropriate modification of the volume of integration.

The measured values of $x F_3$ and F_2 are presented in Table 2 and in Figure 1.

Table 2: The isoscalar structure functions F_2 and $x F_3$ obtained on the assumption of $R = 0$. The difference ΔF_2 between the values of F_2 obtained with $R = .1$ and those obtained with $R = 0$ is also presented. The bin edges are at $x = .0, .1, .2, .3, 4, .5, .65$.

x	$\langle Q^2 \rangle (\text{GeV}^2)$	F_2	<i>stat</i>	<i>syst</i>	ΔF_2	$x F_3$	<i>stat</i>	<i>syst</i>
.052	.55	1.169	.026	.047	.023	.445	.044	.062
.148	1.4	1.097	.026	.022	.022	.583	.044	.017
.248	2.2	.894	.023	.018	.019	.622	.038	.019
.346	2.9	.576	.016	.017	.013	.556	.027	.011
.447	3.4	.390	.014	.012	.009	.336	.023	.007
.563	4.0	.182	.008	.004	.004	.177	.012	.005

The systematic errors presented come from the uncertainties of the correcting factors due to the choice of some input quark distributions in the event simulation program CATAS. These systematic uncertainties were estimated by repeating the calculation of the SF using by turns the Field-Feynman [14] and GRV [15] quark distributions. Note that the systematic errors in Table 2 do not include the normalization error of 4% for F_2 and 11% for $x F_3$. These normalization errors originate from the uncertainties in the ν_μ and $\bar{\nu}_\mu$ flux determination [16].

With the values of $x F_3$, the GLS sum rule (the integral of F_3) has been estimated. Over the interval $0.02 < x < 0.65$ it was calculated by numerical integration of the measured values of $x F_3$ weighted by $1/x$. The contribution from the regions $0 < x < 0.02$ and $0.65 < x < 1$ was




Figure 1. The x -dependence of the isoscalar structure functions $F_2(x)$ and $x F_3(x)$. The statistical and systematic errors are added in quadrature, excluding the normalization error of 4% for F_2 and 11% for $x F_3$. The curve is fit of the form $x F_3(x) = A x^b (1-x)^c$. The best fit values of free parameters $A = 5.36 \pm 1.25 (stat)$, $b = 0.81 \pm 0.10 (stat)$, $c = 3.52 \pm 0.26 (stat)$ were obtained using for each x -bin the mean x of the bin as the actual x -point corresponding to the value of the structure function obtained.

evaluated by integrating over these regions the parametrization of $x F_3$ with the values of free parameters obtained from the fit to the data at $0.02 < x < 0.65$. Finally we obtained

$$\int_0^1 \frac{x F_3(x)}{x} dx = 2.13 \pm 0.38 (stat) \pm 0.26 (syst). \quad (1)$$

The systematic error quoted is the quadrature sum of ± 0.24 due to ν_μ and $\bar{\nu}_\mu$ flux uncertainties, and ± 0.09 due to the choice of some input quark distributions. In accordance with Table 1 we suppose that the measured value (1) of the GLS sum rule corresponds to the averaged value $\overline{Q^2} \sim 1.7 \text{ GeV}^2$.

The experimental data on the $x F_3$ were compared with the QCD prediction for Q^2 -evolution by the Jacobi polynomials method in the next-to-leading order QCD approximation [17, 18, 19, 20]. Making QCD analysis of the $x F_3$ SF, for the first step we do not discuss the problem of validity of application of perturbative QCD predictions for kinematical region of small Q^2 as well as the nuclear effects, heavy quarks thresholds effects and higher order QCD corrections.

In order to take into account the target mass corrections the Nachtmann moments [24] of F_3 and F_2 could be expanded in powers of M_{nucl}^2/Q^2 , and retaining only terms of the order M_{nucl}^2/Q^2 one could obtain:

$$M_{3(2)}(N, Q^2) = M_{3(2)}^{QCD}(N, Q^2) + \frac{N(N+1)}{N+2} \frac{M_{nucl}^2}{Q^2} M_{3(2)}^{QCD}(N+2, Q^2). \quad (2)$$

Here $M_3^{QCD}(N, Q^2)$ and $M_2^{QCD}(N, Q^2)$ are the Mellin moments of $x F_3$ and F_2 :

$$M_3(N, Q^2) = \int_0^1 dx x^{N-2} x F_3(x, Q^2), \quad (3)$$

$$M_2(N, Q^2) = \int_0^1 dx x^{N-2} F_2(x, Q^2), \quad N = 2, 3, \dots$$

The Q^2 -evolution of $M_3^{QCD}(N, Q^2)$ and $M_2^{QCD}(N, Q^2)$ is defined [22, 23] by QCD and is presented here for the nonsinglet case for

simplicity:

$$M_3^{QCD}(N, Q^2) = \left[\frac{\alpha_S(Q_0^2)}{\alpha_S(Q^2)} \right]^{d_N} H_N(Q_0^2, Q^2) M_3^{QCD}(N, Q_0^2), \quad (4)$$

$$d_N = \gamma^{(0),N} / 2\beta_0, \quad N = 2, 3, \dots$$

Here $\alpha_S(Q^2)$ is the strong interaction constant, $\gamma_N^{(0)NS}$ are the nonsinglet leading order anomalous dimensions, and the factor $H_N(Q_0^2, Q^2)$ contains all next-to-leading order QCD corrections [20, 21, 23].

The unknown coefficients $M_3(N, Q_0^2)$ in (4) could be parametrized as the Mellin moments of some function:

$$M_3^{QCD}(N, Q_0^2) = \int_0^1 dx x^{N-2} A x^b (1-x)^c (1+\gamma x), \quad (5)$$

where the constants A , b , c and γ should be determined from the fit to the data. Having at hand the moments (2)–(5) and following the method discussed in [17, 18, 19] we can write the $x F_3$ SF in the form:

$$x F_3^{N_{max}}(x, Q^2) = x^\alpha (1-x)^\beta \sum_{n=0}^{N_{max}} \Theta_n^{\alpha,\beta}(x) \sum_{j=0}^n c_j^{(n)}(\alpha, \beta) M_3^{QCD}(j+2, Q^2),$$

where $\Theta_n^{\alpha,\beta}(x)$ are the Jacobi polynomials and $c_j^{(n)}(\alpha, \beta)$ are the coefficients of the expansion of $\Theta_n^{\alpha,\beta}(x)$ in powers of x :

$$\Theta_n^{\alpha,\beta}(x) = \sum_{j=0}^n c_j^{(n)}(\alpha, \beta) x^j.$$

The accuracy of the SF approximation better than 10^{-3} is achieved for $N_{max} = 12$ in a wide region of the parameters α and β [18, 19].

Using nine Mellin moments for SF reconstruction and taking into account target mass corrections we have determined five free parameters A , b , c , γ and the QCD parameter $\Lambda_{\overline{MS}}$ (Table 3).

Table 3: The results of the NLO QCD fit to the xF_3 SF data for $f = 4$, $Q_0^2 = 3 \text{ GeV}^2$, $N_{MAX} = 12$, $\alpha = 0.7$, $\beta = 3.0$ with the corresponding statistical errors.

	$\langle\langle O \rangle\rangle = 0$	$\langle\langle O \rangle\rangle = 0.53$ [28]
χ^2	2.8	2.05
A	9.28 ± 1.73	0.90 ± 0.67
b	1.06 ± 0.11	0.31 ± 0.18
c	3.22 ± 0.31	3.64 ± 0.21
γ	-0.90 ± 0.21	9.53 ± 5.73
$\Lambda_{\overline{MS}}$ [MeV]	417 ± 51	358 ± 59
GLS sum rule	1.59	2.63
$\alpha_S(M_Z)$	0.123_{-4}^{+3}	0.120_{-4}^{+3}

For the Q^2 - dependence of the GLS sum rule we can write the following theoretical expression³ :

$$GLS(Q^2) = 3 \left[1 - \alpha_s(Q^2)/\pi + O(\alpha_s^2) - \frac{8}{27} \frac{\langle\langle O \rangle\rangle}{Q^2} \right] \quad (6)$$

where α_s is the coupling constant in the \overline{MS} scheme. The general structure of the high-twist (HT) term is known from [26]. The evaluation of this term was carried out in [27], $\langle\langle O \rangle\rangle = 0.33 \pm 0.16 \text{ GeV}^2$, and more recently in [28], $\langle\langle O \rangle\rangle = 0.53 \pm 0.04 \text{ GeV}^2$, using the same three-point function QCD sum rules technique. In order to estimate the uncertainties due to the HT contribution we included a phenomenological term⁴ $h(x) = -\frac{8}{27} \frac{\langle\langle O \rangle\rangle}{Q^2} x$ in the fitting procedure. The first moment of the function $h(x)$ gives some contribution to the GLS sum rule (6) in accordance with [28]. The results of the fit with $\langle\langle O \rangle\rangle = 0.53 \pm 0.04 \text{ GeV}^2$ are presented in Table 3. The value of $\alpha_S(M_Z)$ was calculated for both variants of the fit due to the so-called ‘matching relation’ [31]. We present the GLS sum rule values calculated through (5) with $N = 1$ and with the parameters from Table 3.

³See [25] for higher order QCD corrections to the GLS sum rule.

⁴This shape of $h(x)$ is in qualitative agreement with the theoretical prediction in [29] and experimental estimations in [30] for the x values from Table 2.

We repeated our fit taking into account both the statistical and systematic errors (from Table 2) added in quadrature. With HT from [28], the following estimations have been obtained: $\Lambda_{\overline{MS}} = 359 \pm 71 \text{ MeV}$, $GLS = 2.66$.

In the singlet case the moments of valence quarks, sea quarks and gluons were parametrized at Q_0^2 in the form:

$$M_q^{QCD}(N, Q_0^2) = \int_0^1 dx x^{N-2} [A_v x^{b_v} (1-x)^{c_v} + A_{sea} (1-x)^{c_{sea}}], \quad (7)$$

$$M_g^{QCD}(N, Q_0^2) = \int_0^1 dx x^{N-2} A_g (1-x)^{c_g}, \quad N = 2, 3, \dots$$

Keeping in mind the small number of experimental points we fix A_g from the momentum sum rule $M_q^{QCD}(2, Q^2) + M_g^{QCD}(2, Q^2) = 1$. Following the results [19] of the QCD analysis of F_2 at the momentum transfer $Q^2 = 5 \text{ GeV}^2$ we put $A_{sea} = 0.17$, $c_{sea} = 15$ and $c_g = 9$. The other parameters in (7) as well as Λ were determined from the fit of the data in the leading logarithm QCD approximation and were found to be $A_v = 2.49 \pm 0.311$, $b_v = 0.19 \pm 0.02$, $c_v = 2.80 \pm 0.05$, $\Lambda = (517 \pm 17) \text{ MeV}$ with $\chi^2 = 6.7$ for 6 experimental points and $Q_0^2 = 3 \text{ GeV}^2$. Only statistical errors were taken into account.

Several comments are in order:

- The values (1) and the results on the GLS sum rule in Table 3 are considerably smaller in comparison to the results of previous measurements. (See the summary on the GLS sum rule data in [32] and the latest 3-loop result [33].)
- The parameter $\Lambda_{\overline{MS}}$ is found to be about twice as large as the estimations in [20] and [34]. It is in qualitative agreement with the results of the NLO analysis [35] of the GLS sum rule in the \overline{MS} scheme: $\Lambda_{\overline{MS}}^{(4)} = 317 \pm 23(stat) \pm 99(syst) \pm 62(twist) \text{ MeV}$ with HT and $\Lambda_{\overline{MS}}^{(4)} = 435 \pm 20(stat) \pm 87(syst) \text{ MeV}$ without HT.
- The illustrative nature of the QCD fit to the data on F_2 should be pointed out. The matter is the absence of reliable theoretical

predictions for HT contribution to singlet SF. In spite of this, we obtained the momentum fraction carried by quarks in the nucleon, $M_q^{QCD}(2, Q^2) = 0.46$, to be in agreement with the previous measurements.

- The strong interaction constant at the point of Z boson mass is found to be higher than usual deep inelastic scattering results [36, 37].
- The consideration of the HT contribution decreases χ^2 and appreciably changes the parameters of the fit as well as the GLS sum rule value and $\alpha_S(M_Z)$. For a reliable QCD analysis one must calculate not only the GLS sum rule ($N = 1$) but also the higher SF moments ($N = 2, 3, \dots$). Using in addition a 3-loop QCD analysis one could expect to improve the estimation of $\alpha_S(M_Z)$.

In conclusion let us stress once more that the QCD analysis of SF is sensitive to the HT contribution and in the future it should take into account the nuclear effects, heavy quark threshold effects and higher order QCD corrections. We hope to improve the accuracy of our estimations by processing the additional data on deep-inelastic scattering obtained with the IHEP–JINR Neutrino detector in the wide band beams of ν_μ and $\bar{\nu}_\mu$.

References

- [1] G. Altarelli, in Proc. of the "QCD-20 Years Later" Conf. 9-13 June 1992, Aachen; World Scientific 1993, v. 1., p. 172; Ed. by P. M. Zerwas and H. A. Kastrup.
- [2] D. J. Gross, C. H. Llewellyn-Smith, Nucl. Phys. **B14** (1969) 337.
- [3] S. A. Bunyatov et al., Proc. Int. Conf. Neutrino 82 (Balatonfured, 1982), vol. 2, 249.
- [4] D. G. Baratov et al., Sov. J. of Technical Phys. **47** (1977) 991; A. P. Bugorsky et al., NIM **146** (1977) 367.
- [5] V. B. Anykeyev et al. in Proc. of the 3rd Tallinn Symp. on Neutrino Physics, Tartu 1995, p. 17.
- [6] C. G. Callan and D. J. Gross, Phys. Rev. Lett. **22** (1969) 156.
- [7] A. De Rújula et al., Nucl. Phys. **B154** (1979) 394.
- [8] H. Abramowicz et al., Z. Phys. **C17** (1983) 283.
- [9] A. S. Vovenko et al., NIM **212** (1983) 155.
- [10] A. J. Buras and K. J. F. Gaemers., Nucl. Phys. **B132** (1978) 249; BEBC WA59 Collaboration, K. Varnell et al., Z. Phys. **C36** (1987) 1.
- [11] H. Georgi and H. D. Politzer, Phys. Rev. **D14** (1976) 1829; J. Kaplan and F. Martin, Nucl. Phys. **B115** (1976) 333; R. Brock, Phys. Rev. Lett. **44** (1980) 1027.
- [12] A. Bodek and J. L. Ritchie, Phys. Rev. **D23** (1981) 1070.
- [13] J. Blumlein: in Materials of the VIII Workshop on the IHEP-JINR ν -Detector. Dubna, 1988, p.115.
- [14] R. Field and R. P. Feynman, Phys. Rev. **D15** (1977) 2590.

- [15] M. Glück, E. Reya, A.Vogt, *Z. Phys.* **C48** (1990) 471.
- [16] Y. M. Sapunov and Y. M. Sviridov: in Materials of the XVII Workshop on the IHEP-JINR ν -Detector, Dubna, 1995, p.81.
- [17] G. Parisi, N. Surlas, *Nucl. Phys.* **B151**, 421 (1979); I.S.Barker, C.B.Langensiepen and G.Shaw, *Nucl. Phys.* **B186**, 61 (1981).
- [18] V.G.Krivokhizhin et al., *Z. Phys.* **C36**, 51 (1987); BCDMS Col- lab., A. Benvenuti et al., *Phys. Lett.* **195B** (1987) 97; **223B** (1989) 490.
- [19] V.G.Krivokhizhin et al., *Z. Phys.* **C48**, 347 (1990).
- [20] A.L.Kataev and A.V.Sidorov, *Phys. Lett.* **B331**, 179 (1994).
- [21] A.L.Kataev et al., INP-0919/96, Moscow, 1996, hep-ph/9605367. Submitted to *Phys. Lett.* **B**.
- [22] F.J.Yndurain, Quantum Chromodynamics (An Introduction to the Theory of Quarks and Gluons).- Berlin, Springer-Verlag (1983), 117.
- [23] A.Buras, *Rev. Mod. Phys.* **52** (1980) 199.
- [24] O. Nachtmann, *Nucl. Phys.* **B 63** (1973) 237; H. Georgi, H.D. Politzer, *Phys. Rev.* **D 14** (1976) 1829; S. Wandzura, *Nucl. Phys.* **B 122** (1977) 412.
- [25] A.L. Kataev and V.V. Starshenko, *Mod. Phys. Lett.* **A10** (1995) 235.
- [26] E. V. Shuryak and A. I. Vainshtein, *Nucl. Phys.* **B199** (1982) 451.
- [27] V. M. Braun and A. V. Kolesnichenko, *Nucl. Phys.* **B283** (1987) 723.
- [28] G. G. Ross and R. G. Roberts, *Phys. Lett.* **B322**, 425 (1994).
- [29] M. Dasgupta and B.R. Webber, Cavendish-HEP-96/1 [hep-ph/9604388].

- [30] A.V. Sidorov, JINR E2-96-254 [hep-ph/9607275].
- [31] W. Marciano, *Phys. Rev.* **D29** (1984) 580.
- [32] A.L. Kataev and A.V. Sidorov, Proc. Quarks-94 Int. Seminar, Vladimir, May 1994; eds. D.Yu. Grigoriev, V.A. Matveev, V.A. Rubakov, D.T. Son and A.N. Tavkhelidze; World Scientific 1995, p. 288.
- [33] CCFR-NUTEV Collab., D. Harris et al., FERMILAB-Conf-95-114, March, 1995.
- [34] CCFR Collab. P.Z. Quintas et al, *Phys. Rev. Lett.* **71** (1993) 1307.
- [35] J. Chýla, A. L. Kataev , *Phys.Lett.* **B297** (1992) 385.
- [36] M. Shifman, TPI-MINN-95/22-T, UMN-TH-1416-95, August 1995.
- [37] S. Bethke, in Proc. QCD94 Conference, Montpellier, July 1994, Nucl. Phys. **B** (Proc. Suppl.) **39BC** (1995) 198.

Received by Publishing Department
on August 23, 1996.

Барабаш Л.С. и др.

E1-96-308

Измерение структурных функций xF_3 , F_2 и правила сумм

Гросса — Ллевеллин Смита на нейтринном детекторе

ИФВЭ—ОИЯИ

Изоскалярные структурные функции xF_3 и F_2 измерены как функции двух переменных x и Q^2 для энергий (анти)нейтрино 6-28 ГэВ. На основе измеренных значений xF_3 получена оценка для правила сумм Гросса — Ллевеллин Смита:

$$\int_0^1 F_3 dx = 2,13 \pm 0,38 \text{ (стат)} \pm 0,26 \text{ (сист).}$$
 Из КХД анализа данных по xF_3 получено значение масштабного параметра $\Lambda_{\overline{MS}} = 358 \pm 59$ МэВ. Значение константы сильного взаимодействия $\alpha_s(M_Z) = 0,120_{-4}^{+3}$ оказалось выше результатов большинства экспериментов по глубоконеупругому рассеянию лептонов.

Работа выполнена в Лаборатории ядерных проблем ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1996

Barabash L.S. et al.

E1-96-308

Measurement of xF_3 , F_2 Structure Functions

and Gross—Llewellyn Smith Sum Rule

with IHEP—JINR Neutrino Detector

The isoscalar structure functions xF_3 and F_2 are measured as functions of x averaged over all Q^2 permissible for the range 6 to 28 GeV of incident (anti)neutrino energy. With the measured values of xF_3 , the value of the Gross—Llewellyn Smith

sum rule is found to be
$$\int_0^1 F_3 dx = 2.13 \pm 0.38 \text{ (stat)} \pm 0.26 \text{ (syst).}$$
 The QCD analysis of xF_3 provides $\Lambda_{\overline{MS}} = 358 \pm 59$ MeV. The obtained value of the strong interaction constant $\alpha_s(M_Z) = 0.120_{-4}^{+3}$ is larger than most of the deep inelastic scattering results.

The investigation has been performed at the Laboratory of Nuclear Problems, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1996

Макет Т.Е.Попеко

Подписано в печать 19.09.96
Формат 60 × 90/16. Офсетная печать. Уч.-изд. листов 1,18
Тираж 450. Заказ 49373. Цена 1416 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области