



IE9700011

RPII - 96/4

# Radon in Dwellings - The National Radon Survey Cavan, Dublin, Louth, Monaghan and Wicklow

*J.T. Duffy  
G.M. Mackin  
S.G. Fennell  
J.S. Madden  
A.T. McGarry  
P.A. Colgan*

*October 1996*



**Radiological Protection Institute of Ireland**  
An Institiúid Éireannach um Chosaint Raideolaíoch

3 Clonskeagh Square,  
Clonskeagh Road, Dublin 14.  
Tel: +353 1 269 7766.  
Telex: 30610.  
Fax: +353 1 269 7437.  
E mail: [rpil@rpil.ie](mailto:rpil@rpil.ie)

## *Abstract*

*This report presents the first results of the National Radon Survey carried out by the Radiological Protection Institute of Ireland. The average radon concentrations for the houses measured in counties Cavan, Dublin, Louth, Monaghan and Wicklow ranged from 69 to 138 Bq/m<sup>3</sup> with individual values as high as 1000 Bq/m<sup>3</sup>. The measurement data were grouped on the basis of the 10 km grid squares of the Irish National Grid System and used to predict the percentage of dwellings in each grid square which exceeds the Reference Level of 200 Bq/m<sup>3</sup>. Grid squares where this percentage is predicted to be 10% or higher are designated High Radon Areas. The health effects of exposure to high radon levels are discussed and recommendations are made regarding both new and existing dwellings.*

## **CONTENTS**

- 1. Introduction**
- 2. Radon Sources and Health Effects**
- 3. Survey Methodology**
- 4. Results**
  - 4.1 County Surveys
  - 4.2 High Radon Areas
- 5. Doses and Risk Estimates**
- 6. Conclusions and Recommendations**
- 7. Acknowledgements**
- 8. References**

**Glossary of Terms**

**Figures**

## **1. Introduction**

It is accepted that long-term exposure to radon, a naturally occurring radioactive gas, can cause lung cancer. In a minority of homes and other buildings in Ireland, elevated radon concentrations occur which constitute a significant lung cancer risk for the occupants. When a building with a high radon level is identified, measures are available whereby the level can be substantially reduced.

In 1990, the Government adopted an annual average radon concentration of 200 becquerels per cubic metre ( $\text{Bq/m}^3$ ) as the national Reference Level above which remedial action to reduce the indoor radon level in a dwelling should be considered. Results of a population-weighted radon survey carried out by the Experimental Physics Department of University College Dublin (UCD) between 1985 and 1989 estimate that 4% of the national housing stock have radon concentrations above this Reference Level [McLaughlin and Wasiolek, 1988]. In particular, the UCD survey indicated that the probability of finding a house with elevated indoor radon levels was greatest in the western counties, although houses with elevated radon concentrations were identified in most counties in Ireland.

Following this survey, the Institute undertook follow-up surveys in some of the identified potentially high radon areas including areas in counties Mayo, Galway, Clare and Kerry. More intensive surveys were carried out in selected areas in Co. Galway and south Cork city. The results of these surveys are published in a previous report [Madden *et al.*, 1994].

To assist in identifying areas at greatest risk of high indoor radon levels, the Institute in 1992 initiated a national geographically-based radon survey. The results of this survey will allow the classification of the whole country on the basis of the predicted percentage of dwellings in each area which are above the Reference Level. Areas in which this value is 10% or greater will be designated High Radon Areas.

This report presents the first results of the National Survey and covers the counties of Cavan, Dublin, Louth, Monaghan and Wicklow. Details are given of the percentages of dwellings in these counties estimated to exceed the Reference Level and High Radon Areas are identified.

## **2. Radon Sources and Health Effects**

Radon is a naturally occurring radioactive gas. It is part of the radioactive decay series of uranium, which is present in variable amounts in all rocks and soils. Although radon itself has a relatively short half-life of 3.82 days, it is continually produced from its parent, radium-226, which has a half-life of 1600 years. Radon decays through a series of short lived decay products to stable lead.

Because it is an inert gas, radon can move freely through porous media such as rocks or soil, although only a small fraction of that produced in the ground rises to the surface and enters the atmosphere or seeps into buildings. Radon which surfaces in the open air is quickly diluted. Typical outdoor concentrations are between 4 and 15 Bq/m<sup>3</sup> [NCRP, 1988] and are relatively harmless.

A problem can arise, however, when radon from the ground enters an enclosed space such as a house. There it may reach unacceptably high concentrations as dispersion is restricted. Domestic water and gas supplies, in addition to building materials and infiltration of outdoor air, also contribute to the radon level in a dwelling, but in most cases these are considered minor sources relative to the ground on which the dwelling is constructed [Nero *et al.*, 1983].

The most significant mechanisms by which radon can enter a house are concentration-driven diffusion and pressure-driven flow from the ground beneath the house. The higher temperature indoors and the effect of wind on a dwelling combine to reduce the pressure indoors relative to that in the ground. The pressure difference is usually small, but sufficient to induce a flow of soil gas with radon into the dwelling if the ground is fairly permeable and an easy path for flow exists. Such paths include cracks in the floor or the walls, construction joints or gaps around pipes or cables. In most dwellings with high radon concentrations pressure-driven flow is recognised as the dominant mechanism of influx [Nazaroff *et al.*, 1988].

Two of the short-lived decay products of radon (polonium-218 and polonium-214) emit alpha particle radiation. It is these decay products which, following inhalation and deposition in the lung, deliver the radiation dose that is implicated in the induction of lung cancer [ICRP, 1977].

### **3. Survey Methodology**

The primary objective of the survey was to determine, in detail, the geographical distribution of radon levels in dwellings in Ireland. In order to obtain a good geographical spread the survey was based on small area units; in this case, the 10 km grid squares of the Irish National Grid System. Radon measurements were carried out in selected dwellings and the data were used to predict the percentage of dwellings in each grid square with radon concentrations above the Reference Level. Grid squares where the predicted percentage of dwellings with radon concentrations above 200 Bq/m<sup>3</sup> is 10% or greater are designated High Radon Areas.

It has been shown that the distribution of radon levels in dwellings approximates a log-normal distribution [CEC, 1987] and that such a log-normal approximation appears to hold whether a whole country, or a smaller area such as a 10 km grid square, is considered [NRPB, 1990]. Modelling the distribution of indoor radon concentrations therefore allows the percentage of dwellings exceeding a reference concentration to be estimated [Miles, 1994]. The statistical procedure used in this survey is described by Daly [1994] and is based on the approach of Liberman and Resnikoff [1955].

Although in a truly random survey, a valid prediction can be made based on a sample size of only three dwellings per grid square, a minimum sample size of five dwellings was chosen for this survey. Based on a typical response rate of 20% for previous surveys, initially about 50 householders per grid square were invited to participate. The householders' names and addresses were randomly selected from the Register of Electors by the Economic and Social Research Institute (ESRI). As the Register of Electors is not available on a grid square basis, the names and addresses were selected from the Register for each District Electoral Division (DED). There are, on average, six DED's per grid square and so a minimum of eight householders were selected from each DED.

The householders were then invited by letter to participate in the survey. When agreeing to participate the householders were asked to locate, as accurately as possible, the locations of their homes on the county map provided. This information was then used to assist in assigning the measurement for each dwelling to the correct grid square.

Each participant was issued with two radon detectors, one to be placed in the main living area and the second in an occupied bedroom. The detectors used were of the passive alpha track type and were left in place for a period of twelve months. On return to the Institute, the detectors were processed and the annual average radon concentration for each dwelling was calculated by averaging the results of the two measurements. The householders were individually notified of their measurement results and, where appropriate, recommendations were made regarding the necessity for remedial action.

The survey was carried out on a county by county basis and phased to match the available resources. In grid squares where there were less than five valid measurements, no predictions were made and these squares are to be re-surveyed. For grid squares which overlay the border between two or more counties, the percentage of dwellings above the Reference Level was predicted using all of the available data. In some cases, this prediction required the use of data for counties other than those specifically reported here.

## **4. Results**

### **4.1 County Surveys**

Radon measurements were completed in a total of 691 dwellings in counties Cavan, Dublin, Louth, Monaghan and Wicklow. A summary of the results of these measurements is presented in Table 1. Mean annual indoor radon concentrations for the dwellings measured in these counties ranged from 69 Bq/m<sup>3</sup> in Monaghan to 138 Bq/m<sup>3</sup> in Wicklow. The maximum value of 1032 Bq/m<sup>3</sup> was measured in a house in Co. Wicklow.

**Table 1. Summary of Survey Data**

County	Number of Houses Measured	Measured Average Annual Radon Concentration (Bq/m <sup>3</sup> )	Measured Maximum Annual Radon Concentration (Bq/m <sup>3</sup> )	Measured Number of Dwellings above Reference Level
Cavan	179	72	780	9
Dublin	128	81	260	6
Louth	104	122	751	15
Monaghan	116	69	365	5
Wicklow	164	138	1032	24

A total of 59 of the dwellings surveyed had radon concentrations in excess of the Reference Level of 200 Bq/m<sup>3</sup>. The number of these dwellings in each county, expressed as a percentage of the total number measured in that county, highlights the geographical variation in indoor radon: in Cavan, Dublin and Monaghan the value was between 4% and 5%, similar to the national figure determined by the UCD survey, whereas in Louth and Wicklow the values were 14% and 15%, respectively.

#### **4.2 High Radon Areas**

The radon measurement data presented above were used to predict the percentage of dwellings with radon concentrations in excess of 200 Bq/m<sup>3</sup> in each grid square of counties Cavan, Dublin, Louth, Monaghan and Wicklow according to the method described in Section 3. The results are presented in Figures 1 - 5. The figures are colour coded with each colour corresponding to a percentage band i.e. < 1%, 1-4%, 4-10%, 10-20% and > 20%. The two percentage bands 10-20% and > 20% delimit the High Radon Areas.

From the figures it is again evident that there is considerable geographical variation in indoor radon concentrations. In counties Dublin and Monaghan and to a lesser extent Cavan, the High Radon Areas are limited, with large areas predicted to have < 4% of dwellings with radon concentrations in excess of 200 Bq/m<sup>3</sup>. The opposite is the case in Co. Wicklow where 10 of the 19 grid squares for which predictions have been made are designated High Radon Areas. In Co. Louth, the Cooley Peninsula and the area surrounding Clogherhead are categorised as High Radon Areas.

In estimating the scale of the radon problem in any county, it must be borne in mind that this will depend on the housing density in the county as well as on the extent of High Radon Areas within the county.

## **5. Doses and Risk Estimates**

The epidemiological evidence for the induction of fatal lung cancer by radon decay products has been derived both for miners exposed to radon at work and for persons exposed at home. In this report, estimates of risk from long-term exposure to radon are based on the lifetime fatality coefficient recommended in the latest relevant report of the International Commission on Radiological Protection [ICRP, 1994]. For a person living in a house with a radon concentration of 200 Bq/m<sup>3</sup>, the lifetime excess lung cancer risk is 2%. At higher radon concentrations, the risk is proportionately greater.

This risk estimate may be compared with the observed lifetime risk of lung cancer from all sources of 3% for the whole population of Ireland. It is also of interest to note that for people who smoke, there is growing evidence that the risk from radon is considerably greater than these figures, while for non-smokers the risk from radon is less than the average figure for the population as a whole.

To allow comparison with radiation doses arising from other sources, the radiation dose due to radon exposure can be calculated using an exposure-dose conversion factor of 1 millisievert (mSv) annual radiation dose per 40 Bq/m<sup>3</sup> radon concentration in a dwelling. The derivation of this factor is detailed elsewhere [Madden *et al.*, 1994]. Annual exposure in the home to a radon concentration of 200 Bq/m<sup>3</sup> would therefore give rise to a radiation dose of 5 mSv.

When compared with the estimated average annual radiation dose of approximately 3 mSv per year in Ireland from all sources, the data show that radon exposure in the home is the major source of human exposure to ionising radiation in Ireland. It is also an extremely variable source, with exposures in individual homes in the counties reported here ranging from less than 1 mSv to about 25 mSv.

## **6. Conclusions and Recommendations**

The initial results of the National Radon Survey reported here show that there is significant geographical variation in indoor radon concentrations in counties Cavan, Dublin, Louth, Monaghan and Wicklow. Approximately half the area of Co. Wicklow and smaller areas in counties Louth, Cavan, Monaghan and Dublin are designated as High Radon Areas. In these areas, the Institute would encourage householders to have radon measurements made in their homes in order to identify the individual dwellings with elevated indoor radon levels. Where high radon levels are found, householders are encouraged to take the necessary action to reduce the risks to themselves and their families. While the High Radon Areas can, by definition, be expected to have a higher proportion of homes with high radon levels, it cannot be inferred that elevated indoor radon concentrations do not occur in homes in other areas that are not designated as High Radon Areas.



The Institute would also advise that all new dwellings being built in High Radon Areas include some radon preventive measure. Advice on the most appropriate measure to be taken in either new or existing dwellings is available in the publication "Radon in Buildings" [Ryan and Finn, 1995], which is available from the Government Publications Sales Office.

## **7. Acknowledgements**

The authors wish to thank Dr James McLaughlin, Department of Experimental Physics, University College Dublin, for his expert advice on many aspects of the work. They are grateful to Mr Michael Keatinge\*, Zoology Department, Trinity College Dublin, for his advice in devising the survey and to Dr Leslie Daly, Department of Public Health Medicine and Epidemiology, University College Dublin, for his assistance with statistical evaluation of the data.

The authors are particularly grateful to all the householders who participated in the survey. The assistance of Ms Louisa Smyth in the administration of the survey and of the RPII Information Service personnel is greatly appreciated.

*\*Present address: Fisheries Research Centre, Dublin.*

## 8. References

Commission of the European Communities, 1987. **Exposure to natural radiation in dwellings of the European Communities**. Luxembourg: Commission of the European Communities.

Daly, L., 1994. Personal communication.

International Commission on Radiological Protection, 1977. **Radiation protection in uranium and other mines**. ICRP Publication 24, Oxford: Pergamon Press.

International Commission on Radiological Protection, 1994. **Protection against radon-222 at home and at work**. ICRP Publication 65, Oxford: Pergamon Press.

Liberman, G.J. and Resnikoff, G.J., 1955. Sampling plans for inspection by variables. **American Statistical Association Journal**, June 1955, p.457-516.

Madden, J.S., Duffy, J.T., Mackin, G.A., Colgan, P.A. and McGarry, A.T., 1994. **Radon in dwellings in selected areas of Ireland**. RPII-94/3, Dublin: Radiological Protection Institute of Ireland.

McLaughlin, J.P. and Wasiolek, P., 1988. Radon levels in Irish dwellings. **Radiation Protection Dosimetry**, 24,(1/4), p.383-386.

Miles, J.C.H., 1994. Mapping the proportion of the housing stock exceeding a radon reference level. **Radiation Protection Dosimetry**, 56,(1/4), p.207-210.

National Council on Radiation Protection and Measurements, 1988. **Measurement of radon and daughters in air**. NCRP Report No. 97, Bethesda: NCRP.

National Radiological Protection Board, 1990. Radon affected areas: Cornwall and Devon. **Documents of the NRPB**, 1,(4), p.37-43.

Nazaroff, W.W., Moed, B.A. and Sextro, R.G., 1988. Soil as a source of indoor radon: Generation, migration and entry. In **Radon and its decay products in indoor air**. p.57-112. Nazaroff, W.W. and Nero, A.V. (Jr) (eds.). New York: John Wiley and Sons.

Nero, A.V., Boegal, M.L., Hollowell, C.D., Ingersoll, J.G. and Nazaroff, W.W., 1983. Radon concentrations and infiltration rates measured in conventional and energy-efficient houses. **Health Physics**, 45,(2), p. 401-405.

Ryan, N.M. and Finn, M., 1995. **Radon in buildings**. Department of the Environment. Dublin: Stationery Office.


## Glossary of Terms

**Radon Concentration** - the activity concentration of radon gas in terms of activity per unit volume of air. Activity is given in units of becquerel (Bq), with activity concentration given in becquerels per cubic metre (Bq/m<sup>3</sup>). A becquerel is equal to one nuclear transformation per second.

**Half-Life** - the time taken for the activity of a radionuclide to lose half its value by decay.


**Lifetime Fatality Coefficient** - the risk of death associated with exposure to radon and its decay products over a lifetime of 70 years.

**Radiation Dose** - a general term for the quantity of radiation. It takes account of the type of radiation, the tissues exposed and their sensitivity to radiation. Dose is given in units of sievert (Sv). One Sv produces the same biological effect irrespective of the type of radiation.



Grid: Irish National Grid

### Legend



**Map of County Cavan**  
 showing the percentage of dwellings predicted to exceed  
 200 Bq/m<sup>3</sup> in each 10 km grid square.





Figure 1.



Grid: Irish National Grid

**Legend**

- > 20%
- 10 - 20%
- 4 - 10%
- 1 - 4%
- < 1%
- Insufficient Data


**Map of County Dublin**  
 showing the percentage of dwellings predicted to exceed  
 200 Bq/m<sup>3</sup> in each 10 km grid square.

0 5 10 15 Kilometres

0 5 Miles

Scale 1: 600,000

**Figure 2.**




Grid: Irish National Grid

**Legend**


- > 20%
- 10 - 20%
- 4 - 10%
- 1 - 4%
- < 1%
- Insufficient Data

**Map of County Louth**  
 showing the percentage of dwellings predicted to exceed  
 200 Bq/m<sup>3</sup> in each 10 km grid square.



Scale 1: 600,000

**Figure 3.**



Grid: Irish National Grid

**Legend**

- > 20%
- 10 - 20%
- 4 - 10%
- 1 - 4%
- < 1%
- Insufficient Data


**Map of County Monaghan**  
 showing the percentage of dwellings predicted to exceed  
 200 Bq/m<sup>3</sup> in each 10 km grid square.

0 5 10 15 Kilometres

0 5 Miles

Scale 1: 600,000

**Figure 4.**




Grid: Irish National Grid

**Legend**

- > 20%
- 10 - 20%
- 4 - 10%
- 1 - 4%
- < 1%
- Insufficient Data

**Map of County Wicklow**  
 showing the percentage of dwellings predicted to exceed  
 200 Bq/m<sup>3</sup> in each 10 km grid square.



**Figure 5.**