

CONF-9603173--2/

**ERNEST ORLANDO LAWRENCE
BERKELEY NATIONAL LABORATORY**

**Fragmentation Functions Approach
in pQCD Fragmentation Phenomena**


S. Rolli
Physics Division

RECEIVED

NOV 14 1996

OSTI

July 1996
Presented at the
*XXXI Rencontres de Moriond,
QCD and Hadronic Interactions,*
Les Arcs, France
March 24-31, 1996,
and to be published in
the Proceedings



DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

MASTER

DISCLAIMER

Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor The Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or The Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof, or The Regents of the University of California.

This report has been reproduced directly from the best available copy.

Ernest Orlando Lawrence Berkeley National Laboratory
is an equal opportunity employer.

LBNL-39176
UC-000

**Fragmentation Functions Approach in
pQCD Fragmentation Phenomena**

Simona Rolli

INFN-Pavia, Italy

and

Physics Division
Ernest Orlando Lawrence Berkeley National Laboratory
University of California
Berkeley, California 94720

July 1996

This work was supported in part by the "Maria Rossi" fellowship from Collegio Ghislieri of Pavia, through the U.S. Department of Energy under Contract No. DE-AC03-76SF00098.

Fragmentation Functions approach in pQCD fragmentation phenomena

Simona Rolli

INFN-Pavia, Italy and Lawrence Berkeley Laboratory, Berkeley, USA
E-mail: rolli@fnal.gov

Abstract

Next-to-leading order parton fragmentation functions into light mesons are presented. They have been extracted from real and simulated e^+e^- data and used to predict inclusive single particle distributions at different machines.

Talk given at the XXXI Rencontres de Moriond, QCD and hadronic interactions,
Les Arcs, France March 24-31 1996

1 Introduction

One of the fundamental properties of QCD is the shrinking of the coupling constant as the energy of the interactions grows (asymptotic freedom). This implies that perturbative techniques can be used to study high energy hadronic or leptonic phenomena. In spite of this, we can not rely completely on perturbation theory because the fundamental particles whose interactions become weak at high energy are deeply bound inside the hadrons we use as beams, targets or as observables. The solution is given by the factorization theorems, whereby cross sections can be expressed as the products of factors, each one involving phenomena appearing at different energy scales. A generic hadronic collisions with production of one hadron inclusive can be expressed as the convolution of the partonic hard cross section and structure and fragmentation functions, which represent respectively the parton density inside the hadrons and the hadron density inside the parton. Those parton model distributions acquire a scale dependence when the QCD corrections are included, the scale being the point where collinear initial (final) state divergencies are subtracted. Moreover these distribution are independent of the specific reaction. The universality of these distributions is a key property, since they are not calculated from first principles as they contain non-perturbative information. Nevertheless they evolve with the scale following Altarelli-Parisi type equations and this permits to extract them from a low-energy process and use them to predict rates for another one.

In this talk I'll concentrate on some recent sets of parton fragmentation functions into light mesons. These sets have been extracted from real and simulated e^+e^- data in a full NLO formalism. They have been used to predict inclusive single particle differential distributions at different energies and machine and to study jet fragmentation properties. The outline of the talk is the following: I'll describe briefly how the sets have been derived, then I'll give some results on the inclusive single particle predictions comparing with data when they are available.

2 Fragmentation functions from e^+e^- data

Parton fragmentation functions are defined as the density probability to find a hadron h inside a parton p . They are parametrized in the following form:

$$D_p^h(x, M_f) = N x^\alpha (1-x)^\beta \quad (1)$$

where x is the fraction of parton momentum carried by the hadron, while M_f is the scale the function is evaluated. They satisfied multiplicity, charge and momentum conservation sum rules and evolve following Altarelli Parisi type equation. This means that given a certain set of input functions at the scale M_{f0}^2 we can obtain the functions at any desired scale simply solving the AP equation. To obtain the input sets we need to rely on experimental informations, because FF are not calculable from first principle, owing the fact that they contain non-perturbative informations¹.

To extract the FF we use e^+e^- data. In the language of the QCD-improved parton model, in fact, the x distribution of the process $e^+e^- \rightarrow h + X$ emerges from the x distribution $(d\sigma/dx)(x, \mu^2, s)$ of $e^+e^- \rightarrow a + X$ through convolution with $D_a^h(x, M_f^2)$:

$$\frac{1}{\sigma_{TOT}} \frac{d\sigma(e^+e^- \rightarrow h + X)}{dx} = \sum_a \int_x^1 \frac{dz}{z} D_a^h(z, M_f^2) \frac{1}{\sigma_{TOT}} \frac{d\sigma_a}{dy} \left(\frac{x}{z}, \mu^2, M_f^2 \right); \quad (2)$$

where $\sigma_{TOT} = N_c \sum_{i=1}^{N_f} e_q^2 \sigma_0$ and

¹we can actually calculate in a completely independent way only heavy quarks fragmentation functions, because in this case the mass of the heavy object act as a natural cutoff.

$$\frac{1}{\sigma_{TOT}} \frac{d\sigma_{qi}}{dy} (y, \mu^2, M_f^2) = e_q^2 N_c \frac{\sigma_0}{\sigma_{TOT}} \left\{ \delta(1-y) + \frac{\alpha_s(\mu^2)}{2\pi} \left[P_{qq}^0(y) \ln\left(\frac{s}{M_f^2}\right) + K_q^T(y) + K_q^L(y) \right] \right\} \quad (3)$$

and

$$\frac{1}{\sigma_{TOT}} \frac{d\sigma_g}{dy} (y, \mu^2, M_f^2) = \frac{2\alpha_s(\mu^2)}{2\pi} \left[P_{gg}^{(0,T)}(y) \ln\left(\frac{s}{M_f^2}\right) + K_g^T(y) + K_g^L(y) \right] \quad (4)$$




Figure 1: K^\pm (upper right), K_s^0 (upper left) and η (lower) production using sets from ref.[6,8]




Fig. 2

Figure 2: neutral kaons production at e^+e^- using the sets from ref.[5]

The functions K_q^T , K_q^L , K_g^T and K_g^L are shown in reference [1] and represent the NLO corrections. Using real and simulated[2] data several different sets have been derived for neutral[1] and charged pions[3, 4], neutral and charged kaons[6, 5] and η mesons[8] at different reference scales. In Figure 1 and 2 we give some examples on how well the different sets, when convoluted with the partonic cross sections, agree with e^+e^- data. We refer the readers to the previous references for further details.

3 Applications

As already stated, fragmentation functions are process independent: this means that we can use them to make predictions for process different from the one we used to extract them. We simply need to convolute them with the appropriate partonic cross sections. In the previous references the different sets have been used to give predictions on inclusive single particle production in hadron-hadron, hadron-lepton and photon-photon[9] collisions, and to phenomenologically describe jet fragmentation properties. In Figures 3, 4 and 5 we show again some examples on the NLO predictions for different energy ranges and process types.

Finally, in order to disentangle the fragmentation properties and the hadronization mechanism of high p_t jets, we consider the ratio between the single hadron and jet cross sections, for fixed values of the variable $z = E_{hadr}/E_{jet}$. Then, using the cone jet algorithm and the NLO evaluation of the jet cross sections of ref. [10], we present in Figure 6 a result on jet fragmentation in charged and neutral pions, with the energy of the jet varying between 40




Figure 3: π^0 production at ISR and SppS using sets from ref.[1]




Fig. 5

Figure 4: K_s^0 production at Tevatron using sets from ref.[5]. The dotted line is from ref.[6]

and 70 GeV, and a jet cone radius $R=0.7$ centered around the $\eta = 0$ direction. The overall theoretical uncertainty -which is not reported in figure- can be estimated to be of order 50%. We also show the analogous experimental result on jet fragmentation in charged hadrons from CDF, in reasonable agreement with the theoretical prediction.




Fig. 3

Figure 5: K_s^0 production at HERA, using sets from ref.[5]




Figure 6: jet fragmentation function, into π^0 and $(\pi^+ + \pi^-)$. The experimental points refer to charged hadrons and are from CDF

4 Conclusions

In this talk I reviewed the results on some recent NLO sets of parton fragmentation functions into light mesons. The possibilities to use them to make reliable predictions at different type of machines has been briefly shown.

5 Acknowledgement

This work was supported in part by the "Maria Rossi" fellowship from Collegio Ghislieri of Pavia, through the U.S. Department of Energy under contract DE-AC03-76SF00098.

References

- [1] P. Chiappetta, M. Greco, J.Ph. Guillet, S. Rolli, M. Werlen, *Next-to-leading-order determination of pion fragmentation functions*, Nucl. Phys. B 412 (1994) 3;
- [2] G. Marchesini et al., Comp. Phys. Comm. 67(1992)465;
- [3] M. Greco, S. Rolli, A. Vicini, *Inclusive single particle photoproduction to next-to-leading order*, Z. Phys. C65 (1995) 277;
- [4] J. Binnewies, B. A. Kniehl and G. Kramer, *Next-to-leading order fragmentation functions for pions and kaons*, Z. Phys. C65 (1995) 471
- [5] J. Binnewies, B. A. Kniehl and G. Kramer, *Pion and kaon production in e^+e^- and ep collisions at next-to-leading order* Phys. Rev. D52 (1995) 4947;
- [6] M. Greco, S. Rolli, *Light mesons production at the Tevatron to next-to-leading order* Phys. Rev. D52 (1995) 3853;
- [7] J. Binnewies, B. A. Kniehl and G. Kramer, *Neutral kaon production in e^+e^- and ep $p\bar{p}$ collisions at next-to-leading order*, hep-ph/9506437, Phys. Rev. D (in press);
- [8] M. Greco, S. Rolli, *Next-to-leading-order eta production at hadron colliders*, Z. Phys. C60 (1993) 169;
- [9] J. Binnewies, B. A. Kniehl and G. Kramer, *Inclusive hadron production in photon-photon collisions at next-to-leading order* hep-ph/9601278, DESY-96-001, submitted to Phys. Rev. D.
- [10] F. Aversa, P. Chiappetta, M. Greco e J. Ph. Guillet, Nucl. Phys. B327 (1989) 105