




XJ9800110



ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

E2-97-267

N.A.Chernikov, N.G.Fadeev, N.S.Shavokhina

PROPER TIME AXIS OF A CLOSED
RELATIVISTIC SYSTEM

Submitted to the International Workshop «Non-Euclidean Geometry
in Modern Physics», August 13—16, 1997, Uzhgorod, Ukraine

2
29-19

1997

Let us consider a closed relativistic system. In our consideration the space-time is a four dimensional pseudo-Euclidean space $E_{(1,3)}$ with the following metric:

$$\theta_{ab} dx^a dx^b = dt^2 - \frac{1}{c^2} (dx^2 + dy^2 + dz^2), \quad (1)$$

$$a = 0, 1, 2, 3,$$

c — being the light velocity.

Let us denote by p^a the 4-momentum of a particle, where

$$p^a = p^0 \frac{dx^a}{dt},$$

$$p^0 = m / \sqrt{1 - \frac{1}{c^2} \left(\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right)}, \quad (2)$$

m — being the mass of the particle. Consequently,

$$m^2 = \theta_{ab} p^a p^b. \quad (3)$$

We denote by p_a the following components:

$$p_a = \theta_{ab} p^b. \quad (4)$$

Let us denote by m^{ab} the angular 4-momentum of the particle. It equals

$$m^{ab} = x^a p^b - x^b p^a, \quad (5)$$

where x^a are the coordinates of the particle. This momentum is associated with the origin of coordinates.

Now let us consider a Cauchy hyper-surface Σ in $E_{(1,3)}$ and a system of N particles on it, not yet their interacting. The Cauchy hyper-surface in $E_{(1,3)}$ is a space-like one. Let us define the 4-momentum of the system as

$$P^a = \sum_{k=1}^N p_{(k)}^a \quad (6)$$

and the angular 4-momentum of the system as

$$M^{ab} = \sum_{k=1}^N m_{(k)}^{ab}. \quad (7)$$

Here the index k numbers our particles,

$$m_{(k)}^{ab} = x_{(k)}^a p_{(k)}^b - x_{(k)}^b p_{(k)}^a. \quad (8)$$

The points $x_{(k)}^a$ and the momenta $p_{(k)}^a$ are considered on the hyper-surface Σ .

During the interaction the number of particles N may be changed, but not the vector P^a and the bivector M^{ab} : the last ones are integrals of the equations of motion, since our system is closed.

The angular momenta (7) and (8) are associated with the origin coordinates O . Let us replace the point O with a point $\check{O}(\check{t}, \check{x}, \check{y}, \check{z})$. Correspondingly, we replace $x_{(k)}^a$ with $(x_{(k)}^a - \check{x}^a)$, $m_{(k)}^{ab}$ with

$$\check{m}_{(k)}^{ab} = m_{(k)}^{ab} - \check{x}^a p_{(k)}^b + \check{x}^b p_{(k)}^a, \quad (9)$$

and M^{ab} with

$$\check{M}^{ab} = M^{ab} - \check{x}^a P^b + \check{x}^b P^a. \quad (10)$$

Consequently, if $\check{x}^a = \lambda P^a$, then $\check{M}^{ab} = M^{ab}$. Here λ is an arbitrary number.

Now let us find such points, so that

$$\check{M}^{ab} P_b = 0. \quad (11)$$

Inserting here (10), we receive the following equation for

the proper time axis of our system:

$$(\check{x}^a P^b - \check{x}^b P^a) P_b = M^{ab} P_b. \quad (12)$$

One can see that if a point \check{x}_0^a is a solution of the equation (12), then the point $\check{x}_0^a + \lambda P^a$ is a solution of the same equation too.

In our case

$$P^0 > 0, (P, P) = P^a P_a = \theta_{ab} p^a p^b > 0; \quad (13)$$

the set of solutions of equation (12) is a time-like straight line

$$\check{x}^a = \check{x}_0^a + \lambda P^a \quad (14)$$

in the space-time $E_{(1,3)}$. This straight line is the proper time axis of our system.

Let us write down equation (12) in the form:

$$\check{x}^a (P, P) - P^a (P, \check{x}) = M^{ab} P_b. \quad (15)$$

$$(P, \check{x}) = P_b \check{x}^b.$$

Setting $a = 0$, we have

$$\check{t}(P, P) - P^0(P, \check{x}) = M^{0b} P_b. \quad (16)$$

From (15) and (16) we get

$$\check{x}^a = \frac{P^a}{P^0} \check{t} + \frac{P^0 M^{ab} P_b - P^a M^{0b} P_b}{P^0(P, P)}. \quad (17)$$

This is another form of writing down Eq (14).

If $P^1 = P^2 = P^3 = 0$, then

$$\check{x}^\alpha = \frac{M^{\alpha 0}}{P^0}. \quad (18)$$

In this case the proper time axis is parallel to the coordinate time axis.

If $P^1 = P^2 = P^3 = 0$, and $M^{10} = M^{20} = M^{30} = 0$, then

$$\check{x}^\alpha = 0. \quad (19)$$

In the last case the proper time axis coincides with the coordinate time axis.

Let us consider now the case when the system moves in the plane $z = 0$. Then $p_{(k)}^3 = 0$ and $m_{(k)}^{a3} = 0$. Consequently, in this case $P^3 = 0$, $M^{a3} = 0$ and $\check{M}^{a3} = 0$. If we denote

$$\check{M}_0 = \check{M}^{12}, \quad \check{M}_1 = \check{M}^{20}, \quad \check{M}_2 = \check{M}^{01}, \quad (20)$$

then

$$\begin{aligned} \check{M}^{0b} P_b &= \check{M}_2 P_1 - \check{M}_1 P_2, \\ \check{M}^{1b} P_b &= \check{M}_0 P_2 - \check{M}_2 P_0, \quad \check{M}^{2b} P_b = \check{M}_1 P_0 - \check{M}_0 P_1. \end{aligned} \quad (21)$$

From Eqs. (11) and (21) it follows

$$\frac{\check{M}_0}{P_0} = \frac{\check{M}_1}{P_1} = \frac{\check{M}_2}{P_2}. \quad (22)$$

Inserting here (10), we receive the following equations for the proper time axis of our system in the case $z = 0$:

$$\begin{aligned} & \frac{M_0 - \check{x}^1 P^2 + \check{x}^2 P^1}{P_0} = \\ & = \frac{M_1 - \check{x}^2 P^0 + \check{x}^0 P^2}{P_1} = \frac{M_2 - \check{x}^0 P^1 + \check{x}^1 P^0}{P_2}. \end{aligned} \quad (23)$$

In engineering statics of an absolutely rigid body the problem about reducing of a system of forces to the dynamic screw is considered (see for instance [4]). The dynamic screw consists of a force and a couple, lying in a plane, perpendicular to this force.

In the case $z = 0$, starting from equations (23), we come upon the problem about the dynamic screw, assuming that the light velocity c is equal to the imaginary unit i . In such a way the particle momenta play the role of forces, acting on the rigid body.

In the general case, setting up in Eq.(1) $c^2 = -1$ and starting from (12), we arrive at the problem about the dynamic screw in a four-dimensional Euclidean space.

References

- [1] *Chernikov N.A, Shavokhina N.S.* – A linear potential at high energies. Proc. of 5th Int. Conf. on problems of quantum field theory. Alushta, 1979. JINR, Dubna, P2-12462, p.340.
- [2] *Chernikov N.A, Shavokhina N.S.* – A special formulation of the relativistic two bodies problem. In collected volume: Problems of theory of gravitation and elementary particles. No 14, Moscow, 1986, p. 113.
- [3] *Shavokhina N.S.* – A problem on a world surface which spans world trajectories of particles. Russian Mathematics (Iz. VUZ) Vol. 38, No 2, 1994, p.67.
- [4] *Voronkov I.M.* – A Course of Theoretical Mechanics. Gostechisdat, Moscow, 1955, p. 182.

Received by Publishing Department
on September 4, 1997.

**The Publishing Department
of the Joint Institute for Nuclear Research
offers you to acquire the following books:**

| Index | Title |
|--------------|--|
| 94-55 | Proceedings of the International Bogoliubov Memorial Meeting. Dubna, 1993 (216 p. in Russian and English) |
| E7-94-270 | Proceedings of the Workshop on Physical Experiments and First Results on Heavy Ion Storage and Cooler Rings. Smolenice, 1992 (324 p. in English) |
| E2-94-347 | International Workshop «Symmetry Methods in Physics». In Memory of Professor Ya.A.Smorodinsky. Dubna, 1993 (2 volumes, 602 p. in English) |
| E4-94-370 | Proceedings of the IV International Conference on Selected Topics in Nuclear Structure. Dubna, 1994 (412 p. in English) |
| E4-94-386 | Proceedings of the VI Trilateral German-Russian-Ukrainian Seminar on High-Temperature Superconductivity. Dubna, 1994 (340 p. in English) |
| D2-94-390 | D.I.Blokhintsev. Proceedings of the Seminars. Dedicated to the 85th Anniversary of the Birthday of D.I.Blokhintsev. Dubna, 1995 (271 p. in Russian and English) |
| E3-94-419 | Proceedings of the II International Seminar on Neutron-Nucleus Interactions (ISINN-2) «Neutron Spectroscopy, Nuclear Structure, Related Topics». Dubna, 1994 (363 p. in English) |
| D13-94-491 | Proceedings of the XVI International Symposium on Nuclear Electronics and VI International School on Automation and Computing in Nuclear Physics and Astrophysics. Varna, 1994 (246 p. in Russian and English) |
| D13,14-95-49 | Proceedings of the International Symposium on Muon and Pion Interactions with Matter. Dubna, 1994 (304 p. in Russian and English) |
| D3-95-169 | Proceedings of the International Seminar ADVANCED PULSED NEUTRON SOURCES: Physics of/at ADVANCED Pulsed Neutron Sources. PANS-II. Dubna, 1994 (336 p. in Russian and English) |
| D1-95-305 | Proceedings of the XVII Workshop on Neutrino Detector IHEP—JINR. Dubna, 1995 (178 p. in Russian and English) |
| E3-95-307 | Proceedings of the III International Seminar on Interaction of Neutrons with Nuclei. Neutron Spectroscopy, Nuclear Structure, Related Topics. Dubna, 1995 (356 p. in English) |
| D3,14-95-323 | VII School on Neutron Physics. Lectures. Vol.1. Dubna, 1995 (356 p. in Russian and English) |

| Index | Title |
|---------------|---|
| E10,11-95-387 | Proceedings of the ESONE International Conference RTD'94 on REAL TIME DATA 1994 with Emphasis on Distributed Front-End Processing. Dubna, 1994 (358 p. in English) |
| D15-96-18 | Proceedings of the International Workshop Charge and Nucleon Radii of Exotic Nuclei. Poznan, 1995 (172 p. in Russian and English) |
| E9-96-21 | Proceedings of VII ICFA Beam Dynamics Workshop on «Beam Issues for Multibunch, High Luminosity Circular Colliders». Dubna, 1995 (198 p. in English) |
| E2-96-100 | Proceedings of the 3rd International Symposium «Dubna Deuteron-95». Dubna, 1995 (374 p. in English) |
| E2-96-224 | Proceedings of the VII International Conference «Symmetry Methods in Physics». Dubna, 1996 (2 volumes, 630 p., in English) |
| E-96-321 | Proceedings of the International Conference «Path Integrals: Dubna'96». Dubna, 1996 (392 p. in English) |
| E3-96-336 | Proceedings of the IV International Seminar on Interaction of Neutrons with Nuclei. Dubna, 1996 (396 p. in English) |
| E3-96-369 | Proceedings of the X International Conference «Problems of Quantum Field Theory». Dubna, 1996 (437 p. in English) |
| E3-96-507 | Proceedings of the International Workshop «Polarized Neutrons for Condensed Matter Investigations». Dubna, 1996 (154 p. in English) |
| D1,2-97-6 | Proceedings of the International Workshop «Relativistic Nuclear Physics: from MeV to TeV». Dubna, 1996 (2 volumes 418 p. and 412 p. in English and Russian) |
| E7-97-49 | Proceedings of the 3rd International Conference «Dynamical Aspects of Nuclear Fission». Slovakia, 1996 (426 p. in English) |
| E1,2-97-79 | Proceedings of the XIII International Seminar on High Energy Physics Problems. Relativistic Nuclear Physics and Quantum Chromodynamics. Dubna, 1996 (2 volumes, 364 p. and 370 p. in English) |

Please apply to the Publishing Department of the Joint Institute for Nuclear Research for extra information. Our address is:

Publishing Department
 Joint Institute for Nuclear Research
 Dubna, Moscow Region
 141980 Russia
 E-mail: publish@pds.jinr.dubna.su

Собственная ось времени замкнутой релятивистской системы

Дано определение собственной оси времени замкнутой релятивистской системы сталкивающихся частиц, и найдено решение задачи о ее нахождении. Если скорость света c приравнять мнимой единице i , то в случае плоского движения системы задача о собственной оси времени оказывается эквивалентной известной в инженерной механике задаче о приведении к динаме (к динамическому винту) системы сил, приложенных к твердому телу. В общем случае при $c = i$ задача о собственной оси времени оказывается эквивалентной задаче о приведении к динаме системы сил, приложенных к твердому телу в четырехмерном евклидовом пространстве.

Работа выполнена в Лаборатории теоретической физики им. Н.Н.Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1997

Proper Time Axis of a Closed Relativistic System

The definition of a proper time axis of a closed relativistic system of colliding particles is given. The solution of the proper time axis problem is presented. If the light velocity c equals the imaginary unit i , then in the case of a plane motion of the system the problem about the proper time axis turns out to be equivalent to the known in engineering mechanics problem about the reduction of any system of forces, applied to a rigid body, to the dynamic screw. In the general case, when $c = i$, the problem about the proper time axis turns out to be equivalent to the problem about the reduction to the dynamic screw of a system of forces, applied to a rigid body in a four-dimensional Euclidean space.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1997

Макет Т.Е.Попеко

Подписано в печать 22.09.97
Формат 60 × 90/16. Офсетная печать. Уч.-изд.листов 0,83
Тираж 460. Заказ 50159. Цена 1116 р.

Издательский отдел Объединенного института ядерных исследований
Дубна Московской области