

THE HEAVY QUARK EXPANSION OF QCD

Adam F. Falk
Department of Physics and Astronomy
Johns Hopkins University
3400 North Charles Street
Baltimore, Maryland 21218

ABSTRACT

These lectures contain an elementary introduction to heavy quark symmetry and the heavy quark expansion. Applications such as the expansion of heavy meson decay constants and the treatment of inclusive and exclusive semileptonic B decays are included. Heavy hadron production via nonperturbative fragmentation processes is also discussed.

© 1996 by Adam F. Falk.

1 Heavy Quark Symmetry

In these lectures, I will introduce the ideas of heavy quark symmetry and the heavy quark limit, which exploit the simplification of certain aspects of QCD for infinite quark mass, $m_Q \rightarrow \infty$. We will see that while these ideas are extraordinarily simple from a physical point of view, they are of enormous practical utility in the study of the phenomenology of bottom and charmed hadrons. One reason for this is the existence not just of an interesting new limit of QCD, but of a systematic expansion about this limit. The technology of this expansion is the Heavy Quark Effective Theory (HQET), which allows one to use heavy quark symmetry to make accurate predictions of the properties and behavior of heavy hadrons in which the theoretical errors are under control. While the emphasis in these lectures will be on the physical picture of heavy hadrons which emerges in the heavy quark limit, it will be important to introduce enough of the formalism of the HQET to reveal the structure of the heavy quark expansion as a simultaneous expansion in powers of Λ_{QCD}/m_Q and $\alpha_s(m_Q)$. However, what I hope to leave you with, above all, is an appreciation for the simplicity, elegance, and coherence of the ideas which underlie the technical results which will be presented. The interested reader is also encouraged to consult a number of excellent reviews,¹ which typically cover in more detail the material in the first two sections of these lectures.

1.1 Introduction

Why is an understanding of QCD crucial to the study of the properties of the bottom and charmed quarks? As an example, consider semileptonic b decay, $b \rightarrow c \ell \bar{\nu}$. This process is mediated by a four-fermion operator,

$$\mathcal{O}_{bc} = \frac{G_F V_{cb}}{\sqrt{2}} \bar{c} \gamma^\mu (1 - \gamma^5) b \bar{\nu} \gamma_\mu (1 - \gamma^5) \ell. \quad (1)$$

The weak matrix element is easy to calculate at the quark level,

$$\mathcal{A}_{\text{quark}} = \langle c \ell \bar{\nu} | \mathcal{O}_{bc} | b \rangle = \frac{G_F V_{cb}}{\sqrt{2}} \bar{u}_c(p_c) \gamma^\mu (1 - \gamma^5) u_b(p_b) \bar{u}_\ell(p_\ell) \gamma_\mu (1 - \gamma^5) v_\nu(p_\nu). \quad (2)$$

However, $\mathcal{A}_{\text{quark}}$ is only relevant at very short distances; at longer distances, QCD confinement implies that free b and c quarks are not asymptotic states of the theory. Instead, nonperturbative QCD effects “dress” the quark level transition $b \rightarrow c \ell \bar{\nu}$ to a hadronic transition, such as

$$B \rightarrow D \ell \bar{\nu} \quad \text{or} \quad B \rightarrow D^* \ell \bar{\nu} \quad \text{or} \quad \dots \quad (3)$$

(In these lectures, we will use a convention in which a B meson contains a b quark, not a \bar{b} antiquark.) The hadronic matrix element $\mathcal{A}_{\text{hadron}}$ depends on nonperturbative QCD as well as on $G_F V_{cb}$, and is difficult to calculate from first principles. To disentangle the weak interaction part of this complicated process requires us to develop some understanding of the strong interaction effects.

There are a variety of methods by which one can do this. Perhaps the most popular, historically, has been use of various quark potential models.² While these models are typically very predictive, they are based on uncontrolled assumptions and approximations, and it is virtually impossible to estimate the theoretical errors associated with their use. This is a serious defect if one builds such a model into the experimental extraction of a weak coupling constant such as V_{cb} , because the uncontrolled theoretical errors then infect the experimental result.

Another approach, the one to be discussed here, is to exploit as much as possible the fact that the b and c quarks are *heavy*, by which we mean that $m_b, m_c \gg \Lambda_{\text{QCD}}$. The scale Λ_{QCD} is the typical energy at which QCD becomes nonperturbative, and is of the order of hundreds of MeV. The physical quark masses are approximately $m_b \approx 4.8 \text{ GeV}$ and $m_c \approx 1.5 \text{ GeV}$. The formalism which we will develop will not make as many predictions as do potential models. However, the compensation will be that we will develop a *systematic* expansion in powers of $\Lambda_{\text{QCD}}/m_{b,c}$, within which we will be able to do concrete error analysis. In particular, we will be able to estimate the error associated with the fact that m_c may not be very close to the asymptotic limit $m_c \gg \Lambda_{\text{QCD}}$. Even where this error may be substantial, the fact that it is under control allows us to maintain predictive power in the theory.

1.2 The Heavy Quark Limit

Consider a hadron H_Q composed of a heavy quark Q and “light degrees of freedom,” consisting of light quarks, light antiquarks, and gluons, in the limit $m_Q \rightarrow \infty$. The Compton wavelength of the heavy quark scales as the inverse of the heavy quark mass, $\lambda_Q \sim 1/m_Q$. The light degrees of freedom, by contrast, are characterized by momenta of order Λ_{QCD} , corresponding to wavelengths $\lambda_\ell \sim 1/\Lambda_{\text{QCD}}$. Since $\lambda_\ell \gg \lambda_Q$, the light degrees of freedom cannot resolve features of the heavy quark other than its conserved gauge quantum numbers. In particular, they cannot probe the actual *value* of λ_Q , that is, the value of m_Q .

We draw the same conclusion in momentum space. The structure of the hadron H_Q is determined by nonperturbative strong interactions. The asymptotic freedom of QCD implies that when quarks and gluons exchange momenta p much larger than Λ_{QCD} , the process is perturbative in the strong coupling constant $\alpha_s(p)$. On the other hand, the typical momenta exchanged by the light degrees of freedom with each other and with the heavy quark, are of order Λ_{QCD} , for which a perturbative expansion is of no use. For these exchanges, however, $p < m_Q$, and the heavy quark Q does not recoil, remaining at rest in the rest frame of the hadron. In this limit, Q acts as a static source of electric and chromoelectric gauge fields. The chromoelectric field, which holds H_Q together, is nonperturbative in nature, but it is independent of m_Q . The result is that the properties of the light degrees of freedom depend only on the presence of the static gauge field, independent of the flavor and mass of the heavy quark carrying the gauge charge.*

There is an immediate implication for the spectroscopy of heavy hadrons. Since the interaction of the light degrees of freedom with the heavy quark is independent of m_Q , then so is the spectrum of their excitations. It is these excitations which determine the spectrum of heavy hadrons H_Q . Hence, the *splittings* $\Delta_i \sim \Lambda_{\text{QCD}}$ between the various hadrons H_Q^i are independent of Q , and in the limit $m_Q \rightarrow \infty$, do not scale with m_Q .

For example, the spectra of bottom and charmed mesons are shown schematically in Fig. 1, in the limit $m_b, m_c \gg \Lambda_{\text{QCD}}$. The light degrees of freedom are in exactly the same state in the mesons B_i and D_i , for a given i . The offset $B_i - D_i = m_b - m_c$ is just the difference between the heavy quark masses; in no way does the relationship between the spectra rely on an approximation $m_b \approx m_c$.

We can enrich this picture by recalling that the heavy quarks and light degrees of freedom also carry spin. The heavy quark has spin quantum number $S_Q = \frac{1}{2}$, which leads to a chromomagnetic moment

$$\mu_Q = \frac{g}{2m_Q}. \quad (4)$$

Note that $\mu_Q \rightarrow 0$ as $m_Q \rightarrow \infty$, and the interaction between the spin of the heavy quark and the light degrees of freedom is suppressed. Hence, the light degrees of freedom are insensitive to S_Q ; their state is independent of whether $S_Q^z = \frac{1}{2}$ or

*Top quarks decay too quickly for a static chromoelectric field to be established around them, so the simplifications discussed here are not relevant to them.




Figure 1: Schematic spectra of the bottom and charmed mesons in the limit $m_b, m_c \gg \Lambda_{\text{QCD}}$. The offset of the two spectra is not to scale; in reality, $m_b - m_c \gg \Delta_i \sim \Lambda_{\text{QCD}}$.

$S_Q^z = -\frac{1}{2}$. Thus each of the energy levels in Fig. 1 is actually doubled, one state for each possible value of S_Q^z .

To summarize, what we see is that the light degrees of freedom are the same when combined with any of the following heavy quark states:

$$Q_1(\uparrow), Q_1(\downarrow); Q_2(\uparrow), Q_2(\downarrow); \dots Q_{N_h}(\uparrow), Q_{N_h}(\downarrow), \quad (5)$$

where there are N_h heavy quarks (in the real world, $N_h = 2$). The result is an $SU(2N_h)$ symmetry which applies to the light degrees of freedom.³⁻⁷ A new symmetry means new nonperturbative relations between physical quantities. It is these relations which we wish to understand and exploit.

The light degrees of freedom have total angular momentum J_ℓ , which is integral for baryons and half-integral for mesons. When combined with the heavy quark spin $S_Q = \frac{1}{2}$, we find physical hadron states with total angular momentum

$$J = |J_\ell \pm \frac{1}{2}|. \quad (6)$$

If $S_\ell \neq 0$, then these two states are degenerate. For example, the lightest heavy mesons have $S_\ell = \frac{1}{2}$, leading to a doublet with $J = 0$ and $J = 1$. Indeed, in the

charm system, we find that the states of lowest mass are the spin-0 D and the spin-1 D^* ; the corresponding bottom mesons are the B and B^* .

When effects of order $1/m_Q$ are included, the chromomagnetic interactions split the states of given S_ℓ but different J . This “hyperfine” splitting is not calculable perturbatively, but it is proportional to the heavy quark magnetic moment μ_Q . This gives its scaling with m_Q :

$$\begin{aligned} m_{D^*} - m_D &\sim 1/m_c \\ m_{B^*} - m_B &\sim 1/m_b. \end{aligned} \quad (7)$$

From this fact, we can construct a relation which is a nonperturbative prediction of heavy quark symmetry,

$$m_{B^*}^2 - m_B^2 = m_{D^*}^2 - m_D^2. \quad (8)$$

Experimentally, $m_{B^*}^2 - m_B^2 = 0.49 \text{ GeV}^2$ and $m_{D^*}^2 - m_D^2 = 0.55 \text{ GeV}^2$, so this prediction works quite well. Note that this relation involves not just the heavy quark symmetry, but the systematic inclusion of the leading symmetry violating effects.

So far, we have formulated heavy quark symmetry for hadrons in their rest frame. Of course, we can easily boost to a frame in which the hadrons have arbitrary four-velocity $v^\mu = \gamma(1, \vec{v})$. For heavy quarks Q_1 and Q_2 , the symmetry will then relate hadrons $H_1(v)$ and $H_2(v)$ with the same velocity but with different momenta. This distinguishes heavy quark symmetry from ordinary symmetries of QCD, which relate states of the same momentum. To remind ourselves of this distinction, henceforth we will label heavy hadrons explicitly by their velocity: $D(v)$, $D^*(v)$, $B(v)$, $B^*(v)$, and so on.

1.3 Semileptonic Decay of a Heavy Quark

Now let us return to the semileptonic weak decay $b \rightarrow c \ell \bar{\nu}$, but now consider it in the heavy quark limit for the b and c quarks. Suppose the decay occurs at time $t = 0$. For $t < 0$, the b quark is embedded in a hadron H_b ; for $t > 0$, the c quark is dressed by light degrees of freedom to H_c' . Let us consider the lightest hadrons, $H_b = B(v)$ and $H_c' = D(v')$. Note that since the leptons carry away energy and momentum, in general $v \neq v'$.

What happens to the light degrees of freedom when the heavy quark decays? For $t < 0$, they see the chromoelectric field of a point source with velocity v . At $t = 0$, this point source recoils instantaneously[†] to velocity v' ; the color-neutral leptons do not interact with the light hadronic degrees of freedom as they fly off. The light quarks and gluons then must reassemble themselves about the recoiling color source. This nonperturbative process will generally involve the production of an excited state or of additional particles; the light degrees of freedom can exchange energy with the heavy quark, so there is no kinematic restriction on the excitations (of energy $\sim \Lambda_{\text{QCD}}$) which can be formed. There is also some chance that the light degrees of freedom will reassemble themselves back into a ground-state D meson. The amplitude for this to happen is a function only of the inner product $w = v \cdot v'$ of the initial and final velocities of the color sources. This amplitude, $\xi(w)$, is known as the Isgur-Wise function.³

Clearly, the kinematic point $v = v'$, or $w = 1$, is a special one. In this corner of phase space, where the leptons are emitted back to back, there is no recoil of the source of color field at $t = 0$. As far as the light degrees of freedom are concerned, *nothing happens!* Their state is unaffected by the decay of the heavy quark; they don't even notice it. Hence, the amplitude for them to remain in the ground state is exactly unity. This is reflected in a nonperturbative normalization of the Isgur-Wise function at zero recoil,³

$$\xi(1) = 1. \quad (9)$$

As we will see, this normalization condition is of enormous phenomenological use. It will be extremely important to understand the corrections to this result for finite heavy quark masses m_b , and especially, m_c .

The weak decay $b \rightarrow c$ is mediated by a left-handed current $\bar{c}\gamma^\mu(1 - \gamma^5)b$. Not only does this operator carry momentum, but it can change the orientation of the spin S_Q of the heavy quark during the decay. For a fixed light angular momentum J_ℓ , the relative orientation of S_Q determines whether the physical hadron in the final state is a D or a D^* . However, the light degrees of freedom are insensitive to S_Q , so the nonperturbative part of the transition is the same whether it is a D or a D^* which is produced. Hence, heavy quark symmetry implies relations between the hadronic matrix elements which describe the semileptonic decays $B \rightarrow D\ell\bar{\nu}$ and $B \rightarrow D^*\ell\bar{\nu}$.

[†]The weak decay occurs over a very short time $\delta t \sim 1/M_W \ll 1/\Lambda_{\text{QCD}}$.

It is conventional to parameterize these matrix elements by a set of scalar form factors. These are defined separately for the vector and axial currents, as follows:

$$\begin{aligned}
(D(v') | \bar{c} \gamma^\mu b | B(v)) &= h_+(w)(v + v')^\mu + h_-(w)(v - v')^\mu \\
(D^*(v', \epsilon) | \bar{c} \gamma^\mu b | B(v)) &= h_V(w) i \epsilon^{\mu\alpha\beta} \epsilon_\nu^* v'_\alpha v_\beta \\
(D(v') | \bar{c} \gamma^\mu \gamma^5 b | B(v)) &= 0 \\
(D(v', \epsilon) | \bar{c} \gamma^\mu \gamma^5 b | B(v)) &= h_{A_1}(w)(w + 1) \epsilon^{*\mu} - \epsilon^* \cdot v [h_{A_2}(w)v^\mu + h_{A_3}(w)v'^\mu].
\end{aligned} \tag{10}$$

The set of form factors $h_i(w)$ is the one appropriate to the heavy quark limit. Other linear combinations are also found in the literature. In any case, the form factors are independent nonperturbative functions of the recoil, or equivalently, for fixed m_b and m_c , of the momentum transfer. However, in the heavy quark limit, they correspond to a *single* transition of the light degrees of freedom, being distinguished from each other only by the relative orientation of the spin of the heavy quark. Hence, they may all be written in terms of the single function $\xi(w)$ which describes this nonperturbative transition. As we will derive later, the result is a set of relations,³

$$\begin{aligned}
h_+(w) &= h_V(w) = h_{A_1}(w) = h_{A_2}(w) = \xi(w) \\
h_-(w) &= h_{A_3}(w) = 0,
\end{aligned} \tag{11}$$

which follow solely from the heavy quark symmetry. Of course, all of the form factors which do not vanish inherit the normalization condition (9) at zero recoil. This result is a powerful constraint on the structure of semileptonic decay in the heavy quark limit.

1.4 Heavy Meson Decay Constant

As a final example of the utility of the heavy quark limit, consider the coupling of the heavy meson field to the axial vector current. This is conventionally parameterized in terms of a decay constant; for example, for the B^- meson, we define f_B via

$$\langle 0 | \bar{u} \gamma^\mu \gamma^5 b | B^-(p_B) \rangle = i f_B p_B^\mu. \tag{12}$$

What is the dependence of the nonperturbative quantity f_B on m_B ? To address this question, we rewrite Eq. (12) in a form appropriate to taking the heavy quark limit, $m_B \rightarrow \infty$ (which is equivalent to $m_b \rightarrow \infty$). This entails making explicit

the dependence of all quantities on m_B . First, we trade the B^- momentum for its velocity,

$$p_B^\mu = m_B v^\mu. \tag{13}$$

Second, we replace the usual B^- state, whose normalization depends on m_B ,

$$\langle B(p_1) | B(p_2) \rangle = 2E_B \delta^{(3)}(\vec{p}_1 - \vec{p}_2), \tag{14}$$

by a mass-independent state,

$$|B(v)\rangle = \frac{1}{\sqrt{m_B}} |B(p_B)\rangle, \tag{15}$$

satisfying

$$\langle B(v_1) | B(v_2) \rangle = 2\gamma \delta^{(3)}(\vec{p}_1 - \vec{p}_2). \tag{16}$$

Then Eq. (12) becomes

$$\sqrt{m_B} \langle 0 | \bar{u} \gamma^\mu \gamma^5 b | B^-(v) \rangle = i f_B m_B v^\mu. \tag{17}$$

The nonperturbative matrix element $\langle 0 | \bar{u} \gamma^\mu \gamma^5 b | B^-(v) \rangle$ is independent of m_B in the heavy quark limit. Hence, we see that in this limit, f_B takes the form

$$f_B = m_B^{-1/2} \times (\text{independent of } m_B). \tag{18}$$

This makes explicit the scaling of f_B with m_B . It is more interesting to write this as a prediction for the ratio of charmed and bottom meson decay constants. We find^{3,4,6}

$$\frac{f_B}{f_D} = \sqrt{\frac{m_D}{m_B}} + \mathcal{O}\left(\frac{\Lambda_{\text{QCD}}}{m_D}, \frac{\Lambda_{\text{QCD}}}{m_B}\right). \tag{19}$$

For the physical bottom and charm masses, of course, the correction terms proportional to Λ_{QCD}/m_Q could be important.

2 Heavy Quark Effective Theory

We have already extracted quite a bit of nontrivial information from the heavy quark limit. We have found the scaling of various quantities with m_Q , we have studied the implications for heavy hadron spectroscopy, and we have found nonperturbative relations among the hadronic form factors which describe semileptonic $b \rightarrow c$ decay. However, all of these results have been obtained in the strict

limit $m_Q \rightarrow \infty$. If the heavy quark limit is to be of more than academic interest and is to provide the basis for quantitative phenomenology, we have to understand how to include corrections systematically. There are actually two types of corrections which we would like to include. *Power corrections* are subleading terms in the expansion in Λ_{QCD}/m_Q ; those proportional to Λ_{QCD}/m_c are the most worrisome, because of the relatively small charm quark mass. *Logarithmic corrections* arise from the implicit dependence of quantities on m_Q through the strong coupling constant $\alpha_s(m_Q) \sim 1/\ln(m_Q/\Lambda_{\text{QCD}})$. For the physical values of m_b and m_c , either of these could be important. What we need is a formalism which can accommodate them both.

In short, we need to go from a set of heavy quark symmetry predictions in the $m_Q \rightarrow \infty$ limit to a reformulation of QCD which provides a controlled expansion about this limit. The formalism which does the job is the Heavy Quark Effective Theory, or the HQET. The purpose of the HQET is to allow us to extract, explicitly and systematically, all dependence of physical quantities on m_Q , in the limit $m_Q \gg \Lambda_{\text{QCD}}$. In these lectures, we will develop only enough of the technology to treat the dominant leading effects, providing indications along the way of how one would carry the expansion further.

The HQET, as formulated here, was developed in a series of papers going back to the late 1980s (Refs. 3, 4, 6, and 8–15), which the reader who is interested in tracking its historical development may consult.

2.1 The Effective Lagrangian

Consider the kinematics of a heavy quark Q , bound in a hadron with light degrees of freedom to make a color singlet state. The small momenta which Q typically exchanges with the rest of the hadron are of order $\Lambda_{\text{QCD}} \ll m_Q$, and they never take Q far from its mass shell, $p_Q^2 = m_Q^2$. Hence, the momentum p_Q^μ can be decomposed into two parts,

$$p_Q^\mu = m_Q v^\mu + k^\mu, \quad (20)$$

where $m_Q v^\mu$ is the constant on-shell part of p_Q^μ , and $k^\mu \sim \Lambda_{\text{QCD}}$ is the small, fluctuating “residual momentum.” The on-shell condition for the heavy quark then becomes

$$m_Q^2 = (m_Q v^\mu + k^\mu)^2 = m_Q^2 + 2m_Q v \cdot k + k^2. \quad (21)$$

In the heavy quark limit, we may neglect the last term, and we have the simple condition

$$v \cdot k = 0 \quad (22)$$

for an on-shell heavy quark. Here, the velocity v^μ functions as a label; since soft interactions cannot change v^μ , there is a *velocity superselection rule* in the heavy quark limit, and v^μ is a good quantum number of the QCD Hamiltonian.

We find the same result by taking the $m_Q \rightarrow \infty$ limit of the heavy quark propagator,

$$\frac{i}{\not{p} - m_Q + i\epsilon} \rightarrow \frac{1 + \not{v}}{2} \frac{i}{v \cdot k + i\epsilon}. \quad (23)$$

In this limit, the propagator is independent of m_Q . The projection operators

$$P_\pm = \frac{1 \pm \not{v}}{2} \quad (24)$$

project onto the positive (P_+) and negative (P_-) frequency parts of the Dirac field Q . This is clear in the Dirac representation in the rest frame, in which P_+ and P_- project, respectively, onto the upper two and lower two components of the heavy quark spinor. In the limit $m_Q \rightarrow \infty$, in which Q remains almost on shell, only the “large” upper components of the field Q propagate; mixing via zitterbewegung with the “small” lower components is suppressed by $1/2m_Q$. Hence, the action of the projectors on Q is

$$P_+ Q(x) = Q(x) + O(1/m_Q), \quad P_- Q(x) = 0 + O(1/m_Q). \quad (25)$$

The momentum dependence of the field Q is given by its action on a heavy quark state,

$$Q(x) |Q(p)\rangle = e^{-ip \cdot x} |0\rangle. \quad (26)$$

If we now multiply both sides by a phase corresponding to the on-shell momentum,

$$e^{im_Q v \cdot x} Q(x) |Q(p)\rangle = e^{-ik \cdot x} |0\rangle, \quad (27)$$

the right side of this equation is independent of m_Q . Hence, the left side must be, as well. Combining this observation with the argument of the previous paragraph, we are motivated to define a m_Q -independent effective heavy quark field $h_v(x)$,

$$h_v(x) = e^{im_Q v \cdot x} P_+ Q(x). \quad (28)$$

Note that the effective field carries a velocity label v and is a two-component object. The modifications to the ordinary field $Q(x)$ project out the positive

frequency part and ensure that states annihilated by $h_v(x)$ have no dependence on m_Q . Hence, these are reasonable candidate fields to carry representations of the heavy quark symmetry. Of course, the small components cannot be neglected when effects of order $1/m_Q$ are included. In the HQET, they are represented by a field

$$H_v(x) = e^{im_Q v \cdot x} P_- Q(x). \quad (29)$$

The field $H_v(x)$ vanishes in the $m_Q \rightarrow \infty$ limit.

The ordinary QCD Lagrange density for a field $Q(x)$ is given by

$$\mathcal{L}_{\text{QCD}} = \bar{Q}(x) (i\not{D} - m_Q) Q(x), \quad (30)$$

where $D_\mu = \partial_\mu - igA_\mu^a T^a$ is the gauge covariant derivative. To find the Lagrangian of the HQET, we substitute

$$Q(x) = e^{-im_Q v \cdot x} h_v(x) + \dots \quad (31)$$

into \mathcal{L}_{QCD} and expand. With the aid of the projection identity $P_+ \gamma^\mu P_+ = v^\mu$, we find

$$\mathcal{L}_{\text{HQET}} = \bar{h}_v(x) i v \cdot D h(x). \quad (32)$$

This simple Lagrangian leads to the propagator we derived earlier,

$$\frac{i}{v \cdot k + i\epsilon}, \quad (33)$$

and to an equally simple quark-gluon vertex,

$$igT^a v^\mu A_\mu^a. \quad (34)$$

Note that both the propagator and the vertex are independent of m_Q , reflecting the heavy quark flavor symmetry. They also have no Dirac structure, reflecting the heavy quark spin symmetry. Our intuitive statements about the structure of heavy hadrons have been promoted to explicit symmetries of the QCD Lagrangian in the limit $m_Q \rightarrow \infty$.

It is straightforward to include power corrections to $\mathcal{L}_{\text{HQET}}$. Write $Q(x)$ in terms of the effective fields,

$$Q(x) = e^{-im_Q v \cdot x} [h_v(x) + H_v(x)], \quad (35)$$

and apply the classical equation of motion $(i\not{D} - m_Q)Q(x) = 0$:

$$i\not{D} h_v(x) + (i\not{D} - 2m_Q)H_v(x) = 0. \quad (36)$$

Multiplying by P_- and commuting \not{D} to the right, we find

$$(iv \cdot D + 2m_Q)H_v(x) = i\not{D}_\perp h_v(x), \quad (37)$$

where $D_\perp^\mu = D^\mu - v^\mu v \cdot D$. We then substitute $Q(x)$ into \mathcal{L}_{QCD} as before, eliminate $H_v(x)$, and expand in $1/m_Q$ to obtain

$$\begin{aligned} \mathcal{L}_{\text{HQET}} &= \bar{h}_v i v \cdot D h_v + \bar{h}_v i\not{D}_\perp \frac{1}{iv \cdot D + 2m_Q} i\not{D}_\perp h_v \\ &= \bar{h}_v i v \cdot D h_v + \frac{1}{2m_Q} \left[\bar{h}_v (iD_\perp)^2 h_v + \frac{g}{2} \bar{h}_v \sigma^{\alpha\beta} G_{\alpha\beta} h_v \right] + \dots \end{aligned} \quad (38)$$

The leading corrections have a simple interpretation, which becomes clear in the rest frame, $v^\mu = (1, 0, 0, 0)$. The spin-independent term is

$$\frac{1}{2m_Q} \mathcal{O}_K = \frac{1}{2m_Q} \bar{h}_v (iD_\perp)^2 h_v \rightarrow -\frac{1}{2m_Q} \bar{h}_v (i\vec{D})^2 h_v, \quad (39)$$

which is the negative of the nonrelativistic kinetic energy of the heavy quark. Because of the explicit factor of $1/2m_Q$, this term violates the heavy flavor symmetry. The spin-dependent part is

$$\frac{1}{2m_Q} \mathcal{O}_G = \frac{1}{2m_Q} \frac{g}{2} \bar{h}_v \sigma^{\alpha\beta} G_{\alpha\beta} h_v \rightarrow \frac{1}{4m_Q} \bar{h}_v \sigma^{ij} T^a h_v \times g G_{ij}^a = g\vec{\mu}_Q^a \cdot \vec{B}^a, \quad (40)$$

which is the coupling of the spin of the heavy quark to the chromomagnetic field. Because it has a nontrivial Dirac structure, this term violates both the heavy flavor symmetry and the heavy spin symmetry. For example, \mathcal{O}_G is responsible for the $D - D^*$ and $B - B^*$ mass splittings. These correction terms will be treated as part of the *interaction* Lagrangian, even though \mathcal{O}_K has a piece which is a pure bilinear in the heavy quark field.

2.2 Effective Currents and States

The expansion of the weak interaction current $\bar{c}\gamma^\mu(1-\gamma^5)b$ is analogous. However, here we must introduce separate effective fields for the charm and bottom quarks, each with its own velocity:

$$b \rightarrow h_v^b, \quad c \rightarrow h_v^c. \quad (41)$$

Then a general flavor-changing current becomes, to leading order,

$$\bar{c}\Gamma b \rightarrow \bar{h}_v^c \Gamma h_v^b, \quad (42)$$

where Γ is a fixed Dirac structure. With the leading power corrections, this is

$$\bar{e}\Gamma b \rightarrow \bar{h}_v^c \Gamma h_v^b + \frac{1}{2m_b} \bar{h}_v^c \Gamma (i\not{D}_\perp) h_v^b + \frac{1}{2m_c} \bar{h}_v^c (-i\not{D}_\perp) \Gamma h_v^b + \dots \quad (43)$$

The effective currents, and other operators which appear in the HQET, may often be simplified by use of the classical equation of motion,

$$iv \cdot D h_v(x) = 0. \quad (44)$$

However, it is only safe to apply these equations naively at order $1/m_Q$; at higher order, the application of the equations of motion involves additional subtleties.¹⁶⁻¹⁸

To complete the effective theory, we need m_Q -independent hadron states which are created and annihilated by currents containing the effective fields. For example, there is an effective pseudoscalar meson state $|M(v)\rangle$ which couples to the effective axial current $\bar{q}\gamma^\mu\gamma^5 h_v$, with a coupling F_M which is independent of m_Q ,

$$\langle 0 | \bar{q}\gamma^\mu\gamma^5 h_v | M(v)\rangle = iF_M v^\mu. \quad (45)$$

At lowest order, F_M is related to a conventional decay constant such as f_B by

$$f_B = F_M / \sqrt{m_B}, \quad (46)$$

from which we immediately find the relationship [Eq. (19)] between f_D and f_B .

2.3 Radiative Corrections

We can use the effective Lagrangian $\mathcal{L}_{\text{HQET}}$ to compute the radiative corrections to the matrix element [Eq. (45)]. In particular, we would like to extract the dependence of F_M on $\ln m_Q$. This dependence comes through the one-loop renormalization of the current $\bar{q}\gamma^\mu\gamma^5 Q$. At lowest order, of course, the renormalization is straightforward: we simply compute the set of graphs found in Fig. 2. The result is finite, because the current is (partially) conserved, and we extract a result of the form

$$\bar{q}\gamma^\mu\gamma^5 Q \left(1 - \bar{\gamma}_0 \frac{\alpha_s}{\pi} \ln(m_Q/m_q) + \dots \right). \quad (47)$$

Note that there is no explicit dependence on the renormalization scale μ , since there is no divergence to be subtracted.

The same result may be obtained in the effective theory. In this case, we must match the currents in full QCD onto HQET currents of the form $\bar{q}\gamma^\mu\gamma^5 h_v$.




Figure 2: Tree level plus the one-loop renormalization of the current $\bar{q}\Gamma Q$ in QCD. The box represents the current insertion.

This step will induce a matching coefficient containing the explicit dependence on m_Q , which is absent, by construction, from the operators and Lagrangian of the HQET.[†] In addition, the effective current will not necessarily be conserved, since the ultraviolet properties of QCD and the HQET differ. Hence, the form of the matching, once radiative corrections are included, is

$$\bar{q}\gamma^\mu\gamma^5 Q \rightarrow C(m_Q, \mu) \bar{q}\gamma^\mu\gamma^5 h_v. \quad (48)$$

We can deduce the form of $C(m_Q, \mu)$ by considering the renormalization of the effective current $\bar{q}\gamma^\mu\gamma^5 h_v$, shown in the last three terms of Fig. 3. These diagrams, computed in the effective theory, are independent of m_Q . However, in general they are divergent, so they depend on the renormalization scale μ ; the renormalization takes the form

$$\bar{q}\gamma^\mu\gamma^5 h_v \left(C(m_Q, \mu) - \gamma_0 \frac{\alpha_s}{\pi} \ln(\mu/m_q) + \dots \right), \quad (49)$$

where m_q acts as an infrared cutoff. The μ dependence in the second term must be canceled by $C(m_Q, \mu)$. Since the logarithm depends on a dimensionless ratio, $C(m_Q, \mu)$ must be of the form

$$C(m_Q, \mu) = 1 - \gamma_0 \frac{\alpha_s}{\pi} \ln(m_Q/\mu) + \dots \quad (50)$$

Comparing the dependence on m_Q of $C(m_Q, \mu)$ and the expansion [Eq. (47)], we see that $\bar{\gamma}_0 = \gamma_0$.

However, the effective theory allows us to go beyond leading order and to resum all corrections of the form $\alpha_s^n \ln^n m_Q$. We do this with the renormalization group equations, which express the independence of physical observables on the

[†]In general, the matching procedure at order α_s can also induce new Dirac structures $\bar{q}\Gamma h_v$. They do not affect the leading logarithms discussed here.




Figure 3: Tree level plus the one-loop renormalization of the current $\bar{q}\gamma^\mu\gamma^5 h_\nu$ in HQET. The double line is the propagator of the effective field h_ν .

renormalization scale μ . In this case, they require that the μ dependence of $C(m_Q, \mu)$ cancel that of the one-loop diagrams in Fig. 3, under small changes in μ :

$$\mu \frac{d}{d\mu} C(m_Q, \mu) = \gamma_0 \frac{\alpha_s(\mu)}{\pi}. \quad (51)$$

Here, the total derivative with respect to μ includes the implicit dependence on μ of the coupling constant $\alpha_s(\mu)$:

$$\begin{aligned} \mu \frac{d}{d\mu} &= \mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g} \\ \beta(g) &= -\beta_0 \frac{g^3}{16\pi^2} + \dots \\ \beta_0 &= 11 - \frac{2}{3} N_f = \frac{25}{3} \text{ for } N_f = 4, \end{aligned} \quad (52)$$

where N_f is the number of light flavors. We compute the anomalous dimension γ_0 from the divergent parts of the one-loop diagrams shown in Fig. 3. From the three terms we find, respectively,^{4,6}

$$\gamma_0 = \frac{8}{3} + \frac{1}{2} \left(-\frac{8}{3} \right) + \frac{1}{2} \left(\frac{16}{3} \right) = 4. \quad (53)$$

The solution of the renormalization group equation is

$$C(m_Q, \mu) = \left(\frac{\alpha_s(m_Q)}{\alpha_s(\mu)} \right)^{\gamma_0/2\beta_0} = \left(\frac{\alpha_s(m_Q)}{\alpha_s(\mu)} \right)^{6/25}. \quad (54)$$

This then yields the leading logarithmic correction to the ratio f_B/f_D :

$$\frac{f_B}{f_D} = \sqrt{\frac{m_D}{m_B}} \left(\frac{\alpha_s(m_b)}{\alpha_s(m_c)} \right)^{6/25}. \quad (55)$$

The radiative correction is approximately a ten percent effect. In fact, it has a simple physical interpretation. For virtual gluons of “intermediate” energy,

$m_c < E_g < m_b$, the bottom quark is heavy but the charm quark is light. Such gluons contribute to the difference between f_B and f_D even in the heavy quark limit.

In summary, then, the purpose of the HQET is to make explicit all dependence of observable quantities on m_Q . The logarithmic dependence, through $\alpha_s(m_Q)$, arises from intermediate virtual gluons with $m_c < E_g < m_b$. We obtain these corrections by computing perturbatively with the HQET Lagrangian, then using the renormalization group to resum the logarithms to all orders. The power dependence, $1/m_Q$, is extracted systematically in the heavy quark expansion. We have seen how to expand the Lagrangian and the states to subleading order; the application of the expansion to a physical decay rate will be presented in the next section.

These lectures are meant to be pedagogical, so we will only treat the leading corrections to a few processes. However, the state of the art goes significantly beyond what will be presented here. For many quantities, not only the leading logarithms, $\alpha_s^n \ln^n m_Q$, but the subleading (two-loop) logarithms, of order $\alpha_s^{n+1} \ln^n m_Q$, have been resummed. Similarly, many power corrections are known to relative order $1/m_Q^2$. It is particularly important phenomenologically to take into account the corrections of order $1/m_c^2$.

3 Exclusive B Decays

We now have the tools we need for an HQET treatment of the exclusive semileptonic transitions $B \rightarrow D \ell \bar{\nu}$ and $B \rightarrow D^* \ell \bar{\nu}$. Earlier, we argued on physical grounds that in the heavy quark limit, all of the hadronic matrix elements which appear in these decays are related to a single nonperturbative function $\xi(w)$. Now we will sharpen this analysis to actually *derive* these relations, and to include radiative and power corrections. In fact, almost all of our effort will go into the power corrections, since the radiative corrections to the transition currents are computed just as in the previous section.

3.1 Matrix Element Relations at Leading Order

The transitions in question require the nonperturbative matrix elements

$$(D(v')|\bar{c}\gamma^\mu b|B(v)), \quad (D^*(v',\epsilon)|\bar{c}\gamma^\mu b|B(v)), \quad (D^*(v',\epsilon)|\bar{c}\gamma^\mu\gamma^5 b|B(v)), \quad (56)$$

parameterized in terms of form factors as in Eq. (10). Our first task is to derive the relations between these form factors, as promised earlier. These relations depend on the heavy quark symmetry, that is, on the fact that the spin quantum numbers of Q and of the light degrees of freedom are separately conserved by the soft physics. Hence, we need a representation of the heavy meson states in which they have well-defined transformations separately under the angular momentum operators S_Q and J_L . In particular, the representation must reflect the fact that a rotation by S_Q can exchange the pseudoscalar meson $M(v)$ with the vector meson $M^*(v,\epsilon)$.

The solution is to introduce a “superfield” $\mathcal{M}(v)$, defined as the 4×4 Dirac matrix^{11,19}

$$\mathcal{M}(v) = \frac{1 + \not{v}}{2} [\gamma^\mu M_\mu^*(v, \epsilon) - \gamma^5 M(v)]. \quad (57)$$

Under heavy quark spin rotations S_Q , $\mathcal{M}(v)$ transforms as

$$\mathcal{M}(v) \rightarrow D(S_Q)\mathcal{M}(v), \quad (58)$$

and under Lorentz rotations Λ , as

$$\mathcal{M}(v) \rightarrow D(\Lambda)\mathcal{M}(\Lambda^{-1}v)D^{-1}(\Lambda). \quad (59)$$

Here $D(\dots)$ is the spinor representation of $SO(3,1)$. The superfield satisfies the matrix identity

$$P_+ \mathcal{M}(v) P_- = \mathcal{M}(v), \quad (60)$$

so it transforms the same way as the product of spinors $h_v \bar{q}$, representing a heavy quark and a light antiquark moving together at velocity v^μ .

A current which mediates the decay of one heavy quark (Q) into another (Q') is of the form $\bar{h}_{v'} \Gamma h_v$. Under a rotation by S_Q , the effective field h_v transforms as

$$\bar{h}_v \rightarrow D(S_Q) \bar{h}_v, \quad (61)$$

while $h_{v'}$ is unchanged. The current would remain invariant if we took Γ to transform as

$$\Gamma \rightarrow \Gamma D^{-1}(S_Q). \quad (62)$$

On the other hand, the matrix element of superfields

$$\langle \mathcal{M}'(v') | \bar{h}_{v'} \Gamma h_v | \mathcal{M}(v) \rangle \quad (63)$$

is invariant if we rotate *both* h_v and $\mathcal{M}(v)$ by the *same* S_Q . With the transformation law [Eq. (58)] for $\mathcal{M}(v)$, it follows that the S_Q -invariant matrix element must be proportional to $\Gamma \mathcal{M}(v)$. When we also consider rotations under $S_{Q'}$, we find that the matrix element is restricted to the general form

$$\langle \mathcal{M}'(v') | \bar{h}_{v'} \Gamma h_v | \mathcal{M}(v) \rangle = -[M_M M_{M'}]^{1/2} \text{Tr} [\bar{\mathcal{M}}'(v') \Gamma \mathcal{M}(v) \hat{F}(v, v')]. \quad (64)$$

The product of masses in front is a convention which restores the relativistic normalization of the states. Note that the heavy quark symmetry allows an arbitrary 4×4 Dirac matrix $\hat{F}(v, v')$ to act on the “light quark” part of the superfields. Its presence reflects the fact that, other than the constraint of Lorentz symmetry, the behavior of the spin of the light degrees of freedom during the decay is unknown.

A general expansion of $\hat{F}(v, v')$ in terms of scalar functions $F_i(w = v \cdot v')$ takes the form

$$\hat{F}(v, v') = F_1(w) + F_2(w)\not{v} + F_3(w)\not{v}' + F_4(w)\not{v}\not{v}'. \quad (65)$$

However, the identity [Eq. (60)] applied to the matrix element [Eq. (64)] yields

$$\hat{F}(v, v') = P_- \hat{F}(v, v') P_- = [F_1(w) - F_2(w) - F_3(w) + F_4(w)] P_- P_-'. \quad (66)$$

In other words, $F(v, v')$ is actually a *scalar*, which we identify with the Isgur-Wise function,

$$\hat{F}(v, v') = \xi(w). \quad (67)$$

As an exercise, let us apply this formalism to the matrix elements for $B \rightarrow (D, D^*) \ell \bar{\nu}$. For a given matrix element, we pick out the part of the superfield $\mathcal{M}(v)$ which is relevant. Hence, we find

$$\begin{aligned} \langle D(v') | \bar{c}\gamma^\mu b | B(v) \rangle &= (M_D(v') | \bar{h}_{v'}^c \gamma^\mu h_v^b | M_B(v) \rangle \\ &= -[m_D m_B]^{1/2} \text{Tr} [\gamma^5 P_+^c \gamma^\mu P_+ (-\gamma^5)] \xi(w) \\ &= [m_D m_B]^{1/2} \xi(w) (v + v')^\mu \end{aligned} \quad (68)$$

$$\begin{aligned}
\langle D^*(v', \epsilon) | \bar{c} \gamma^\mu \gamma^5 b | B(v) \rangle &= \langle M_D^*(v', \epsilon) | \bar{h}_v^c \gamma^\mu \gamma^5 h_v^b | M_B(v) \rangle \\
&= -[m_D \cdot m_B]^{1/2} \text{Tr} \left[\not{P}'_+ \gamma^\mu \gamma^5 P_+ (-\gamma^5) \right] \xi(w) \\
&= [m_D \cdot m_B]^{1/2} \xi(w) [(w+1)\epsilon^\mu - \epsilon^\nu \cdot (v+v')^\mu] \quad (69)
\end{aligned}$$

$$\begin{aligned}
\langle D^*(v', \epsilon) | \bar{c} \gamma^\mu b | B(v) \rangle &= \langle M_D^*(v', \epsilon) | \bar{h}_v^c \gamma^\mu h_v^b | M_B(v) \rangle \\
&= -[m_D \cdot m_B]^{1/2} \text{Tr} \left[\not{P}'_+ \gamma^\mu P_+ (-\gamma^5) \right] \xi(w) \\
&= [m_D \cdot m_B]^{1/2} \xi(w) i \epsilon^{\mu\nu\alpha\beta} \epsilon_\nu' v_\alpha v_\beta, \quad (70)
\end{aligned}$$

reproducing explicitly the relations [Eq. (11)] between the independent form factors $h_i(w)$. We can also derive the normalization condition at $w = 1$. Consider the matrix element of the b number current $\bar{b} \gamma^\mu b$ between B -meson states. In QCD, the matrix element of this current is exactly normalized,

$$\langle B(v) | \bar{b} \gamma^\mu b | B(v) \rangle = 2p_B^\mu = 2m_B v^\mu. \quad (71)$$

But in HQET, we have

$$\begin{aligned}
\langle B(v) | \bar{b} \gamma^\mu b | B(v) \rangle &= \langle M_B(v) | \bar{h}_v \gamma^\mu h_v | M_B(v) \rangle \\
&= m_B \xi(v \cdot v) (v+v)^\mu \\
&= 2m_B v^\mu \xi(1). \quad (72)
\end{aligned}$$

Hence, the normalization condition at zero recoil,

$$\xi(1) = 1, \quad (73)$$

follows directly from the conservation of the heavy quark number current.

3.2 Power Corrections to the Matrix Elements

The matrix elements we have derived are computed in the strict limit $m_{b,c} \rightarrow \infty$. How are they affected by corrections of order $1/m_b$ and $1/m_c$? There are two sources of $1/m_Q$ corrections in the effective theory: the corrections [Eq. (43)] to the heavy quark currents, and the corrections [Eq. (38)] to the Lagrangian.

When radiative corrections are included, the expansion of the heavy quark current $\bar{c} \Gamma b$ in terms of HQET operators has a form which is somewhat more

general than Eq. (43),

$$\bar{c} \Gamma b \rightarrow a_0(\alpha_s) \bar{h}_v^c \Gamma h_v^b + \frac{a_1(\alpha_s)}{2m_b} \bar{h}_v^c \Gamma \not{D}_\alpha h_v^b + \frac{a_1'(\alpha_s)}{2m_c} \bar{h}_v^c (-i \overleftarrow{D}_\alpha) \Gamma h_v^b + \dots \quad (74)$$

The matrix elements of the power corrections are constrained by heavy quark symmetry in a manner completely analogous to the leading current. In terms of traces over the superfields, we have¹³

$$\langle \mathcal{M}'(v') | \bar{h}_v \Gamma^\alpha h_v | \mathcal{M}(v) \rangle = -[M_M M_{M'}]^{1/2} \text{Tr} \left[\overline{\mathcal{M}'}(v') \Gamma^\alpha \mathcal{M}(v) \hat{G}_\alpha(v, v') \right], \quad (75)$$

where $\hat{G}_\alpha(v, v')$ is another arbitrary 4×4 Dirac matrix. The matrix element

$$\langle \mathcal{M}'(v') | \bar{h}_v (-i \overleftarrow{D}_\alpha) \Gamma h_v | \mathcal{M}(v) \rangle \quad (76)$$

may also be written in terms of $\hat{G}_\alpha(v, v')$, using charge conjugation.

The $1/m_Q$ corrections \mathcal{O}_K and \mathcal{O}_G to the Lagrangian contribute somewhat differently. In order to apply heavy quark symmetry, the matrix elements of the local currents, both leading and subleading, must be written in terms of the *effective* states $|M(v)\rangle$. However, these states are not eigenstates of the Hamiltonian, once \mathcal{O}_K and \mathcal{O}_G are included in the Lagrangian. Hence, we must allow for the possibility that if an effective state $|M(v)\rangle$ is created at time $t = -\infty$, then \mathcal{O}_K or \mathcal{O}_G could act on the state before its decay at $t = 0$. This possibility is accounted for by including time-ordered products in which \mathcal{O}_K or \mathcal{O}_G is inserted along the incoming or outgoing heavy quark line. If we are keeping terms of order $1/m_Q$, only one insertion of \mathcal{O}_K or \mathcal{O}_G needs to be included. The time-ordered products are of the form¹³

$$\begin{aligned}
\langle D^{(*)}(v) | \bar{c} \Gamma b | B(v) \rangle &= \dots + \frac{1}{2m_c} \langle \mathcal{M}'(v') | i \int dy T \{ \bar{h}_v^c \Gamma h_v^b, \mathcal{O}_K^{v'} + \mathcal{O}_G^{v'} \} | \mathcal{M}(v) \rangle \\
&\quad + \frac{1}{2m_b} \langle \mathcal{M}'(v') | i \int dy T \{ \bar{h}_v^c \Gamma h_v^b, \mathcal{O}_K^v + \mathcal{O}_G^v \} | \mathcal{M}(v) \rangle, \quad (77)
\end{aligned}$$

where the ellipses refer to the current corrections computed earlier. The evaluation of the matrix elements of the time-ordered products will lead to still more nonperturbative functions like $\hat{F}(v, v')$ and $\hat{G}_\alpha(v, v')$.

3.3 Corrections at Zero Recoil

It is straightforward, but not very illuminating, to expand all of the new nonperturbative functions which arise at order $1/m_Q$ in terms of scalar form factors. In

the end, the corrections may be parameterized in terms of four functions of the velocity transfer w , and a single nonperturbative parameter $\bar{\Lambda}$, all proportional to the mass scale Λ_{QCD} . The new parameter has a simple interpretation as the “energy” of the light degrees of freedom and is given by

$$\bar{\Lambda} = \lim_{m_b \rightarrow \infty} (m_B - m_b). \quad (78)$$

Instead of a general treatment, however, we will consider the $1/m_Q$ corrections at the zero recoil point $w = 1$. This is clearly the most important case, because it is at this point that the nonperturbative matrix elements are absolutely normalized in the heavy quark limit. What happens to this normalization condition when $1/m_Q$ corrections are included?

Let us study the corrections to the current in detail. They are described by the nonperturbative function $\hat{G}_\alpha(v, v')$. At $v = v'$, $\hat{G}_\alpha(v, v)$ may be expanded as

$$\hat{G}_\alpha(v, v) = G_1 v_\alpha + G_2 \gamma_\alpha + G_3 v_\alpha \not{v} + G_4 \gamma_\alpha \not{v}. \quad (79)$$

But $\hat{G}_\alpha(v, v)$ is subject to the same constraint as $\hat{F}(v, v')$,

$$\hat{G}_\alpha(v, v) = P_- \hat{G}_\alpha(v, v) P_- = (G_1 - G_2 - G_3 + G_4) v_\alpha P_- \equiv G v_\alpha P_-, \quad (80)$$

and it, too, is equivalent to a Dirac scalar (the same is *not* true of the general function $\hat{G}_\alpha(v, v')$). Now consider the matrix element where we take $\Gamma^\alpha = v^\alpha$. Then we have

$$\begin{aligned} (\mathcal{M}'(v) | \bar{h}_v i v \cdot D h_v | \mathcal{M}(v)) &= - [M_M M_{M'}]^{1/2} \text{Tr} [\bar{\mathcal{M}}'(v) v^\alpha \mathcal{M}(v)] G v_\alpha \\ &= - G [M_M M_{M'}]^{1/2} \text{Tr} [\bar{\mathcal{M}}'(v) \mathcal{M}(v)]. \end{aligned} \quad (81)$$

But this matrix element *vanishes* by the classical equation of motion in the effective theory,

$$v \cdot D h_v(x) = 0. \quad (82)$$

Hence, $G = 0 = \hat{G}_\alpha(v, v)$. There are no $1/m_Q$ corrections from the current to the normalization condition at zero recoil.^{13,20}

The same is true of insertions of the corrections \mathcal{O}_K and \mathcal{O}_G to the Lagrangian: their contribution vanishes at $w = 1$. To show this requires the imposition of the conservation of the b number current at order $1/m_b$, much as we derived the normalization of the Isgur-Wise function at leading order. The complete derivation is found in the literature.¹³

In the end, we have an extremely exciting result, known as *Luke's Theorem*. There are no corrections at zero recoil to the hadronic matrix elements responsible for the semileptonic decays $B \rightarrow D \ell \bar{\nu}$ and $B \rightarrow D^* \ell \bar{\nu}$. The leading power corrections to the normalization of zero recoil matrix elements are only of order $1/m_c^2$. Given that $\Lambda_{\text{QCD}}/m_c \sim 30\%$ and $\Lambda_{\text{QCD}}^2/m_c^2 \sim 10\%$, the implication is that the leading order predictions at $w = 1$ are considerably more accurate than one might have expected. In addition, away from zero recoil, the $1/m_c$ corrections must be suppressed at least by $(w - 1)$.

On closer inspection, this result is more interesting for $B \rightarrow D^* \ell \bar{\nu}$ than for $B \rightarrow D \ell \bar{\nu}$. This is because the leading order matrix element for $B \rightarrow D \ell \bar{\nu}$ vanishes kinematically at zero recoil for a massless lepton in the final state. Hence, in this case, the $1/m_c$ corrections are not suppressed as a fractional correction to the lowest order term.²¹

3.4 Extraction of $|V_{cb}|$ from $B \rightarrow D^* \ell \bar{\nu}$

An immediate application of these results is the extraction of $|V_{cb}|$ from the exclusive decay $B \rightarrow D^* \ell \bar{\nu}$. This process is mediated by the weak operator \mathcal{O}_{bc} [Eq. (1)], whose matrix element factorizes as

$$\langle D^* \ell \bar{\nu} | \mathcal{O}_{bc} | B \rangle = \frac{G_F V_{cb}}{\sqrt{2}} \langle D^* | \bar{c} \gamma^\mu (1 - \gamma^5) b | B \rangle \langle \ell \bar{\nu} | \bar{\ell} \gamma_\mu (1 - \gamma^5) \nu | 0 \rangle. \quad (83)$$

The leptonic matrix element may be computed perturbatively, while we treat the hadronic matrix element in the heavy quark expansion. The result is a differential decay rate of the form²²

$$\begin{aligned} \frac{d\Gamma}{dw} &= \frac{G_F^2}{48\pi^3} |V_{cb}|^2 (m_B - m_{D^*})^2 m_{D^*}^3 (w + 1)^3 \sqrt{w^2 - 1} \\ &\quad \times \left[1 + \frac{4w}{w + 1} \frac{m_B^2 - 2wm_b m_{D^*} + m_{D^*}^2}{(m_B - m_{D^*})^2} \right] F^2(w). \end{aligned} \quad (84)$$

All of the HQET analysis goes into the factor $F(w)$, which has an expansion

$$F(w) = \xi(w) + (\text{radiative corrections}) + (\text{power corrections}). \quad (85)$$

We extract $|V_{cb}|$ by studying the differential decay rate near $w = 1$, where the hadronic matrix elements are known. Of course, this requires extrapolation of the experimental data, since the rate vanishes kinematically at $w = 1$. For massless

leptons, only the matrix element $\langle D^* | \bar{c} \gamma^\mu \gamma^5 b | B \rangle$ of the axial current contributes at this point. The analysis of this quantity in the HQET yields an expansion of the form

$$F(1) = \eta_A \left[1 + \frac{0}{m_c} + \frac{0}{m_b} + \delta_{1/m^2} + \dots \right]. \quad (86)$$

The correction δ_{1/m^2} , which contains terms proportional to $1/m_c^2$, $1/m_b^2$, and $1/m_c m_b$, is intrinsically nonperturbative. It has been estimated from a variety of models to be small and negative,^{17,23,24}

$$\delta_{1/m^2} \approx -0.055 \pm 0.035. \quad (87)$$

Note that the model dependence in the result has been relegated to the estimation of the sub-subleading terms. The radiative correction η_A has now been computed to two loops,^{25,26}

$$\eta_A = 0.960 \pm 0.007. \quad (88)$$

The result is a value for $F(1)$ with errors at the level of 5%,

$$F(1) = 0.91 \pm 0.04. \quad (89)$$

This is the theory error which the experimental determination of $|V_{cb}|$ will inherit. It is dominated by the uncertainty in the nonperturbative corrections, and it is difficult to see how this can be improved much in the future.

All that is left experimentally is to extrapolate the data to $w = 1$ and extract

$$\lim_{w \rightarrow 1} \frac{1}{\sqrt{w-1}} \frac{d\Gamma}{dw}. \quad (90)$$

Once the kinematic factors in Eq. (84) have been included, this amounts to a direct measurement of the combination $|V_{cb}|F(1)$. Four experiments recently have reported results for this quantity²⁷⁻³⁰:

$$\begin{aligned} \text{ALEPH :} & \quad (31.4 \pm 2.3 \pm 2.5) \times 10^{-3} \\ \text{ARGUS :} & \quad (38.8 \pm 4.3 \pm 2.5) \times 10^{-3} \\ \text{CLEO :} & \quad (35.1 \pm 1.9 \pm 2.0) \times 10^{-3} \\ \text{DELPHI :} & \quad (35.0 \pm 1.9 \pm 2.3) \times 10^{-3}. \end{aligned} \quad (91)$$

Neubert³¹ has combined these measurements to find a “world average” of $(35.3 \pm 1.8) \times 10^{-3}$, from which we extract

$$|V_{cb}| = (38.8 \pm 2.0_{\text{exp}} \pm 1.6_{\text{th}}) \times 10^{-3}. \quad (92)$$

This value of $|V_{cb}|$ has almost no dependence on hadronic models. In contrast to model-based “measurements,” here the theoretical error is meaningful, in that it is based on a systematic expansion in small quantities.

4 Inclusive B Decays

An exclusive semileptonic B decay, such as $B \rightarrow D \ell \bar{\nu}$, is one in which the final hadronic state is fully reconstructed. An inclusive decay, by contrast, is one in which only certain kinematic features, and perhaps the flavor, of the hadron are known. In this case, we need a theoretical analysis in which we sum over all possible hadronic final states allowed by the kinematics. Fortunately, this is possible within the structure of the HQET.

As in the case of exclusive decays, the key theme is the separation of short-distance physics, associated with the heavy quark, from long-distance physics, associated with the light degrees of freedom. We will also rely on heavy quark spin and flavor symmetry. However, the new ingredient will be the idea of “parton-hadron duality,” which, as we will see, also relies on the heavy quark limit $m_b \gg \Lambda_{\text{QCD}}$.

4.1 The Inclusive Decay $B \rightarrow X_c \ell \bar{\nu}$

Let us consider the inclusive decay

$$B(p_B) \rightarrow X_c(p_X) \ell(p_\ell) \nu(p_\nu), \quad (93)$$

where all that is known about the state X_c is its energy and momentum, and the fact that it contains a charm quark. This decay is mediated by the weak operator \mathcal{O}_{bc} . It is easy to generalize our discussion to inclusive decays of other heavy quarks, such as $b \rightarrow u \ell \nu$ and $c \rightarrow s \bar{\ell} \nu$, by replacing \mathcal{O}_{bc} with the appropriate weak operator.

The treatment of exclusive decays required both the b and c quarks to be heavy. For inclusive decays, we can relax this condition on the c quark, requiring only $m_b \gg \Lambda_{\text{QCD}}$. What does the weak decay of the b , at time $t = 0$, look like to the light degrees of freedom? For $t < 0$, there is a heavy hadron composed of a point-like color source and light quarks and gluons. At $t = 0$, the point source disappears, releasing both its color and a large amount of energy into

the hadronic environment. Eventually, for $t > 0$, this new collection of strongly interacting particles will materialize as a set of physical hadrons. The probability of this hadronization is unity; there is no interference between the hadronization process and the heavy quark decay. There are subleading effects in powers of Λ_{QCD}/m_b , but they do not alter the probability of hadronization. Rather, they reflect the fact that the b quark is not exactly a static source of color: it has a small nonrelativistic kinetic energy and it carries a spin, both of which affect the kinematic properties of its decay.

As in the case of exclusive decays, we will compute the inclusive semileptonic width $\Gamma(B \rightarrow X_c \ell \bar{\nu})$ as a double expansion in $\alpha_s(m_b)$ and Λ_{QCD}/m_b .^{18,32-34} The expansion in $\alpha_s(m_b)$ reflects the applicability of perturbative QCD to the short-distance part of the process. The heavy quark expansion will be continued to relative order $1/m_b^2$, as there is an analogue of Luke's Theorem which eliminates power corrections to the rate of order $1/m_b$. The $1/m_b^2$ corrections will be written in terms of three nonperturbative parameters. The first, $\bar{\Lambda}$, is defined in Eq. (78). It is essentially the mass of the light degrees of freedom in the heavy hadron, but we will see that it is plagued by an ambiguity of order Λ_{QCD} in the definition of the b quark mass. The other two parameters are the expectation values in the B meson of the leading corrections \mathcal{O}_K and \mathcal{O}_G to $\mathcal{L}_{\text{HQET}}$. They are defined as¹⁷

$$\begin{aligned}\lambda_1 &= \frac{1}{2m_B} \langle B | \mathcal{O}_K | B \rangle \\ \lambda_2 &= -\frac{1}{6m_B} \langle B | \mathcal{O}_G | B \rangle,\end{aligned}\quad (94)$$

where we take the usual relativistic normalization of the states. Hence, λ_1 may be thought of roughly as the negative of the b quark kinetic energy, and λ_2 as the energy of its hyperfine interaction with the light degrees of freedom.

Now let us outline the computation. The inclusive decay involves a sum over all possible final states, which is actually a sum over exclusive modes (such as $D, D^*, D\pi, \dots$), followed by a phase space integral for each mode. We write

$$\Gamma(B \rightarrow X_c \ell \bar{\nu}) = \sum_{X_c} \int d[\text{P.S.}] \left| \langle X_c \ell \bar{\nu} | \mathcal{O}_{bc} | B \rangle \right|^2. \quad (95)$$

There is an Optical Theorem for QCD, which follows from the analyticity of the scattering matrix as a function of the momenta of the asymptotic states. Its content is that a transition rate is proportional to the imaginary part of the

forward scattering amplitude with two insertions of the transition operator,

$$\Gamma(B \rightarrow X_c \ell \bar{\nu}) = -2 \text{Im} i \int dx e^{ik \cdot x} \langle B | T \{ \mathcal{O}_{bc}^\dagger(x), \mathcal{O}_{bc}(0) \} | B \rangle \equiv 2 \text{Im} T. \quad (96)$$

In what follows, we will write the time-ordered product $T\{\mathcal{O}_{bc}^\dagger, \mathcal{O}_{bc}\}$ as a series of local operators, using the Operator Product Expansion. As we will see, the applicability of this expansion, and its computation in perturbation theory, will rest on the limit $m_b \gg \Lambda_{\text{QCD}}$. We will then use this limit again to expand the matrix elements of these local operators in the HQET.

The first step is to factorize the integration over the lepton momenta, which can be performed explicitly. Written as a product of currents, \mathcal{O}_{bc} takes the form

$$\mathcal{O}_{bc} = \frac{G_F V_{cb}}{\sqrt{2}} J_{bc}^\mu J_{\ell\mu}, \quad (97)$$

where

$$\begin{aligned}J_{bc}^\mu &= \bar{c} \gamma^\mu (1 - \gamma^5) b \\ J_{\ell}^\mu &= \bar{\ell} \gamma^\mu (1 - \gamma^5) \nu.\end{aligned}\quad (98)$$

Then T can be decomposed as an integral over the total momentum $q^\mu = p_\ell^\mu + p_{\bar{\nu}}^\mu$ transferred to the leptons,

$$T = \frac{1}{2} G_F^2 |V_{cb}|^2 \int dq T^{\mu\nu}(q) L_{\mu\nu}(q). \quad (99)$$

Here the lepton tensor is

$$\begin{aligned}L_{\mu\nu}(q) &= \int d[\text{P.S.}] \langle 0 | J_{\ell\mu}^\dagger | \ell \bar{\nu} \rangle \langle \ell \nu | J_{\ell\nu} | 0 \rangle \\ &= \frac{1}{3\pi} (q_\mu q_\nu - q^2 g_{\mu\nu}),\end{aligned}\quad (100)$$

and the hadron tensor is

$$T^{\mu\nu}(q) = -i \int dx e^{iq \cdot x} \langle B | T \{ J_{bc}^{\mu\dagger}(x), J_{bc}^\nu(0) \} | B \rangle. \quad (101)$$

We will need the imaginary part, $\text{Im} T^{\mu\nu}$. Where is it nonvanishing? In quantum field theory, a scattering amplitude develops an imaginary part when there can be a real intermediate state, that is, the intermediate particles can all go on their mass shell. Whether this is possible, of course, depends on the kinematics of the external states.

In this case, there are two avenues for creating a physical intermediate state.³² The first is to act on the external state $|B\rangle$ with the transition current J_{bc}^μ . The state which is created has no net b number and a single charm quark; the simplest possibility is the decay process $b \rightarrow c$. The momentum of the intermediate state is $p_X = p_B - q$; the condition that it could be on mass shell is simply

$$p_X^2 = (p_B - q)^2 \geq m_D^2. \quad (102)$$

If we define scaled variables

$$p_B^\mu = m_B v^\mu, \quad \hat{q}^\mu = q^\mu / m_B, \quad \hat{m}_D = m_D / m_B, \quad (103)$$

this condition becomes

$$v \cdot \hat{q} \geq \frac{1}{2} (1 + \hat{q}^2 - \hat{m}_D^2). \quad (104)$$

Another possibility is to act on $|B\rangle$ with the conjugate operator $J_{bc}^{\mu\dagger}$. This operation would produce an intermediate state with two b quarks and one \bar{c} . For this state to be on shell, the momentum transfer has to satisfy

$$p_X^2 = (p_B + q)^2 \geq (2m_B + m_D)^2, \quad (105)$$

that is,

$$v \cdot \hat{q} \geq \frac{1}{2} (3 - \hat{q}^2 + 4\hat{m}_D + \hat{m}_D^2). \quad (106)$$

The physical intermediate states are shown as cuts in the $v \cdot \hat{q}$ plane in Fig. 4. Also shown is the contour corresponding to the phase space integration over the lepton momentum q . For physical (massless) leptons which are the product of a heavy quark decay, this integral runs over the top of the lower cut, for the range

$$\sqrt{\hat{q}^2} + i\epsilon \leq v \cdot \hat{q} \leq \frac{1}{2} (1 + \hat{q}^2 - \hat{m}_D^2) + i\epsilon. \quad (107)$$

As indicated by the dotted line, we can continue this contour around the end of the cut and back along the bottom, to $v \cdot \hat{q} = \sqrt{\hat{q}^2} - i\epsilon$. Since $T^{\mu\nu}(v \cdot \hat{q}^*) = -T^{\mu\nu}(v \cdot \hat{q})$ for real \hat{q}^2 , we compensate for extending the contour by dividing the new integral by two.

We now encounter our central problem. The integral over $v \cdot \hat{q}$ runs over physical intermediate hadron states, which are color-neutral bound states of quarks and gluons. Hence, the integrand depends intimately on the details of QCD at long distances, which is intrinsically nonperturbative. A perturbative calculation of $T^{\mu\nu}$, which is all we have at our disposal, would appear to be of no use.




Figure 4: Analytic structure of $T^{\mu\nu}$ in the complex $v \cdot \hat{q}$ plane, for fixed, real \hat{q}^2 . The integration contour is over the “physical cut,” corresponding to real decay into leptons. The unphysical cuts correspond to other processes.

The solution is to deform the contour away from the cut, into the complex $v \cdot \hat{q}$ plane, as shown in Fig. 5. Since the scale of momenta is set by m_b , the contour is now a distance $\sim m_b$ away from the resonances.³² Since $m_b \gg \Lambda_{\text{QCD}}$, it is reasonable to hope that a perturbative treatment in this region is valid. Essentially, we are saved because we do not need to know $T^{\mu\nu}(q)$ for every value of q , just suitable integrals of $T^{\mu\nu}$. That we can use such arguments to compute perturbatively the *average* value of a hadronic quantity, where at each point the quantity depends on nonperturbative physics, is known as (global) *parton-hadron duality*.

Parton-hadron duality has the status of being somewhat more than an assumption, since it is known to hold in QCD in the limit $m_b \rightarrow \infty$, but somewhat less than an approximation, since it is not known how to compute systematically the leading corrections to it. In any case, the limit $m_b \gg \Lambda_{\text{QCD}}$ plays a crucial role here. By deforming the integration contour a distance $\sim m_b$ away from the resonance regime, we find the correspondence in QCD of our earlier intuitive statement: the probability of the decay products materializing as physical hadrons is unity, independent of the kinematics of the short-distance process. The local redistribution of probability in phase space due to the presence of hadronic resonances is irrelevant to the total decay. Finally, we should note that since we do not have control over the corrections to local duality, it might work better in some processes than in others, for reasons that need not be apparent from within the calculation.




Figure 5: The deformation of the integration contour into the complex $v \cdot \hat{q}$ plane.

Hence, one must be particularly wary of drawing dramatic conclusions from any surprising results of these inclusive calculations.³⁵

Let us perform the operator product expansion at tree level and for decay kinematics. The Feynman diagram is given in Fig. 6, which yields the expression

$$T^{\mu\nu} = \bar{b}\gamma^\mu(1 - \gamma^5) \frac{\not{p}_b - \not{q} + m_c}{(p_b - q)^2 - m_c^2 + i\epsilon} \gamma^\nu(1 - \gamma^5)b. \quad (108)$$

We now write

$$\begin{aligned} p_b^\mu &= m_b v^\mu + k^\mu = m_b(v^\mu + \hat{k}^\mu) \\ \hat{q}^\mu &= q^\mu/m_b \\ \hat{m}_c &= m_c/m_b \\ b(x) &= e^{-im_b v \cdot x} h_v(x) + O(1/m_b), \end{aligned} \quad (109)$$

and expand in powers of $1/m_b$. Since the operator product expansion is in terms of the effective field h_v , a factor of k^μ corresponds to an insertion of the covariant derivative iD^μ . Operator ordering ambiguities are to be resolved by considering graphs with external gluon fields.




Figure 6: The operator product expansion at tree level.

As an example of the procedure, let us expand the propagator to order $1/m_b^2$. (There are also corrections to the currents at this order, which are included in a full calculation.) It is convenient to define the scaled hadronic invariant mass,

$$\hat{s} = (p_b - q)^2/m_b^2 = 1 - 2v \cdot \hat{q} + \hat{q}^2. \quad (110)$$

Then we find a contribution to $T^{\mu\nu}$ of the form

$$T^{\mu\nu} = \frac{1}{m_b} \bar{h}_v \gamma^\mu (\not{p} - \not{q}) \gamma^\nu (1 - \gamma^5) \left[\frac{1}{\hat{s} - \hat{m}_c^2 + i\epsilon} + \frac{2\hat{k} \cdot \hat{q} - \hat{q}^2}{(\hat{s} - \hat{m}_c^2 + i\epsilon)^2} + \dots \right] h_v + \dots \quad (111)$$

From this expression, we can read off the operators which appear in the operator product expansion. Since

$$\text{Im} \frac{1}{(\hat{s} - \hat{m}_c^2 + i\epsilon)^n} \propto \delta^{(n-1)}(\hat{s} - \hat{m}_c^2), \quad (112)$$

we see that the effect of taking the imaginary part in each term is to put the charm quark on its mass shell. The leading term is a quark bilinear,

$$\frac{1}{m_b} \bar{h}_v \gamma^\mu (\not{p} - \not{q}) \gamma^\nu (1 - \gamma^5) h_v. \quad (113)$$

It is straightforward to compute its matrix element in the HQET using the trace formalism,

$$\begin{aligned} \langle B | \bar{h}_v \gamma^\mu (\not{p} - \not{q}) \gamma^\nu (1 - \gamma^5) h_v | B \rangle \\ = 2m_B (2v^\mu v^\nu - g^{\mu\nu} - v^\mu \hat{q}^\nu - v^\nu \hat{q}^\mu + g^{\mu\nu} v \cdot \hat{q} + \dots). \end{aligned} \quad (114)$$

The ellipses denote terms of order $1/m_b^2$; there are no corrections of order $1/m_b$, by Luke's Theorem. Finally, we contract the tensor $T^{\mu\nu}$ with $L_{\mu\nu}$ and perform the phase space integration [Eq. (99)]. In the end, the result is the same as we would have gotten directly by computing free quark decay.

Of course, if we only intended to reproduce the free quark decay result, we would never have introduced so much new formalism. The value of the HQET framework is that it allows us to go beyond leading order and compute the next terms in the series in $1/m_b^2$. For example, consider the operators induced by the expansion of the propagator (111). The correction of order $1/m_b$ comes from the operator

$$\frac{1}{m_b^2} \frac{2}{(\hat{s} - \hat{m}_c^2 + i\epsilon)^2} \bar{h}_v \gamma^\mu (\not{p} - \not{q}) \gamma^\nu (1 - \gamma^5) \hat{q} \cdot iD h_v. \quad (115)$$

However, the matrix element of this operator is of the form

$$\langle B | \bar{h}_v \Gamma^\alpha(v, q) i D_\alpha h_v | B \rangle, \quad (116)$$

which, as we have seen, vanishes by the classical equation of motion. In fact, since all $1/m_b$ corrections, from any source, have a single covariant derivative, they all vanish in the same way. This is the analogue of Luke's Theorem for inclusive decays.³² The correction of order $1/m_b^2$ in Eq. (111) is

$$-\frac{1}{m_b^3} \frac{1}{(\hat{s} - \hat{m}_c^2 + i\epsilon)^2} \bar{h}_v \gamma^\mu (\not{p} - \not{q}) \gamma^\nu (1 - \gamma^5) (iD)^2 h_v. \quad (117)$$

The matrix element of this operator is related by the heavy quark symmetry to λ_1 , the expectation value of \mathcal{O}_K . The full expansion of $T^{\mu\nu}$ also induces operators with explicit factors of the gluon field, whose matrix elements are related to λ_2 .

We now present the result for the inclusive semileptonic decay rate, up to order $1/m_b^2$ in the heavy quark expansion, and with the complete radiative correction of order α_s . We also include that part of the two-loop correction which is proportional to $\beta_0 \alpha_s^2$. Since $\beta_0 \approx 9$, perhaps this term dominates the two-loop result. In any case, it is interesting for other reasons, as we will see below.

Let us first consider the decay $B \rightarrow X_u \ell \bar{\nu}$, for which the decay rate simplifies since $m_u = 0$. We find^{33,34,36,37,40}

$$\begin{aligned} \Gamma(B \rightarrow X_u \ell \bar{\nu}) = \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_b^5 \left[1 + \left(\frac{25}{6} - \frac{2\pi^2}{3} \right) \frac{\alpha_s(m_b)}{\pi} \right. \\ - (2.98\beta_0 + C_u) \left(\frac{\alpha_s(m_b)}{\pi} \right)^2 + \dots \\ \left. + \frac{\lambda_1 - 9\lambda_2}{2m_b^2} + \dots \right]. \quad (118) \end{aligned}$$

When we include the charm mass, it is convenient to write the unknown quark masses in terms of the measured meson masses and the parameters of the HQET. In terms of the spin averaged mass $\bar{m}_B = (m_B + 3m_{B^*})/4$, we have

$$m_b = \bar{m}_B - \bar{\Lambda} + \frac{\lambda_1}{2m_B} + \dots, \quad (119)$$

and analogously for m_c . We then find^{33,34,38,39}

$$\begin{aligned} \Gamma(B \rightarrow X_c \ell \bar{\nu}) = \frac{G_F^2 |V_{cb}|^2}{192\pi^3} m_B^5 \times 0.369 \left[1 - 1.54 \frac{\alpha_s(m_b)}{\pi} \right. \\ - (1.43\beta_0 + C_c) \left(\frac{\alpha_s(m_b)}{\pi} \right)^2 + \dots \\ - 1.65 \frac{\bar{\Lambda}}{m_B} \left(1 - 0.87 \frac{\alpha_s(m_b)}{\pi} \right) \\ \left. - 0.95 \frac{\bar{\lambda}^2}{m_B^2} - 3.18 \frac{\lambda_1}{m_B^2} + 0.02 \frac{\lambda_2}{m_B^2} + \dots \right]. \quad (120) \end{aligned}$$

All the coefficients which appear in this expression are known functions of \bar{m}_D/\bar{m}_B and are evaluated at the physical point $\bar{m}_D/\bar{m}_B = 0.372$. In both $B \rightarrow X_u \ell \bar{\nu}$ and $B \rightarrow X_c \ell \bar{\nu}$, the power corrections proportional to λ_1 and λ_2 are numerically small, at the level of a few percent.

4.2 Renormalons and the Pole Mass

The inclusive decay rate depends on the heavy quark mass m_b , either explicitly, as in Eq. (118), or implicitly through $\bar{\Lambda}$, as in Eq. (120). At tree level, m_b is just the coefficient of the $\bar{b}b$ term in the QCD Lagrangian, but beyond that, we are faced with the question of what exactly we mean by m_b . Should we take an $\overline{\text{MS}}$ mass, such as $\bar{m}_b(m_b)$? Or should we take the pole mass m_b^{pole} , or maybe some other quantity? The various prescriptions for m_b can vary by hundreds of MeV, and since the total rate is proportional to m_b^5 , the question is of practical importance if we hope to make accurate phenomenological predictions.

At a fixed order in QCD perturbation theory, the answer is clear. The heavy quark masses which appear come from poles in quark propagators, so we should take m_b^{pole} (and m_c^{pole}). This is also the prescription for the mass which cancels out the on-shell part of the heavy quark field in the construction of $\mathcal{L}_{\text{HQET}}$. Hence, the difference of heavy quark pole masses is known quite well,

$$\begin{aligned} m_b^{\text{pole}} - m_c^{\text{pole}} &= \left(\bar{m}_B - \bar{\Lambda} + \frac{\lambda_1}{2m_B} + \dots \right) - \left(\bar{m}_D - \bar{\Lambda} + \frac{\lambda_1}{2m_D} + \dots \right) \\ &= 3.34 \text{ GeV} + O(\Lambda_{\text{QCD}}^2/m_Q^2). \quad (121) \end{aligned}$$

Since $\Gamma(B \rightarrow X_c \ell \bar{\nu})$ depends approximately as $m_b^2(m_b - m_c)^3$, the uncertainty due to quark mass dependence is reduced.

The problem, of course, is that there is no sensible nonperturbative definition of m_b^{pole} , since due to confinement, there is no actual pole in the quark propagator. Hence, a direct experimental determination of a value for m_b^{pole} to insert into the theoretical expressions Eq. (118) and Eq. (120) is not possible. How, then, can we do phenomenology?

One approach would be to define m_b^{pole} to be the pole mass as computed in perturbation theory, truncate at some order, and then estimate the theoretical error from the uncomputed higher order terms. However, it turns out that even *within* perturbation theory, the concept of a quark pole mass is ambiguous. Consider a particular class of diagrams which contribute to m_b^{pole} , shown in Fig. 7. The perturbation theory is developed as an expansion in the small parameter $\alpha_s(m_b)$, so we hope that it will be well-behaved. Each of the bubbles represents an insertion of the gluon self-energy, which is proportional at lowest order to $\alpha_s(m_b)\beta_0$. Of course, the infinite sum of the graphs in Fig. 7 can be absorbed into the one-loop graph, with a compensating change in the coupling from $\alpha_s(m_b)$ to $\alpha_s(q)$, where q is the loop momentum. The result is an expansion for m_b^{pole} of the form

$$m_b^{\text{pole}} = \bar{m}_b(m_b) \left[1 + a_1 \alpha_s(m_b) + (a_2 \beta_0 + b_2) \alpha_s^2(m_b) + (a_3 \beta_0^2 + b_3 \beta_0 + c_3) \alpha_s^3(m_b) + \dots \right]. \quad (122)$$

The graphs in Fig. 7 contribute the terms proportional to $\alpha_s^{n+1}(m_b)\beta_0^n$. Since $\beta_0 \approx 9$, these terms are “intrinsically” larger than ones with fewer powers of β_0 , and we might hope that their sum approximates the full series. However, it is important to realize that the only limit of QCD in which such terms actually dominate is that of a large number of light quark flavors, in which case the sign of β_0 is opposite to that of QCD. Although this is a physical limit of an Abelian theory, we are certainly not close to that limit here. The ansatz of keeping only the terms proportional to $\alpha_s^{n+1}(m_b)\beta_0^n$ is known as “naive non-Abelianization” (NNA).⁴¹

What is most interesting about the series of terms shown in Fig. 7,

$$\sum a_n \alpha_s^n(m_b) \beta_0^{n-1}, \quad (123)$$

is that it does not converge. Already in the graphs kept in the NNA ansatz, we are sensitive to the fact that QCD is an asymptotic, rather than a convergent expansion. For large n , the coefficients a_n diverge as $n!$, much stronger than any




Figure 7: The radiative corrections to m_b^{pole} of order $\alpha_s(m_b)^{n+1}\beta_0^n$.

convergence due to the powers $\alpha_s^n(m_b)$. The series can only be made meaningful if this divergence is subtracted. As with many subtraction prescriptions, there is a residual finite ambiguity.[§] This ambiguity, known as an “infrared renormalon,” leads to an ambiguity in the pole mass of order^{41–43}

$$\delta m_b^{\text{pole}} \sim 100 \text{ MeV}. \quad (124)$$

By the definition [Eq. (78)], $\bar{\Lambda}$ also inherits this ambiguity.

The expressions Eq. (118) and Eq. (120) are plagued by two problems. The first is the renormalon ambiguity in m_b^{pole} and $\bar{\Lambda}$. The second is that the perturbative expansion for the rate Γ is itself divergent, and *also* has an infrared renormalon. In the expansion

$$\Gamma = \Gamma_0 \left[\sum a'_n \alpha_s^n(m_b) \beta_0^{n-1} + (\text{power corrections}) \right], \quad (125)$$

the coefficients a'_n also diverge as $n!$ However, it turns out that these two problems actually cure each other, because the infrared renormalons in m_b^{pole} and in the perturbation series for Γ cancel.^{42,44} We can exploit this cancellation to improve the predictive power of the theoretical computation of the rate. Without this improvement, the infrared renormalons render the expressions Eq. (118) and Eq. (120) of dubious phenomenological utility.

The most reliable approach, theoretically, is to eliminate m_b^{pole} or $\bar{\Lambda}$ explicitly from the rate by computing and measuring another quantity which also depends on it. For example, let us consider the charmless decay rate $\Gamma(B \rightarrow X_u \ell \bar{\nu})$ and

[§]In a formal treatment, this ambiguity arises from a choice of contour in the Borel plane.

the average invariant mass $\langle s_H \rangle$ of the hadrons produced in the decay. Each of these expressions suffers from a poorly behaved perturbation series in the NNA approximation. Ignoring terms of relative order $1/m_b^2$ and writing the rate in terms of $\bar{\Lambda}$ instead of m_b^{pole} , we find to five-loop order,⁴¹

$$\begin{aligned}\Gamma &= \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_B^5 \left[1 - 2.41 \frac{\alpha_s}{\pi} - 2.98 \left(\frac{\alpha_s}{\pi} \right)^2 \beta_0 - 4.43 \left(\frac{\alpha_s}{\pi} \right)^3 \beta_0^2 \right. \\ &\quad \left. - 7.67 \left(\frac{\alpha_s}{\pi} \right)^4 \beta_0^3 - 15.7 \left(\frac{\alpha_s}{\pi} \right)^5 \beta_0^4 + \dots - 5 \frac{\bar{\Lambda}}{m_B} + \dots \right] \\ &= \frac{G_F^2 |V_{ub}|^2}{192\pi^3} m_B^5 \left[1 - 0.061 - 0.120 - 0.107 - 0.111 - 0.136 + \dots \right. \\ &\quad \left. - 5\bar{\Lambda}/m_B + \dots \right],\end{aligned}\quad (126)$$

for $\alpha_s(m_b) = 0.21$ and $\beta_0 = 9$. As we see, not only does the perturbation series fail to converge, it does not even have an apparent smallest term, where one should truncate to minimize the error of the asymptotic series. The series for $\langle s_H \rangle$ exhibits a similar behavior,⁴⁰

$$\begin{aligned}\langle s_H \rangle &= m_B^2 \left[0.20 \frac{\alpha_s}{\pi} + 0.35 \left(\frac{\alpha_s}{\pi} \right)^2 \beta_0 + 0.64 \left(\frac{\alpha_s}{\pi} \right)^3 \beta_0^2 + 1.29 \left(\frac{\alpha_s}{\pi} \right)^4 \beta_0^3 \right. \\ &\quad \left. + 2.95 \left(\frac{\alpha_s}{\pi} \right)^5 \beta_0^4 + \dots + \frac{7}{10} \frac{\bar{\Lambda}}{m_B} + \dots \right] \\ &= m_B^2 \left[0.0135 + 0.0141 + 0.0156 + 0.0189 + 0.0261 + \dots \right. \\ &\quad \left. - 7\bar{\Lambda}/10m_B + \dots \right].\end{aligned}\quad (127)$$

However, the situation improves dramatically if we eliminate $\bar{\Lambda}$ and write Γ directly in terms of $\langle s_H \rangle$,

$$\Gamma = \frac{G_F^2 |V_{ub}|^2 m_B^5}{192\pi^3} \left[1 - 7.14 \frac{\langle s_H \rangle}{m_B^2} - 0.064 - 0.020 - 0.0002 - 0.022 - 0.047 + \dots \right].\quad (128)$$

By truncating this series at its smallest term, 0.0002, we obtain a new expression in which the theoretical errors are under control. The price is that we must now measure a second quantity, $\langle s_H \rangle$, in the same decay.

An analogous approach works for inclusive decays to charm, in which case the new quantity which we must measure is $\langle s_H - \bar{m}_D^2 \rangle$. Let us apply this analysis to the extraction of V_{cb} from the inclusive decay $B \rightarrow X_c \ell \bar{\nu}$. Eliminating $\bar{\Lambda}$ as before, we find⁴⁰

$$\Gamma(B \rightarrow X_c \ell \nu) = \frac{G_F^2 |V_{cb}|^2}{192\pi^3} m_B^5 \times 0.369 \left[1 - 1.17 \frac{\alpha_s(m_b)}{\pi} - 0.74 \beta_0 \frac{\alpha_s^2(m_b)}{\pi^2} \right. \\ \left. - 7.17 \frac{\langle s_H - \bar{m}_D^2 \rangle}{m_B^2} + \dots \right],\quad (129)$$

plus small terms of order $\lambda_{1,2}/m_B^2$. The current data on $\langle s_H - \bar{m}_D^2 \rangle$ are still inconclusive, but they will improve in the future. As an illustration of the utility of the method, however, let us consider the physical limit $\langle s_H - \bar{m}_D^2 \rangle \geq 0$. In that case, we can already obtain the bound

$$V_{cb} \geq 0.037 \times \left(\frac{\tau_B}{1.54 \text{ ps}} \right)^{-1/2},\quad (130)$$

where we have used the observed B semileptonic branching ratio of 10.7%. As the data on $\langle s_H - \bar{m}_D^2 \rangle$ improve, so will this bound. Note that it is entirely consistent with the value [Eq. (92)] obtained from exclusive decays.

An alternative approach is to express the width Γ in terms of the running mass $\bar{m}_b(m_b)$ instead of another inclusive observable.^{42,45} Since the $\overline{\text{MS}}$ mass is a short-distance quantity, this also eliminates the infrared renormalon, which is associated with long-distance physics. However, from a phenomenological point of view, it begs the question of how the running mass is to be determined from experiment. Possibilities include quarkonium spectroscopy, QCD sum rules, and lattice calculations, but in all of these cases, it is unclear both how to determine reliably the accuracy of the method, and how to deal with renormalon ambiguities in a manner that is *consistent* with their treatment in the calculation of Γ . With additional progress, however, such an approach might eventually prove fruitful.

5 Heavy Hadron Production in Fragmentation

So far, we have applied the idea of heavy quark symmetry to heavy hadron spectroscopy and to heavy hadron decay. We can also apply it to heavy hadron production via fragmentation. Although there is not yet available the wealth of detailed data for production processes that there is for spectroscopy and decays,

the study of heavy quark fragmentation is an elegant application of the heavy quark limit in a new regime.¹⁶

5.1 The Physics of Heavy Quark Fragmentation

The production of a heavy hadron proceeds in two steps. First, the heavy quark itself must be created; because of its large mass, this process takes place over a time scale τ_Q which is very short, $\tau_Q \leq 1/m_Q$. Second, some light degrees of freedom assemble themselves about the heavy quark to make a color-neutral heavy hadron, a process which involves nonperturbative strong interactions and typically takes much longer, $\tau_F \sim 1/\Lambda_{\text{QCD}} \gg \tau_Q$. If the heavy quark is produced with a large velocity in the center-of-mass frame, and if there is plenty of available energy, then production of these light degrees of freedom will be local in phase space and independent of the initial state. This is the fragmentation regime. We will see that heavy quark symmetry simplifies the description of heavy hadron production via fragmentation, because, as before, it allows us to separate certain properties of the heavy quark from those of the light degrees of freedom. This is particularly important in the production of excited heavy hadrons, for which the behavior of the spin of the light degrees of freedom can be quite interesting.

Heavy quark symmetry simplifies the picture of fragmentation in several ways. First, since the fragmentation process involves the exchange only of soft momenta, the velocity v^μ of the hadron H_Q which is produced is the same as the velocity of the heavy quark Q . Second, the angular momenta S_Q and J_ℓ of the heavy quark and the light degrees of freedom are well-defined and can be followed individually through the processes of production and decay. Third, the typical precession time τ_S of S_Q and J_ℓ about each other is of order $\tau_S \sim m_Q/\Lambda_{\text{QCD}}^2 \gg 1/\Lambda_{\text{QCD}}$, since the precession is induced by the chromomagnetic interaction. Hence, $\tau_S \gg \tau_F$, and during the fragmentation process, S_Q can essentially be taken to be frozen in direction.

Suppose that Q fragments initially to a state with light degrees of freedom with spin J_ℓ . This could correspond to two possible hadronic states: H_Q , with spin $J = J_\ell - \frac{1}{2}$, or H_Q^* , with spin $J = J_\ell + \frac{1}{2}$. Let the states $H_Q^{(*)}$ have lifetime τ . How should we treat the interaction of S_Q with J_ℓ ?

The answer depends critically on the relationship of τ to the precession time τ_S . While the order of magnitude of τ_S is essentially fixed at $m_Q/\Lambda_{\text{QCD}}^2$, the lifetime

τ can vary widely, depending on the channel in which $H_Q^{(*)}$ primarily decays. Of course, if $H_Q^{(*)}$ decays weakly, then $\tau \gg \tau_S$. But if $H_Q^{(*)}$ decays strongly, then τ depends critically on the phase space available for the transition. In the absence of phase space factors, $\tau \sim 1/\Lambda_{\text{QCD}} \ll \tau_S$. But since the pion, the lightest hadron, is not massless, it is possible to have a situation where an allowed strong decay is either forbidden by phase space (as with $B^* \rightarrow B + \pi$) or severely suppressed (as with $D^* \rightarrow D + \pi$).

The treatment of a given hadron doublet depends on the relative size of τ and τ_S . Let us consider the two extreme possibilities. The first corresponds to a strong decay with plenty of phase space, so $\tau_S \gg \tau$. Here, H_Q and H_Q^* are formed and then decay before the angular momenta S_Q and J_ℓ have a chance to interact. The fact that there is no precession means that there is no change in the orientation of S_Q and J_ℓ . If, for example, either spin was polarized before or during the fragmentation process, this polarization would be undisturbed by passing through the resonance $H_Q^{(*)}$. The strong decay of the heavy hadron then would reflect the orientation of the light degrees as they were produced.

Note that it is the very same spin exchange interaction which is inhibited here which is responsible for the splitting Δ_H between H_Q and H_Q^* . Hence, under these conditions, the resonances are almost completely *overlapping*, with widths $\Gamma = 1/\tau$ satisfying $\Gamma \gg \Delta_H$. This is another consequence of the effective decoupling of S_Q and J_ℓ , which are independent good quantum numbers of the resonances. It also provides a convenient criterion for determining directly whether we are in the regime $\tau_S \gg \tau$: we cannot distinguish separate resonances for H_Q and H_Q^* .

The second possibility is the opposite extreme, $\tau_S \ll \tau$. This corresponds to heavy hadrons which decay weakly or electromagnetically, or to strong decays which are severely suppressed by phase space. Here the spins S_Q and J_ℓ have plenty of time to interact, precessing about each other many times before H_Q and H_Q^* decay. There is at least a partial degradation of any initial polarization of S_Q , as well as a degradation of any information about the fragmentation process which may be carried by the light degrees of freedom. The signature of this situation is that the states H_Q and H_Q^* are well-separated resonances, since the chromomagnetic interactions have ample opportunity to produce a hyperfine splitting much larger than the width, $\Delta_H \gg \Gamma$. In contrast with the first case, here the heavy and light spins are resolved into states of definite total spin $J = |J_\ell \pm \frac{1}{2}|$. This resolution destroys information about the individual angular momenta S_Q and J_ℓ ,

and the decays of H_Q and H_Q^* only partially reflect the features of the production process.

5.2 Production and Decay of D_1 and D_2^*

We will illustrate these ideas with two examples. The first concerns the production and decay of the excited charmed mesons D_1 and D_2^* . These are states with light degrees of freedom with angular momentum and parity $J_\ell^P = \frac{3}{2}^+$; in the quark model, this would correspond to a light antiquark with a unit of orbital angular momentum. The physical states are the D_1 , with mass 2423 MeV and width 18 MeV, and the D_2^* , with mass 2458 MeV and width 21 MeV. The splitting between them is 35 MeV; although this is not much larger than the individual widths, for simplicity, we will treat them in this limiting case $\tau \gg \tau_S$.

Let us review the sequence of events. First, the charm quark is created in some hard interaction, with a time scale $\tau_c \sim 1/m_c$. Second, light degrees of freedom with $J_\ell = \frac{3}{2}$ are created in a fragmentation process, over a time $\tau_F \sim 1/\Lambda_{\text{QCD}}$, forming a color-neutral charmed hadron. Third, the angular momenta S_c and J_ℓ precess about each other, resolving the individual D_1 and D_2^* states, which requires a time $\tau_S \sim m_c/\Lambda_{\text{QCD}}^2$. Then, finally, after a time τ greater than $m_c/\Lambda_{\text{QCD}}^2$, the D_1 or D_2^* emits a pion and decays to a D or D^* . One question we might ask about this process is, what is the alignment of the light degrees of freedom produced in the fragmentation? Of course, the answer depends on nonperturbative QCD and is difficult to calculate from first principles. Still, we can explore the question experimentally.

Since the total angular momentum of the light degrees of freedom is $J_\ell = \frac{3}{2}$, they can be produced with helicity $h = \pm\frac{3}{2}$ or $h = \pm\frac{1}{2}$ along the fragmentation axis. While parity invariance of the strong interactions requires that the probabilities for helicities h and $-h$ are identical, the relative production of light degrees of freedom with $|h| = \frac{3}{2}$ versus $|h| = \frac{1}{2}$ is determined in some complicated way by nonperturbative dynamics. Let the quantity $w_{3/2}$ denote the probability that $|h| = \frac{3}{2}$,

$$w_{3/2} = P(h = \frac{3}{2}) + P(h = -\frac{3}{2}). \quad (131)$$

Then $1 - w_{3/2}$ is the probability that $|h| = \frac{1}{2}$. Completely isotropic production corresponds to $w_{3/2} = \frac{1}{2}$. The observable $w_{3/2}$ is a new nonperturbative parameter of QCD, which is well-defined only in the heavy quark limit.

This new parameter can be measured directly in the strong decay of the D_2^* or D_1 . For example, consider the angular distribution of the pion with respect to the fragmentation axis in the decay $D_2^* \rightarrow D + \pi$. This is a decay of the light degrees of freedom in the excited hadron, so it will depend on their initial orientation (that is, on $w_{3/2}$) and on the details of the precession of J_ℓ around S_Q during the lifetime of the D_2^* . Following the direction of J_ℓ through the sequence of fragmentation, precession, and decay, we find the distribution⁴⁶

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{1}{4} \left[1 + 3\cos^2\theta - 6w_{3/2}(\cos^2\theta - \frac{1}{3}) \right]. \quad (132)$$

This distribution is isotropic only when $w_{3/2} = \frac{1}{2}$, that is, when the light degrees of freedom are produced isotropically in the fragmentation process. Similar distributions are found in the decays $D_2^* \rightarrow D^* + \pi$ and $D_1 \rightarrow D^* + \pi$.

A fit of ARGUS data⁴⁷ to the expression Eq. (132) seems to indicate that a small value of $w_{3/2}$ is preferred; while the errors are large, we find that $w_{3/2} < 0.24$ at the 90% confidence level. It would be nice to confirm this result with a sharper measurement, and not only for the charmed mesons but in the bottom system as well. Since $w_{3/2}$ is intrinsically nonperturbative, we do not have any real theoretical understanding of why it should be small, although one can construct a qualitative understanding based on a string picture of fragmentation.⁴⁸

5.3 Polarization of Λ_b at SLC/LEP

The second example springs from a practical question: What is the polarization of Λ_b baryons produced at the Z pole? This question is motivated by the fact that b quarks produced in the decay of the Z are 94% polarized left-handed. Since the Λ_b is composed of a b quark and light degrees of freedom with $J_\ell = 0$, the orientation of a Λ_b is identical to the orientation of the b quark inside it. Similarly, the b quark spin does not precess inside a Λ_b . Hence, if a b quark produced at the Z fragments to a Λ_b , then those baryons should inherit the left-handed polarization of the quarks and reveal it in their weak decay.

Unfortunately, life is not that simple. Two recent measurements of Λ_b polarization $P(\Lambda_b)$ from LEP are^{49,50}

$$\begin{aligned} \text{DELPHI :} & \quad 0.08 \begin{matrix} +0.35(\text{stat.}) \\ -0.29(\text{stat.}) \end{matrix} \begin{matrix} +0.18(\text{sys.}) \\ -0.16(\text{sys.}) \end{matrix} \\ \text{ALEPH :} & \quad 0.26 \begin{matrix} +0.20(\text{stat.}) \\ -0.25(\text{stat.}) \end{matrix} \begin{matrix} +0.12(\text{sys.}) \\ -0.13(\text{sys.}) \end{matrix}, \end{aligned}$$

both a long way from $P(\Lambda_b) = 0.94$. The reason is that not all b quarks which wind up as Λ_b baryons get there directly. There is a competing process, in which they fragment first to the excited baryons Σ_b and Σ_b^* , which then decay to Λ_b via pion emission. These new states have light degrees of freedom with $J_\ell = 1$. If they have a lifetime $\tau > \tau_S$, then the b quark will have time to precess about J_ℓ and its polarization will be degraded. The result will be a net sample of Λ_b 's with a polarization less than 94%, as is in fact observed.

In addition to the requirement that $\tau > \tau_S$ for the $\Sigma_b^{(*)}$, any depolarization of Λ_b 's by this mechanism depends on two unknown quantities. First, there is the production rate f of $\Sigma_b^{(*)}$ relative to Λ_b . Isospin and spin counting enhance f by a factor of nine, while the mass splitting between $\Sigma_b^{(*)}$ and Λ_b suppresses it; studies based on the Lund Monte Carlo⁵¹ indicate $f \approx 0.5$ with a very large uncertainty. Of course, it would be preferable to measure f directly. Second, there is the orientation of the spin J_ℓ with respect to the fragmentation axis. This orientation, which is nonperturbative in origin, allows the helicities $h = 1, 0, -1$. By analogy with the treatment of the D_1 and D_2^* , let us define

$$w_1 = P(h = 1) + P(h = -1). \quad (133)$$

In this case, isotropic production corresponds to $w_1 = \frac{2}{3}$. We may measure w_1 from the angle of the pion with respect to the fragmentation axis in the decay $\Sigma_b^* \rightarrow \Lambda_b + \pi$ (Ref. 46),

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\cos\theta} = \frac{1}{4} \left[1 + 3\cos^2\theta - \frac{2}{3}w_1(\cos^2\theta - \frac{1}{3}) \right]. \quad (134)$$

It turns out that the decay $\Sigma_b \rightarrow \Lambda_b + \pi$ is isotropic in $\cos\theta$ for any value of w_1 .

The polarization retention of the Λ_b may be computed in terms of f and w_1 . As before, we will consider the simpler situation in which the Σ_b and the Σ_b^* do not overlap, so $\tau \gg \tau_S$. Then the polarization of the observed Λ_b 's is $P(\Lambda_b) = R(f, w_1)P(b)$, where $P(b) = 94\%$ is the initial polarization of the b quarks, and⁴⁶

$$R(f, w_1) = \frac{1 + \frac{1}{9}(1 + 4w_1)f}{1 + f}. \quad (135)$$

Note that for $f = 0$ (no $\Sigma_b^{(*)}$'s are produced), $R(0, w_1) = 1$ and there is no depolarization. For the Lund value $f = 0.5$, R ranges between 0.70 and 0.85 for $0 \leq w_1 \leq 1$.

Can the very low measured values of $P(\Lambda_b)$ be accommodated by the present data on the $\Sigma_b^{(*)}$? The situation is still unclear, because not much is known about the Σ_b and Σ_b^* states. DELPHI⁵² has reported the observation of these states, but with masses which deviate significantly from those that one would expect based on heavy quark symmetry and the observed Σ_c and Σ_c^* (Ref. 54). Along with the masses, DELPHI also reports $w_1 \approx 0$ and $1 < f < 2$. If this is confirmed, then a polarization in the range $P(\Lambda_b) \approx 40\% - 50\%$ is easy to accommodate. On the other hand, the CLEO analysis⁵³ of the decay of the Σ_c^* yields $w_1 = 0.71 \pm 0.13$, consistent with isotropic fragmentation. Recall that by heavy quark symmetry, w_1 measured in the charm and bottom systems must be the same, so this result is inconsistent with the report from DELPHI. Clearly, further measurements are needed to resolve this situation.

6 Concluding Remarks

Unfortunately, we have had time in these lectures only to introduce a very few of the many applications of heavy quark symmetry and the HQET to the physics of heavy hadrons. Since its development less than ten years ago, it has become one of the basic tools of QCD phenomenology. Much of the popularity and utility of the HQET certainly come from its essential simplicity. The elementary observation that the physics of heavy hadrons can be divided into interactions characterized by short and long distances gives us immediately a clear and compelling intuition for the properties of heavy-light systems. The straightforward manipulations which lead to the HQET then allow this intuition to form the basis for a new systematic expansion of QCD. The deeper understanding of heavy hadrons which we thereby obtain will become increasingly important as the end of the second millennium approaches and the B Factory Era begins.

Acknowledgments

I would like to thank the organizers of the Summer Institute for the opportunity to present these lectures, and for arranging a most interesting and pleasant summer school. This work was supported by the National Science Foundation under Grant No. PHY-9404057 and National Young Investigator Award No. PHY-9457916, by

the Department of Energy under Outstanding Junior Investigator Award No. DE-FG02-94ER40869, and by the Alfred P. Sloan Foundation.

References

- [1] H. Georgi in *Proceedings of TASI-91*, edited by R. K. Ellis *et al.* (World Scientific, Singapore, 1991); B. Grinstein in *High Energy Phenomenology*, edited by R. Huerta and M. A. Peres (World Scientific, Singapore, 1991); N. Isgur and M. B. Wise in *Heavy Flavors*, edited by A. Buras and M. Lindner (World Scientific, Singapore, 1992); T. Mannel in *QCD - 20 Years Later*, edited by P. Zerwas and H. Kastrup (World Scientific, Singapore, 1993); M. Neubert, Phys. Rep. **245**, 259 (1994); M. Shifman, preprint TPI-MINN-95-31-T, to appear in *QCD and Beyond, Proceedings of TASI-95*.
- [2] For example, see M. Wirbel, B. Stech, and M. Bauer, Z. Phys. C **29**, 637 (1985); J. G. Körner and G. A. Schuler, Z. Phys. C **38**, 511 (1988); N. Isgur, D. Scora, B. Grinstein, and M. B. Wise, Phys. Rev. D **39**, 799 (1989).
- [3] N. Isgur and M. B. Wise, Phys. Lett. B **232**, 113 (1989); Phys. Lett. B **237**, 527 (1990).
- [4] M. B. Voloshin and M. A. Shifman, Yad. Fiz. **45**, 463 (1987); Yad. Fiz. **47**, 511 (1988).
- [5] S. Nussinov and W. Wetzel, Phys. Rev. D **36**, 130 (1987).
- [6] H. D. Politzer and M. B. Wise, Phys. Lett. B **206**, 681 (1988); Phys. Lett. B **208**, 504 (1988).
- [7] E. V. Shuryak, Phys. Lett. B **93**, 134 (1980).
- [8] E. Eichten and B. Hill, Phys. Lett. B **234**, 511 (1990); Phys. Lett. B **243**, 427 (1990).
- [9] H. Georgi, Phys. Lett. B **240**, 447 (1990).
- [10] B. Grinstein, Nucl. Phys. B **339**, 253 (1990).
- [11] A. F. Falk, H. Georgi, B. Grinstein, and M. B. Wise, Nucl. Phys. B **343**, 1 (1990).
- [12] A. F. Falk and B. Grinstein, Phys. Lett. B **247**, 406 (1990).
- [13] M. Luke, Phys. Lett. B **252**, 247 (1990).
- [14] A. F. Falk, B. Grinstein, and M. Luke, Nucl. Phys. B **357**, 185 (1991).
- [15] T. Mannel, W. Roberts, and Z. Ryzak, Nucl. Phys. B **355**, 38 (1991).
- [16] H. D. Politzer, Nucl. Phys. B **172**, 349 (1980).
- [17] A. F. Falk and M. Neubert, Phys. Rev. D **47**, 2965 (1993).
- [18] A. F. Falk, M. Luke, and M. J. Savage, Phys. Rev. D **49**, 3367 (1994).
- [19] A. F. Falk, Nucl. Phys. B **378**, 79 (1992).
- [20] P. Cho and B. Grinstein, Phys. Lett. B **285**, 153 (1992).
- [21] M. Neubert and V. Rieckert, Nucl. Phys. B **382**, 97 (1992).
- [22] M. Neubert, Phys. Lett. B **264**, 455 (1991).
- [23] M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. D **51**, 2217 (1995); Erratum, D **52**, 3149 (1995).
- [24] M. Neubert, Phys. Lett. B **338**, 84 (1994).
- [25] A. Czarnecki, Phys. Rev. Lett. **76**, 4124 (1996).
- [26] For earlier one-loop and one-loop improved calculations, see A. F. Falk and B. Grinstein, Phys. Lett. B **247**, 406 (1990); Phys. Lett. B **249**, 314 (1990); M. Neubert, Phys. Rev. D **46**, 2212 (1992); Phys. Rev. D **51**, 5924 (1995); Phys. Lett. B **341**, 367 (1995).
- [27] D. Buskulic *et al.* (ALEPH Collaboration), Phys. Lett. B **359**, 236 (1995).
- [28] H. Albrecht *et al.* (ARGUS Collaboration), Z. Phys. C **57**, 533 (1993).
- [29] B. Barish *et al.* (CLEO Collaboration), Phys. Rev. D **51**, 1041 (1995).
- [30] P. Abreu *et al.* (DELPHI Collaboration), CERN report CERN-PPE/96-11 (1996).
- [31] M. Neubert, Int. J. Mod. Phys. A **11**, 4173 (1996).
- [32] J. Chay, H. Georgi, and B. Grinstein, Phys. Lett. B **247**, 399 (1990).
- [33] I. I. Bigi, N. G. Uraltsev, and A. I. Vainshtein, Phys. Lett. B **293**, 430 (1992); I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. Lett. **71**, 496 (1993).
- [34] A. V. Manohar and M. B. Wise, Phys. Rev. D **49**, 1310 (1994).
- [35] A. Falk, I. Dunietz, and M. B. Wise, Phys. Rev. D **51**, 1183 (1995).

- [36] B. Guberina, R. D. Peccei, and R. Ruckl, Nucl. Phys. B **171**, 333 (1980).
- [37] M. Luke, M. J. Savage, and M. B. Wise, Phys. Lett. B **343**, 329 (1995).
- [38] Y. Nir, Phys. Lett. B **221**, 184 (1989).
- [39] M. Luke, M. J. Savage, and M. B. Wise, Phys. Lett. B **345**, 301 (1995).
- [40] A. F. Falk, M. Luke, and M. J. Savage, Phys. Rev. D **53**, 2491 (1996); Phys. Rev. D **53**, 6316 (1996).
- [41] M. Beneke and V. M. Braun, Nucl. Phys. B **426**, 301 (1994); M. Beneke, V. M. Braun, and V. I. Zakharov, Phys. Rev. Lett. **73**, 3058 (1994).
- [42] I. I. Bigi, M. A. Shifman, N. G. Uraltsev, and A. I. Vainshtein, Phys. Rev. D **50**, 2234 (1994).
- [43] M. Neubert and C. T. Sachrajda, Nucl. Phys. B **438**, 235 (1995).
- [44] M. Luke, A. V. Manohar, and M. J. Savage, Phys. Rev. D **51**, 4924 (1995).
- [45] P. Ball, M. Beneke, and V. M. Braun, Phys. Rev. D **52**, 3929 (1995).
- [46] A. F. Falk and M. E. Peskin, Phys. Rev. D **49**, 3320 (1994).
- [47] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **221**, 422 (1989).
- [48] C. K. Chow, Cornell report CLNS-96-1432 (1996).
- [49] P. Brückman (DELPHI Collaboration), DELPHI 96-95 CONF, 24 (1996).
- [50] D. Buskulic *et al.* (ALEPH Collaboration), Phys. Lett. B **365**, 437 (1996).
- [51] T. Sjöstrand, Comput. Phys. Commun. **39**, 347 (1996).
- [52] M. Feindt *et al.* (DELPHI Collaboration), CERN report CERN-PPE/95-139 (1995).
- [53] G. Brandenburg *et al.* (CLEO Collaboration), Cornell report CLEO CONF. 96-13 (1996).
- [54] A. F. Falk, Phys. Rev. Lett. **77**, 223 (1996).