

RECEIVED

CONF-9607147-3

ANL/PHY/CP-91409

OCT 30 1996

Few-valence-particle excitations around doubly magic ^{132}Sn

P.J. Daly¹, C.T. Zhang¹, P. Bhattacharyya¹, R. Broda^{1,*}, Z.W. Grabowski¹, D. Nisius¹, I. Ahmad², M.P. Carpenter², L.R. Morss², W.R. Phillips³, J.L. Durell³, M.J. Leddy³, A.G. Smith³, W. Urban³, B.J. Varley³, N. Schulz⁴, E. Lubkiewicz⁴, M. Bentalba⁴, and J. Blomqvist⁵

¹ Chemistry and Physics Depts., Purdue University, West Lafayette, IN 47907, USA

² Physics Division, Argonne National Laboratory, Argonne, IL 60439, USA

³ Dept. of Physics and Astronomy, University of Manchester, M13 9PL Manchester, UK

⁴ Centre de Recherches Nucleaires, Universite Louis Pasteur, F-67037 Strasbourg, France

⁵ Dept. of Physics Frescati, Royal Institute of Technology, S-10405 Stockholm, Sweden

Received August 6, 1996

The submitted manuscript has been created by the University of Chicago as Operator of Argonne National Laboratory ("Argonne") under Contract No. W-31-109-ENG-38 with the U.S. Department of Energy. The U.S. Government retains for itself, and others acting on its behalf, a paid-up, nonexclusive, irrevocable worldwide license in said article to reproduce, prepare derivative works, distribute copies to the public, and perform publicly and display publicly, by or on behalf of the Government.

Abstract. Prompt γ -ray cascades in neutron-rich nuclei around doubly-magic ^{132}Sn have been studied using a ^{248}Cm fission source. Yrast states located in the $N = 82$ isotones ^{134}Te and ^{135}I are interpreted as valence proton and neutron particle-hole core excitations with the help of shell model calculations employing empirical nucleon-nucleon interactions from both ^{132}Sn and ^{208}Pb regions.

PACS numbers: 21.60.Cs, 23.20.Lv, 25.85.Ca, 27.60.+j

What we know about the ^{132}Sn nucleus and its neighbors comes mainly from β^- decay studies of short-lived radionuclides produced in fission of actinides; consequently our knowledge about simple excitation modes, single particle energies, effective nucleon-nucleon interactions and other basic properties in this region is far from complete. The large multidetector γ -ray arrays, which can separate the prompt γ -ray cascades within a single fission product nucleus (of moderate yield) from the bulk of prompt γ -rays, have now opened new prospects for detailed studies of yrast excitations in ^{132}Sn and the few valence particle nuclei around it. The spectroscopy of ^{132}Sn and its neighbors should in many ways resemble that of the well studied nuclei around $Z = 82$, $N = 126$ ^{208}Pb , and comparisons of experimental data from these and other magic regions would help the development of a "universal" theoretical description of shell model properties.

The measurements reported here were performed at Eurogam II using a ^{248}Cm source which delivered $\sim 6.3 \times 10^4$ fissions/sec. Eurogam II at the time consisted of 52 escape-suppressed spectrometers incorporating 124 Ge detector elements, here augmented by four LEPS spectrometers; in total, 2×10^9 threefold or higher-fold coincidence events were recorded. The excellent quality and high selectivity of the triple coincidence γ -ray data made it possible to identify even weak transitions in the nuclei of interest extending over the $Z = 50 - 54$, $N = 80 - 84$ range. Cross coincidences observed between γ -rays from partner light and heavy fission products were often of key importance in establishing isotopic assignments for

previously unknown cascades; in other cases, some overlap with the γ -rays known from β -decay provided vital first clues. The findings for only two of these nuclei - the two- and three-proton $N = 82$ isotones ^{134}Te and ^{135}I - are presented here, but results for other fission products around ^{132}Sn will also be forthcoming.

In the two-proton nucleus ^{134}Te , many members of the $\pi g_{7/2}^2$, $\pi g_{7/2} d_{5/2}$ and $\pi g_{7/2} h_{11/2}$ multiplets are known from ^{134}Sb β^- decay studies, especially the recent work of Omtvedt et al. [1]. The present fission product measurements identified two dominant high-energy γ -rays feeding the 1691 keV $\pi g_{7/2}^2$ 6^+ state in ^{134}Te , one the 2322 keV $9^- \rightarrow 6^+$ E3 transition known from β -decay [1], the other a 2866 keV γ -ray from a ^{134}Te level at 4557 keV. Gating on this 2866 keV γ -ray revealed many new ^{134}Te γ -rays, and the full $\gamma\gamma\gamma$ results established the level sequence above 4557 keV shown to the left in the ^{134}Te scheme (Fig. 1). Since the only possible two-proton state with $I > 9$ is $(\pi h_{11/2}^2) 10^+$, expected in ^{134}Te above 7 MeV, the obvious conclusion is that these new states must involve excitation of the ^{132}Sn core. We interpret them as $\pi g_{7/2}^2 \nu f_{7/2} h_{11/2}^{-1}$ states, with strong support from shell model calculations described below.

Nothing was known up to now about high-spin states in the $N = 82$ nucleus ^{135}I , but a ^{135}Te β^- -decay study [2] has located an $11/2^+$ level at 1134 keV above the ^{135}I $\pi g_{7/2}$ ground state. In the present work, gates on 1134 keV γ -rays identified other strong γ -rays in ^{135}I , and the full $\gamma\gamma\gamma$ results established the ^{135}I level scheme presented in Fig. 1. Since no information about transition multipolarities was derived from the data, the spin-parity assignments and the interpretation of the ^{135}I levels below 4 MeV as $\pi g_{7/2}^3$, $\pi g_{7/2}^2 d_{5/2}$ and $\pi g_{7/2}^2 h_{11/2}$ states are largely based on shell model calculations for which the nucleon-nucleon interactions were taken directly from the ^{134}Te level spectrum. It is no surprise that the yrast excitations of ^{135}I are found to resemble closely those of the other three-proton nucleus ^{211}At .

The sequence of levels above 4241 keV in ^{135}I must involve core excitations, and we naturally interpret them as $\pi g_{7/2}^3 \nu f_{7/2} h_{11/2}^{-1}$ states directly related to the core-excited states in ^{134}Te . Particle-hole states of $\nu f_{7/2} h_{11/2}^{-1}$

* On leave from Institute of Nuclear Physics, Krakow, Poland

DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.




Fig. 1. The yrast level spectra established for ^{134}Te and ^{135}I . Widths of the transition arrows are proportional to the observed γ -ray intensities, except for transitions below the 164 ns isomer in ^{134}Te . Configuration assignments in both nuclei are also shown.

character with $I^\pi=2^+$ to 8^+ are known[3] in ^{132}Sn , and their energies provided some of the two-body interactions needed for calculating $\pi g_{7/2} \nu f_{7/2} h_{11/2}^{-1}$ states. In addition $\pi g_{7/2} \nu h_{11/2}^{-1}$ and $\pi g_{7/2} \nu f_{7/2}$ interactions were also needed, and these had to be estimated from corresponding multiplets in ^{208}Bi and ^{210}Bi , with scaling as $A^{-1/3}$ to take account of nuclear size variation[4]. Calculations of $\pi g_{7/2} \nu f_{7/2} h_{11/2}^{-1}$ energies were performed using the OXBASH shell model code, and the results were found to be in excellent agreement with experiment for both ^{134}Te and ^{135}I , thus providing persuasive support for the proposed interpretations.

In summary, neutron-rich fission product nuclei around doubly magic ^{132}Sn have now become accessible for detailed study by prompt γ -ray measurements using multi-detector arrays. Yrast excitations to above 5.5 MeV excitation energy in the two- and three-proton nuclei ^{134}Te and ^{135}I have been established and interpreted with the help of precise shell model calculations using empirical nucleon-nucleon interactions. These results open possibilities for exploring simple excitation modes in the ^{132}Sn

region under conditions that are comparable with but not identical to those in the well-studied ^{208}Pb region.

Acknowledgement. This work was supported by the U.S. Department of Energy under contract nos. DE-FG02-87ER40346 and W-31-109-ENG-38, by the Science and Engineering Council of the UK under grant no. GRH 71161, and by Polish Scientific Committee grant no. 1044-PO3-96-10. The authors are indebted for the use of ^{248}Cm to the Office of Basic Energy Sciences, US Dept. of Energy through the transplutonium element production facilities at the Oak Ridge National Laboratory.

References

1. J. P. Omtvedt et al., *Phys. Rev. Lett.* **75**, 3090 (1995).
2. M. Samri et al., *Z. Phys.* **A321**, 255 (1985).
3. B. Fogelberg et al., *Phys. Scr.* **T56**, 79 (1995).
4. J. Blomqvist, in *Proc. 4th Int. Conf. on Nuclei Far From Stability*, Helsingor, 1981 p. 536.

This article was processed by the author using the L^AT_EX style file *cljour2* from Springer-Verlag.