



## 26. 非閉じ込め相転移付近でのメソンの振舞い

## Mesons Above The Deconfining Transition

QCD-TARO Collaboration

Ph. de Forcrand<sup>1</sup>, M. García Pérez<sup>2</sup>, T. Hashimoto<sup>3</sup>, S. Hioki<sup>4</sup>, H. Matsufuru<sup>5,6</sup>, O. Miyamura<sup>6</sup>,  
A. Nakamura<sup>7</sup>, I.-O. Stamatescu<sup>5,8</sup>, T. Takaishi<sup>9</sup> and T. Umeda<sup>6</sup><sup>1</sup>SCSC, ETH-Zürich, CH-8092 Zürich, Switzerland<sup>2</sup>Dept. Física Teórica, Universidad Autónoma de Madrid, E-28049 Madrid, Spain<sup>3</sup>Department of Applied Physics, Faculty of Engineering, Fukui University, Fukui 910-8507, Japan<sup>4</sup>Department of Physics, Tezukayama University, Nara 631-8501, Japan<sup>5</sup>Institut für Theoretische Physik, Univ. Heidelberg D-69120 Heidelberg, Germany<sup>6</sup>Department of Physics, Hiroshima University, Higashi-Hiroshima 739-8526, Japan<sup>7</sup>Res. Inst. for Information Science and Education, Hiroshima University, Higashi-Hiroshima 739-8521, Japan<sup>8</sup>FEST, Schmeilweg 5, D-69118 Heidelberg, Germany<sup>9</sup>Hiroshima University of Economics, Hiroshima 731-01, Japan

**Abstract:** We analyze *temporal* and *spatial* meson correlators in quenched lattice QCD at  $T > 0$ . Above  $T_c$  we find different masses and (spatial) “screening masses”, signals of plasma formation, and indication of persisting “mesonic” excitations.

With increasing temperature we expect the physical picture of QCD to change according to a phase transition where chiral symmetry restoration and deconfinement may simultaneously occur. For model independent non-perturbative results one attempts lattice Monte Carlo studies [1]. Since in the Euclidean formulation the  $O(4)$  symmetry is broken at  $T > 0$  (see, e.g., [2]), physics appears different, depending on whether we probe the space (“ $\sigma$ ”:  $\mathbf{x}$ ) or time (“ $\tau$ ”:  $t$ ) direction: the string tension, e.g., measured from  $\sigma\sigma$  Wilson loops does not vanish above  $T_c$ , in contrast to the one measured from  $\sigma\tau$  loops. *Therefore we need to investigate hadronic correlators with full “space-time” structure, in particular the propagation in the Euclidean time.* The latter, however, poses special problems because of the inherently limited *physical length* of the lattice in the time direction  $l_\tau = 1/T$ . We shall discuss briefly this question and introduce our procedure.

1) Lattice problems: Large  $T$  can be achieved using small  $N_\tau = l_\tau/a$  ( $a$ : lattice spacing), however this

leads to systematic errors [3]. Moreover, having the  $t$ -propagators at only a few points makes it difficult to characterize the unknown structure in the corresponding channels. To obtain a fine  $t$ -discretization and thus detailed  $t$ -correlators, while avoiding prohibitively large lattices (we need large spatial size to avoid finite size effects, typically  $l_\sigma \sim 3l_\tau$ ), we use different lattice spacings in space and in time,  $a_\sigma/a_\tau = \xi > 1$ . For this we employ anisotropic Yang-Mills and fermionic actions [4]:

$$S_{\text{YM}} = -\frac{\beta}{3} \left( \frac{1}{\gamma_G} \text{ReTr} \square_{\sigma\sigma} + \gamma_G \text{ReTr} \square_{\sigma\tau} \right) \quad (1)$$

$$S_F = \frac{1}{2\kappa_\sigma} \bar{\Psi} W \Psi, \quad \frac{1}{\kappa_\sigma} = 2(m_0 + 3 + \gamma_F), \quad (2)$$

$$W = 1 - \kappa_\sigma \left( \sum_i \Gamma_i^+ U_i T_i + \gamma_F \Gamma_4^+ U_4 T_4 + \text{“h.c.”} \right) \quad (3)$$

( $\Gamma_\mu^\pm = 1 \pm \gamma_\mu$ ,  $\gamma_\mu^2 = 1$ ,  $T_\mu$  are lattice translation operators and  $m_0$  the bare quark mass).  $\xi$  is determined from  $T \simeq 0$  correlators  $F$  (“calibration”) by requiring isotropy in the physical distance:  $F^\sigma(z) = F^\tau(t = \xi z)$ . In a quenched simulation at some  $\gamma_G$ ,  $\xi$  is fixed by the Yang-Mills calibration, then  $\gamma_F$  is tuned to give the same  $\xi$  for  $T \simeq 0$  hadron correlators. We vary  $T = \xi/N_\tau a_\sigma$  by varying  $N_\tau$ .

2) Physical problems: Increasing the temperature is expected to induce significant changes in the struc-

ture of the hadrons (see, e.g. [5] for reviews). Two “extreme” pictures are frequently used for the intermediate and the high  $T$  regimes, respectively: the weakly interacting meson gas, where we expect the mesons to become effective resonance modes with a small mass shift and width due to the interaction; and the quark-gluon plasma (QGP), where the mesons should eventually disappear and at very high  $T$  perturbative effects should dominate. These genuine temperature effects should be reflected in the low energy structure in the mesonic channels. But this structure cannot be observed directly, due to the inherently coarse *energy* resolution  $1/l_\tau = T$ . Our strategy is the following: we fix at  $T = 0$  a mesonic source which gives a large (almost 100%) projection onto the ground state. Then we use this source to determine the changes induced by the temperature on the ground state. For  $T > T_c$  we do not have a good justification to use that source as representative of the meson but we assume that it still projects onto the dominant low energy structure in the spectral function. This is a reasonable procedure if the mesons interact weakly with other hadron-like modes in the thermal bath and the changes in the correlators are small. Large changes will signal the breakdown of this weakly interacting gas picture and there we shall try to compare our observations with the QGP picture.

On a *periodic* lattice the contribution of a pole in the mesonic spectral function to the  $t$ -propagator is  $\cosh(m(t - N_\tau/2))$  (this  $m$  is therefore called “pole-mass”). A broad structure or the admixture of excited states leads to a superposition of such terms. Fitting a given  $t$ -propagator by  $\cosh(m(t)(t - N_\tau/2))$  at pairs of points  $t, t + 1$  defines an “effective mass”  $m(t)$  which is constant if the spectral function has only one, narrow peak. We shall simply speak about  $m(t)$  as “mass”: it connects directly to the (pole) mass of the mesons below  $T_c$ , while above  $T_c$  it will help analyze the dominant low energy structure in the frame of our strategy above. By contrast, we shall speak of “screening mass” ( $m^{(\sigma)}$ ) when extracted from *spatial propagators*.  $m^{(\sigma)}$  is different from the  $T > 0$  mass (the propagation in the space directions represents a  $T = 0$  problem with finite size effects).

We use lattices of  $12^3 \times N_\tau$  with  $N_\tau = 72, 20, 16$  and 12 at  $\beta = 5.68$ ,  $\gamma_G = 4$ , no.conf = 60. We find [6]  $T_c$  at  $N_\tau$  slightly above 18, which fixes for

the above lattices  $T \simeq 0, 0.93T_c, 1.15T_c$  and  $1.5T_c$  and  $a_\tau \sim 0.044 \text{ fm} = (4.5 \text{ GeV})^{-1}$ . We calculate correlators for 3 quark masses, with  $\xi \simeq 5.3(1)$ . The various parameters are given in the Table. Details

$\kappa_\sigma$	$\gamma_F$	$m_{Ps}$	$m_V$	$a$	$p$
0.081	4.05	0.178(1)	0.196(1)	0.379	1.289
0.084	3.89	0.149(1)	0.175(1)	0.380	1.277
0.086	3.78	0.134(1)	0.164(1)	0.380	1.263

Table 1: Simulation parameters used at every  $T$  and meson masses at  $T \simeq 0$  (in units  $a_\tau^{-1}$ ). The source parameters  $a$ ,  $p$  eq. (5) are extracted from the  $T \simeq 0$  wave function.

will be given in a forthcoming paper [7]. We use periodic (anti-periodic) boundary conditions in the spatial (temporal) directions and gauge-fix to Coulomb gauge. We investigate correlators of the form:

$$G_M(\mathbf{x}, t) = \sum_{\mathbf{z}, \mathbf{y}_1, \mathbf{y}_2} w_1(\mathbf{y}_1) w_2(\mathbf{y}_2) \times \langle \text{Tr} [S(\mathbf{y}_1, 0; \mathbf{z}, t) \gamma_M \gamma_5 S^\dagger(\mathbf{y}_2, 0; \mathbf{z} + \mathbf{x}, t) \gamma_5 \gamma_M^\dagger] \rangle \quad (4)$$

$$w_{1,2}(\mathbf{y}) \sim \delta(\mathbf{y}) \text{ (point)}; \quad w_{1,2}(\mathbf{y}) \sim \exp(-a\mathbf{y}^p) \text{ (exp)} \quad (5)$$

with

$S$  the quark propagator and  $\gamma_M = \{\gamma_5, \gamma_1, 1, \gamma_1 \gamma_5\}$  for  $M = \{Ps, V, S, A\}$  (pseudoscalar, vector, scalar and axial-vector, respectively). We use point and smeared (exponential) *quark* sources and point sink. We fix the exponential (*exp*) source taking the parameters  $a, p$  in (5) from the observed dependence on  $\mathbf{x}$  of the temporal  $Ps$  correlator with *point-point* source at  $T \simeq 0$  (see Table). The results of a variational analysis using *point-point*, *point-exp* and *exp-exp* sources indicate that the latter ansatz projects practically entirely onto the ground state at  $T = 0$ . This is well seen from the effective mass in Fig. 1.

Therefore we use throughout the *exp-exp* sources in the Table, according to our strategy for defining hadron operators at high  $T$  [8]. All masses are given in units  $a_\tau^{-1}$ , i.e. we plot *mass*  $\times a_\tau$ . Errors are statistical only.

*a) Effective masses.* In Fig. 2 we show the effective mass  $m(t)$  of the  $Ps$  and  $V$  time-propagators at  $T \simeq 0.93T_c, 1.15T_c$  and  $1.5T_c$ .




Figure 1: Effective pseudoscalar mass  $m(t)$  in units of  $a_\tau^{-1}$  vs  $t$ , for various sources at  $T \simeq 0$ .




Figure 2: From left to right, effective mass  $m(t)$  in units of  $a_\tau^{-1}$  at  $T \simeq 0.93, 1.15$  and  $1.5T_c$  (open symbols) vs  $t$ . Also shown are the effective masses from the same correlators calculated using free quarks.  $\kappa_\sigma = 0.086$ .

We notice practically no change from  $T = 0$  (Fig. 1) to  $0.93T_c$ , while above  $T_c$  clear changes develop:  $m(t)$  depends strongly on  $t$ , it increases significantly, and the  $Ps$  and  $V$  reverse their positions. Because of the large changes at high  $T$  we compare here with the unbound quark picture of the high  $T$  regime of QGP. For this we calculate mesonic correlators using, with the same source, free quark propagators  $S_0$  instead of  $S$  in (4), with  $\gamma_F = \xi = 5.3$ , illustratively,  $m_0 = m_q a_\sigma = m_q a_\tau \xi = 0.1$ . Generally above  $T_c$ ,  $m(t)$  shows stronger  $t$ -dependence, which means that the spectral function selected by the *exp-exp* source no longer has just one, narrow contribution, as for  $T < T_c$ .

*b) Wave functions.* We have also analyzed the “wave functions”, i.e. the behaviour of the temporal correlators with the quark-antiquark distance at the sink. In Fig. 3 we compare the  $Ps$  wave func-




Figure 3:  $Ps$  wave functions (*Set-B*,  $\kappa_\sigma = 0.086$ , *exp-exp* source) normalized at  $r = 0$  vs quark separation  $r$  at  $t = 2, 4$  and  $6$ , using full and free propagators. Also plotted is the initial distribution of separations as given by the source,  $\int d^3y w(y+r)w(y)$ .  $T \simeq 0.93T_c$  (left) and  $T \simeq 1.5T_c$  (right).

tions normalized at  $x = 0$ ,  $G_{Ps}(x, t)/G_{Ps}(0, t)$ , at several  $t$  for  $T \simeq 0.93T_c$  (which is very similar to  $T \simeq 0$ ) and  $T \simeq 1.5T_c$  at our lightest quark mass ( $\kappa_\sigma = 0.086$ ) and for the free quark case ( $m_q a_\sigma = 0.1$ ,  $\gamma_F = \xi = 5.3$ ).

Our *exp-exp* source appears somewhat too broad at  $T \simeq 0.93T_c$ : the quarks go nearer each other while propagating in  $t$ . Interestingly enough, this is also the case at  $T \simeq 1.5T_c$ : the spatial distribution shrinks and stabilizes, indicating that even at this high temperature there is a tendency for quark and anti-quark to stay together. This is in clear contrast to the free quark case which never shows such a behaviour regardless of the source (in Fig. 3 we use the *exp-exp* source and  $m_0 = 0.1$ ; for heavy free quarks we expect a “wave function” similar to the source at all  $t$ ). Hence the only effect of the temperature on the wave function is to make it slightly broader. The same holds also for the other mesons at all quark masses.

*c) Masses and screening masses.* We fit the  $Ps$ ,  $V$ ,  $S$  and  $A$  time-correlators to single hyperbolic functions at the largest 3  $t$ -values [9]. Above  $T_c$  we assume that these masses characterize the dominant low energy structure selected by our source (e.g., the putative, unstable states described by the wave functions discussed above). We also extract screening masses from spatial correlators at the largest 3  $x$ . Since the spatial physical distance is large we use *point-point*




Figure 4: Temperature dependence of the mass  $m$  (open symbols) and screening mass  $m^{(\sigma)}$  (full symbols) in units  $a_\tau^{-1}$ ,  $\kappa_\sigma = 0.086$  (upper plot) and in the chiral limit (lower plot). The vertical gray lines indicate  $T_c$ . The data correspond to  $T \simeq 0, 0.93T_c, 1.15T_c$  and  $1.5T_c$ .

source for all  $T$  (a gauge invariant extended source leads to similar results). The results for  $m$  and  $m^{(\sigma)}$  at  $\kappa_\sigma = 0.086$  are shown in Fig. 4 ( $m$ , and to a smaller extent also  $m^{(\sigma)}$  may be overestimated).

We extrapolate  $m$  and  $m^{(\sigma)}$  in  $1/\kappa = 2m_0 + 8$  to the chiral limit from the 3 quark masses analyzed [10]. Up to  $T_c$  screening masses and masses remain similar, while above  $T_c$  the former become much larger than the latter, both at finite quark mass and in the chiral limit – see Fig. 4 [11]. In the high  $T$  regime of QGP one expects  $m^{(\sigma)} \sim 2\pi/N_\tau \sim 0.4(0.5)$  in units of  $a_\tau^{-1}$  for  $T \simeq 1.15T_c(1.5T_c)$ , respectively, to be compared with the values in Fig. 4 of  $\sim 0.3(0.4)$ .

*In conclusion*, in this quenched QCD analysis the changes of the meson properties with temperature appear to be small below  $T_c$ , while above  $T_c$  they become important and rapid, but not abrupt. Here we observe apparently opposing features: On the one hand, the behaviour of the propagators, in particular the change in the ordering of the mass splittings could be accounted for by free quark propagation [13]

in the mesonic channels above  $T_c$ , which would also explain the variation of  $m(t)$ . On the other hand, the behaviour of the wave functions obtained from the 4-point correlators suggests that there can be low energy excitations in the mesonic channels above  $T_c$ , remnants of the mesons below  $T_c$ . They would be characterized by a mass giving the location of the corresponding bump in the spectral function. The variation of  $m(t)$  with  $t$  and with the source would then indicate a resonance width increasing with  $T$ , although it may simply reflect the uncertainty in our treatment of the low energy states [14]. Remember that our source is not chosen arbitrarily but such as to reproduce a “pure” meson source at  $T = 0$ ; at high  $T$ , however, it may allow admixture of other contributions, and  $m$  becomes increasingly ambiguous. We see chiral symmetry restoration above  $T_c$  both in the masses and in the screening masses, with the latter increasing faster than the former and remaining below the free gas limit at  $T \simeq 1.5T_c$ . The exact amount of splitting among the channels and the precise ratio between  $m$  and  $m^{(\sigma)}$  may, however, be affected also by uncertainties in our  $\xi$  calibration. Finally, note that this is a quenched simulation, with incomplete dynamics.

A possible physical picture is this: Mesonic excitations subsist above  $T_c$  (up to at least  $1.5T_c$ ) as unstable modes (resonances), in interaction with unbound quarks and gluons. Our results are consistent with this, but there may be also other possibilities (cf [15, 16], cf [5] and references therein). E.g., in a study of meson propagators including dynamical quarks but without wavefunction information [16], one found masses and (spatial) screening masses  $\propto T$  above  $T_c$  and indication for QGP with “deconfined, but strongly interacting quarks and gluons”. The complex, non-perturbative structure of QGP (already indicated by equation of state studies up to far above  $T_c$  [17]) is also confirmed by our analysis of general mesonic correlators. From our study however, the detailed low energy structure of the mesonic channels appears to present further interesting, yet unsolved aspects.

Further work is needed to remove the uncertainties still affecting our analysis. This concerns particularly the  $\xi$  calibration and the question of the definition of hadron operators at high  $T$ , which appear to have been the major deficiencies, besides the smaller lat-

tices, affecting earlier results [18]. We shall also try to extract information directly about the spectral functions [19].

**Acknowledgments:** We thank JSPS, DFG and the European Network “Finite Temperature Phase Transitions in Particle Physics” for support. H.M. thanks T. Kunihiro and I.O.S. thanks F. Karsch for interesting discussions. We thank F. Karsch for reading the manuscript. We are indebted to two anonymous referees for useful comments. The calculations have been done on AP1000 at Fujitsu Parallel Comp. Res. Facilities and Intel Paragon at INSAM, Hiroshima Univ.

## References

- [1] For a recent review see: E. Laermann, Nucl. Phys. B (Proc.Suppl.) **63A-C** (1998) 114.
- [2] N.P. Landsman and Ch.G. van Weert, Phys. Rep. **145** (1987) 141.
- [3] J. Engels, F. Karsch and H. Satz, Nucl. Phys. **B205**[FS5] (1982) 239.
- [4] F. Karsch, Nucl. Phys. **B205**[FS5] (1982) 285.
- [5] H. Meyer-Ortmanns, Rev. Mod. Phys. **68** (1996) 473; H. Satz, hep-ph/9706342.
- [6] From the peak of the Polyakov loop susceptibility; see QCD-TARO: M. Fujisaki et al., Nucl. Phys. B (Proc.Suppl.) **53** (1997) 426.
- [7] QCD-TARO: Ph. de Forcrand et al., in preparation.
- [8] Source dependence has various origins. At fixed physical distance, say,  $ta_\tau = 5a_\tau$ , we find a variation in  $m(t)$  between *point-exp*, *exp-exp* and *wall* source of  $\sim 20\%$  below  $T_c$  and  $\sim 25\%$  above  $T_c$ . This source dependence disappears exponentially with  $t$  at low  $T$ , see Fig. 1. By choosing an optimal source, we try to minimize that part of the  $m(t)$  variation which is present at  $T = 0$ .
- [9] For the  $S$ -meson we only use the connected correlator.
- [10] We assume that the  $Ps$  extrapolates in mass squared for  $T < T_c$ . Above  $T_c$  the  $\kappa$  dependence is very weak. See [7].
- [11] A similar behaviour is obtained in T. Hatsuda and T. Kunihiro, Phys. Rep. **247** (1994) 221 from an NJL effective model; an increasing  $m_\pi$  is also obtained in J. Berges, D.U. Jungnickel and C. Wetterich, hep-ph/9705474.
- [12] G. Boyd, Sourendu Gupta and F. Karsch, Nucl. Phys. **B385** (1992) 481.
- [13] We mean  $S_0$  with some effective  $m_q$  induced by the interaction. The larger  $m(t)$  in  $Ps$  channel as compared with  $V$  is a peculiarity of the free Wilson quarks (3). Its counterpart for staggered fermions is an oscillating behaviour, quoted in [16] also as indicator of free quark propagation.
- [14] We remark that using free quark propagators the source dependence is much stronger.
- [15] C. DeTar and J. Kogut, Phys. Rev. **D36** (1987) 2828; C. DeTar, Phys. Rev. **D37** (1988) 2328.
- [16] G. Boyd, Sourendu Gupta, F. Karsch, E. Laermann, Z. f. Physik **C 64** (1994) 331.
- [17] G. Boyd et al., Nucl. Phys. **B 460** (1996) 419.
- [18] T. Hashimoto, A. Nakamura and I.-O. Stamatescu, Nucl. Phys. **B400** (1993) 267.
- [19] QCD-TARO: Ph. de Forcrand et al., Nucl. Phys. B (Proc.Suppl.) **63** (1998) 460.