




XJ0000210



**ОБЪЕДИНЕННЫЙ  
ИНСТИТУТ  
ЯДЕРНЫХ  
ИССЛЕДОВАНИЙ**

**Дубна**

E2-2000-176

A.P.Bakulev, S.V.Mikhailov

**NEW SHAPES OF LIGHT-CONE DISTRIBUTIONS  
OF THE TRANSVERSELY POLARIZED  $\rho$ -MESONS**

Submitted to «European Physical Journal C»

**31 / 49**

**2000**

# INTRODUCTION

In this paper, we complete our investigation of the leading twist light-cone distribution amplitudes (DAs) for lightest transversely polarized mesons with quantum numbers  $J^{PC} = 1^{--}$  ( $\rho_{\perp}$ ,  $\rho'_{\perp}$ ),  $1^{+-}$  ( $b_{1\perp}$ ) in the framework of QCD sum rules (SRs) with nonlocal condensates (NLC). These DAs are important ingredients of the “factorization” formalism [1] for any hard exclusive reactions involving  $\rho$ -mesons. For this reason, the DAs have been attractive for theorists for a long time: the main points are presented in [2, 3], a detailed revised version of the standard approach is in [4], and a generalization to the next twists is in [5]. The leading twist DA  $\varphi_{\rho, \rho', b_1}^T(x, \mu^2)$  parameterizes the matrix elements of the tensor current with transversely polarized  $\rho(770)$ - and  $\rho'(1465)$ -mesons ( $J^{PC} = 1^{--}$ )

$$\langle 0 | \bar{u}(z)\sigma_{\mu\nu}d(0) | \rho_{\perp}(p, \lambda) \rangle \Big|_{z^2=0} = i f_{\rho_{\perp}}^T \langle \varepsilon_{\mu}(p, \lambda)p_{\nu} - \varepsilon_{\nu}(p, \lambda)p_{\mu} \rangle \int_0^1 dx e^{ix(zp)} \varphi_{\rho_{\perp}}^T(x, \mu^2) + \dots, \quad (1)$$

and the  $b_1(1235)$ -meson ( $J^{PC} = 1^{+-}$ )

$$\langle 0 | \bar{u}(z)\sigma_{\mu\nu}d(0) | b_1^+(p, \lambda) \rangle \Big|_{z^2=0} = f_{b_1}^T \varepsilon_{\mu\nu\alpha\beta} \varepsilon^{\alpha}(p, \lambda) p^{\beta} \int_0^1 dx e^{ix(zp)} \varphi_{b_1}^T(x, \mu^2) + \dots \quad (2)$$

(here dots represent higher-twist contributions, explicitly defined in Appendix A, see Eqs.(A.9)-(A.10) and ref.[5]). In the above definitions,  $p_{\nu}$  and  $\varepsilon_{\mu}(p, \lambda)$  are the momentum and the polarization vector of a meson, respectively, and  $\mu^2$  is normalization point. In the framework of the standard approach, one should restrict oneself to an estimate of the second moment ( $\langle \xi^2 \rangle$ ) of the DA to restore its shape<sup>1</sup>. In other words, the variety of different DA shapes is reduced to the 1-parameter family of “admissible” DAs:  $\varphi(x; a_2) = 6x(1-x) \left[ 1 + a_2 C_2^{3/2} (2x-1) \right]$ . This family includes both the asymptotic DA ( $a_2 = 0$ ) and Chernyak-Zhitnitsky model [2] for the pion DA ( $a_2^{\pi|CZ} = -2/3$ ). For the pion case, one can think it is rather enough: most of debates (see [2, 9, 10, 11, 12] and refs. therein) about the shape of this DA are concerned just with the value of coefficient  $a_2$  – is it close to 0 or to  $a_2^{\pi|CZ}$ ? In our opinion, advocated since 1986 [9], the shape of the pion DA is not far from the asymptotic one [6, 13, 7]. Only recently, researchers have tried to extract the next Gegenbauer coefficient [12] and other parameters of the pion DA [14] from experimental data. But, in general, there is no principle to exclude a more rich structure for a hadron DA. In this case, the standard approach is definitely out of its applicability range, and one should use more refined techniques, e.g., the QCD SRs with NLC.

This work was started in [7] where the “mixed parity” NLC SR for DAs of  $\rho$ - and  $b_1$ -mesons, the particles possessing different P-parity, was analyzed. We concluded that, to obtain a reliable result, one should reduce model uncertainties due to the nonlocal gluon contribution. Separate SRs for each P-parity channel should be preferable for this purpose, and here we construct these “pure parity” SRs for corresponding DAs. The SR of this type possesses a low sensitivity to the gluon model but involves contributions from higher twists<sup>2</sup>. To construct a refined, “pure parity” SR for twist 2 DA, one must resolve the corresponding system of equations (see Appendix A). We realize this solution using the duality transformation, introduced in our previous work [15]. The negative parity NLC SR for the transversely polarized  $\rho$ -,  $\rho'$ -mesons works rather well and allows us to estimate the 2-nd, 4-th, 6-th, and 8-th moments of the leading twist DAs. The positive parity SR for the transversely polarized  $b_1$ -meson can provide only the value of the  $b_1$ -meson tensor coupling,  $f_{b_1}^T$ . We suggest the models for these DAs and check their self-consistency,

<sup>1</sup>We should note in this respect that the standard approach could not provide a reliable estimate even for the second moment of DA, see [6, 7, 8]

<sup>2</sup>as was noted in [4].

based upon both “pure” and “mixed” NLC SR. The DA shape  $\varphi_{\rho_{\pm}}^T(x)$  differs noticeably from the known one. Finally, we inspect how they can influence the  $B \rightarrow \rho e \nu$  decay form factors.

The calculation technique is the same as in [6, 7]; therefore, the corresponding details are omitted below. But, for the readers’ convenience, some important features of the NLC SRs approach would be briefly recalled. The approach introduced in [9] was successfully applied for determining light meson dynamic characteristics, DAs, form factors, see. *e.g.*, [6, 7, 13]. The original tools of NLC SR are nonlocal objects like  $M(z^2) = \langle \bar{q}(0)E(0, z)q(z) \rangle$ <sup>3</sup>, rather than  $\langle \bar{q}(0)q(0) \rangle$ . NLC  $M(z^2)$  can be expanded over the standard (local) condensates,  $\langle \bar{q}(0)q(0) \rangle$ ,  $\langle \bar{q}(0)\nabla^2 q(0) \rangle$ , and over “higher dimensions”. So, one can come back to the standard SR by truncating this series. But, in virtue of the truncation, one loses an important physical property of nonperturbative vacuum – the possibility of vacuum quarks (gluons) to flow through vacuum with a nonzero momentum  $k_{q(g)} \neq 0$ . The parameter  $\langle k_q^2 \rangle$ , fixing the average virtuality of vacuum quarks, was estimated from the mixed condensate of dimension 5 (see Appendix B, [6]),  $\langle k_q^2 \rangle = \lambda_q^2 \approx 0.4 - 0.5 \text{ GeV}^2$  [16, 17]. This value is of an order of the hadronic scale,  $\lambda_q^2 \sim m_\rho^2 \approx 0.6 \text{ GeV}^2$ , therefore the nonlocality effect can be large, and it should be taken into account in QCD SR. Since neither QCD vacuum theory exist yet, nor higher dimension condensates are estimated, it is clear that merely the models of NLC can be suggested (Appendix B). Here we apply the simplest ansatz to NLC [6, 7] that takes into account only the main effect  $\langle k_q^2 \rangle = \lambda_q^2 \neq 0$  and leads to the Gaussian decay for NLC, while the quantity  $1/\lambda_q$  reveals itself as a length of the quark-gluon correlation in QCD vacuum [6]. It is important to note that the nonlocal character of the quark condensate was recently confirmed in direct lattice calculations [18, 19]. The latter measurement in [19] confirms the validity of the Gaussian ansatz (at a small distance) as well as the value of the parameter  $\lambda_q^2$ .

## “DUALITY” TRANSFORMATION

To obtain sum rule, we start with a 2-point correlator  $\Pi^{\mu\nu;\alpha\beta}(q)$  of tensor currents  $J_{(N)}^{\mu\nu}(x) = \bar{u}(x)\sigma^{\mu\nu}(z\nabla)^N d(x)$  ( $z$  is a light-like vector,  $z^2 = 0$ ),

$$\Pi_{(N)}^{\mu\nu;\alpha\beta}(q) = i \int d^4x e^{iqx} \langle 0 | T [ J_{(0)}^{\mu\nu+}(x) J_{(N)}^{\alpha\beta}(0) ] | 0 \rangle \quad (3)$$

whose properties were partially analyzed in [3, 4, 15]. It is well known that the correlator at  $N = 0$  can be decomposed in invariant form factors  $\Pi_{\pm}$ , [3, 4]

$$\Pi_{(0)}^{\mu\nu;\alpha\beta}(q) = \Pi_{-}(q^2)P_1^{\mu\nu;\alpha\beta} + \Pi_{+}(q^2)P_2^{\mu\nu;\alpha\beta} \quad (4)$$

where the projectors  $P_{1,2}$ , obeying the projector-type relations

$$(P_i \cdot P_j)^{\mu\nu;\alpha\beta} \equiv P_i^{\mu\nu;\sigma\tau} P_j^{\sigma\tau;\alpha\beta} = \delta_{ij} P_i^{\mu\nu;\alpha\beta} \quad (\text{no sum over } i), \quad P_i^{\mu\nu;\mu\nu} = 3, \quad (5)$$

are presented in Appendix A. For the general case  $N \neq 0$ , a similar decomposition involves 4 new independent tensors  $Q_i$ ; they appear due to a new vector  $z^\alpha$  introduced into the composite tensor current operator,

$$\begin{aligned} \Pi_{(N)}^{\mu\nu;\alpha\beta}(q) &= \Pi_{-}(q^2, qz)P_1^{\mu\nu;\alpha\beta} + \Pi_{+}(q^2, qz)P_2^{\mu\nu;\alpha\beta} + K_1(q^2, qz)Q_1^{\mu\nu;\alpha\beta} \\ &+ K_3(q^2, qz)Q_3^{\mu\nu;\alpha\beta} + K_z(q^2, qz)Q_z^{\mu\nu;\alpha\beta} + K_q(q^2, qz)Q_q^{\mu\nu;\alpha\beta}. \end{aligned} \quad (6)$$

Contributions of DAs, defined in Eqs.(A.9)-(A.10), to different tensor structures in decomposition (6) are mixed, see Eqs.(A.11)-(A.12). The most effective way to disentangle them in

<sup>3</sup>Here  $E(0, z) = P \exp(i \int_0^z dt_\mu A_\mu^a(t)\tau_a)$  is the Schwinger phase factor required for gauge invariance.

practical OPE calculations is to use explicit properties of different OPE terms under the duality transformation  $\hat{D}$  (introduced in our previous work [15]) mapping any rank-4 tensor  $T^{\mu\nu;\alpha\beta}$  to another rank-4 tensor  $T_D^{\mu\nu;\alpha\beta} = (\hat{D}T)^{\mu\nu;\alpha\beta}$  with

$$D_{\mu'\nu';\alpha'\beta'}^{\mu\nu;\alpha\beta} = \frac{-1}{4} \epsilon^{\mu\nu}{}_{\mu'\nu'} \epsilon^{\alpha\beta}{}_{\alpha'\beta'} \quad \text{and} \quad \hat{D}^2 = 1. \quad (7)$$

Our projectors  $P_1^{\mu\nu;\alpha\beta}$ ,  $P_2^{\mu\nu;\alpha\beta}$ ,  $Q_1^{\mu\nu;\alpha\beta}$ ,  $Q_3^{\mu\nu;\alpha\beta}$ ,  $Q_z^{\mu\nu;\alpha\beta}$ , and  $Q_q^{\mu\nu;\alpha\beta}$  transform into each other under the action of  $\hat{D}$ :

$$(\hat{D}P_1)^{\mu\nu;\alpha\beta} = P_2^{\mu\nu;\alpha\beta}; \quad (\hat{D}Q_1)^{\mu\nu;\alpha\beta} = [P_1 + P_2 - Q_3]^{\mu\nu;\alpha\beta}; \quad (8)$$

$$(\hat{D}P_2)^{\mu\nu;\alpha\beta} = P_1^{\mu\nu;\alpha\beta}; \quad (\hat{D}Q_3)^{\mu\nu;\alpha\beta} = [P_1 + P_2 - Q_1]^{\mu\nu;\alpha\beta}; \quad (9)$$

$$(\hat{D}Q_z)^{\mu\nu;\alpha\beta} = -Q_z^{\mu\nu;\alpha\beta}; \quad (\hat{D}Q_q)^{\mu\nu;\alpha\beta} = [Q_q - Q_z + Q_1 + Q_3 - P_1 - P_2]^{\mu\nu;\alpha\beta}. \quad (10)$$

We have shown in [15] that all terms in OPE could be divided into two classes, self-dual ( $\hat{D}X_{SD} = X_{SD}$ ) and anti-self-dual ( $\hat{D}X_{ASD} = -X_{ASD}$ ). For example, the perturbative term is of ASD type, whereas the 4-quark scalar condensate contribution to OPE is of SD type.

Below we introduce the shorthand notation for contributions of DAs to decomposition (6):  $v_0$ ,  $v_1$ , and  $v_2$  stand for  $1^{--}$  ( $\rho_\perp$ ,  $\rho'_\perp$ ); and  $u_0$ ,  $u_1$ , and  $u_2$ , for  $1^{+-}$  ( $b_1$ ), see Appendix A for details. For SD parts of OPE  $u_i = -v_i$ , and the system of equations simplifies to:

$$\frac{\Pi_{\mp}(q^2, qz)}{2(qz)^N q^2} = \mp v_0 - v_1 - v_2; \quad \frac{K_{1,3}(q^2, qz)}{2(qz)^N q^2} = \mp v_1 + v_2; \quad \frac{K_q(q^2, qz)(-2K_z(q^2, qz))}{4(qz)^N q^2} = v_2, \quad (11)$$

whereas for ASD parts  $u_i = v_i$ , and we have:

$$\frac{\Pi_{\mp}(q^2, qz)}{2(qz)^N q^2} = \mp v_0 + v_1 + v_2; \quad \frac{K_{1,3}(q^2, qz)}{2(qz)^N q^2} = -v_1 - v_2; \quad \frac{K_z(q^2, qz)}{2(qz)^N q^2} = +v_2; \quad K_q(q^2, qz) = 0. \quad (12)$$

By these formulas, it is possible to determine  $\rho$ - and  $b_1$ -meson DA contributions of leading and higher twists.

## THE ‘‘MIXED PARITY’’ SUM RULE

The usual way [2, 4] to extract the moments of the function  $\varphi^T(x)$  appeals to a correlator  $J_{(N,0)}(q^2)$  of currents  $J_{(N)}^{\mu\alpha}(0)z^\alpha$  and  $J_{(0)}^{\mu\beta}(x)z^\beta$  defined as

$$-2i^n (zq)^{N+2} J_{(N,0)}(q^2) \equiv \Pi_{(N)}^{\mu\nu;\alpha\beta}(q) (z^\nu z^\beta g^{\mu\alpha}) = \frac{\Pi_-(q^2) - \Pi_+(q^2)}{q^2} (qz)^2, \quad (13)$$

the latter equality in (13) follows from (6) and Eqs.(A.7) in Appendix A. This correlator contains the contributions from states with different parity,  $\Pi_-(q^2)$  and  $\Pi_+(q^2)$  (see the analysis in [4]), therefore, the contamination from  $b_1$ -meson ( $J^{PC} = 1^{+-}$ ) in the phenomenological part of the corresponding SR is mandatory. The contamination makes it difficult to reliably extract the meson characteristics from this ‘‘mixed’’ SR.

The main feature of the theoretical part of  $J_{(N,0)}(q^2)$  is to cancel the self-dual part, represented by the four-quark condensate, in the anti-self-dual expression (13). The remaining ‘‘condensate’’ parts of  $J_{(N,0)}$  in (13) contain the same 5 universal elements  $\Delta\Phi_\Gamma(x; M^2)$  as for the  $\rho^\perp$ -,  $\pi$ -cases and, besides, an additional gluon contribution  $\Delta\Phi'_G(x; M^2)$  (see Appendix B)

that were analyzed in [7]. This term affects the values of the moments rather strong. So, here we get rid of the four-quark condensate that is not known very well due to a possible vacuum dominance violation. But, the price we pay for it is a high sensitivity to an ill-known gluon contribution  $\Delta\Phi'_G(x; M^2)$ .

The method of calculation of the NLC contributions  $\Delta\Phi_\Gamma(x; M^2)$  to the theoretical part of SR is described in [6, 7]. The corrected final results are presented in Appendix B that contains all the needed explicit expressions of  $\Delta\Phi_\Gamma(x; M^2)$  for the simplest physically motivated Gaussian ansatz. The final SR including DAs of  $\rho$ -meson and next resonances  $\rho'$  and  $b_1$  into the phenomenological (left) part is as follows:

$$\begin{aligned} (f_\rho^T)^2 \varphi_\rho^T(x) e^{-m_\rho^2/M^2} + (\rho \rightarrow \rho') + (\rho \rightarrow b_1) &= \int_0^{s_b^T} \rho_T^{\text{mixed}}(x, s; s_\rho^T, s_b^T) e^{-s/M^2} ds \\ &+ \Delta\tilde{\Phi}_G(x; M^2) + \Delta\tilde{\Phi}'_G(x; M^2) + \Delta\tilde{\Phi}_V(x; M^2) + \Delta\tilde{\Phi}_T(x; M^2), \end{aligned} \quad (14)$$

where  $s_\rho^T$  and  $s_b^T$  are the effective continuum thresholds in  $\rho$ - and  $b_1$ -channels. Recall again that the variation of the ill-known part of gluon contribution  $\Delta\tilde{\Phi}'_G(x; M^2)$  can reduce the second moment significantly [7]. In that paper, we suggest the following naive model: instead of the constant contribution  $\Delta\varphi'_G(x; M^2) \equiv (\alpha_s GG)/(6\pi M^2)$  (as in the standard approach), we put

$$\Delta\tilde{\Phi}'_G(x; M^2) = \Delta\varphi'_G(x; M^2) \frac{\theta(\Delta < x) \theta(x < 1 - \Delta)}{1 - 2\Delta}.$$

This simulation eliminates end-point ( $x = 0, 1$ ) effects due to the influence of the vacuum gluon nonlocality inspired by the analysis in [20] and our experience in the nonlocal quark case. The corresponding SR leads to estimate  $\langle \xi^2 \rangle_\rho^T = 0.329(11)$  (see Fig.2(a)). However, this value drastically changes,  $\langle \xi^2 \rangle_\rho^T \rightarrow 0.231(8)$ , if we take the local expression  $\Delta\varphi'_G(x; M^2)$  unchanged. Therefore, the estimate  $\langle \xi^2 \rangle_\rho^T = 0.329$  contains a significant model uncertainty, and the real value seems to be smaller.

Which prediction for this quantity can be obtained within the standard QCD SR approach? As one can see from Fig.2(b), the value of  $\langle \xi^2 \rangle_\rho^T$  cannot be estimated with a reasonable accuracy, because the SR **does not have real stability**. Nevertheless, the authors of [4] bravely deduce an estimate  $\langle \xi^2 \rangle_\rho^T [\text{B\&B}] = 0.27(4)$ . We discuss this attempt in comparison with processing other SRs in greater detail in section 5.

## THE ‘‘PURE PARITY’’ SUM RULES

Using the approach of Section 2, we calculate OPE terms for  $\Pi_\mp$ ,  $K_{1,3}$ , and  $K_{z,q}$  correlators and extract the contributions to DAs of the  $\rho$ - and  $b_1$ -mesons. This allows us to write down the SRs for DAs of the  $\rho$ - and  $b_1$ -mesons separately:

$$\begin{aligned} (m_\rho f_\rho^T)^2 \varphi_\rho^T(x) e^{-m_\rho^2/M^2} + (m_{\rho'} f_{\rho'}^T)^2 \varphi_{\rho'}^T(x) e^{-m_{\rho'}^2/M^2} &= \frac{1}{2} \int_0^{s_\rho^T} \rho_T^{\text{pert}}(x; s) s e^{-s/M^2} ds \\ &+ \Delta\tilde{\Phi}_G(x; M^2) + \Delta\tilde{\Phi}_S(x; M^2) + \Delta\tilde{\Phi}_V(x; M^2) + \Delta\tilde{\Phi}_T(x; M^2); \end{aligned} \quad (15)$$

$$\begin{aligned} (m_{b_1} f_{b_1}^T)^2 \varphi_{b_1}^T(x) e^{-m_{b_1}^2/M^2} &= \frac{1}{2} \int_0^{s_b^T} \rho_T^{\text{pert}}(x; s) s e^{-s/M^2} ds \\ &+ \Delta\tilde{\Phi}_G(x; M^2) - \Delta\tilde{\Phi}_S(x; M^2) + \Delta\tilde{\Phi}_V(x; M^2) + \Delta\tilde{\Phi}_T(x; M^2). \end{aligned} \quad (16)$$

where  $s_{\rho,b}^T$  are the effective continuum thresholds in the  $\rho$ - and the  $b_1$ -meson cases, respectively. The perturbative spectral density  $\rho_T^{\text{pert}}(x; s)$  is presented in an order of  $O(\alpha_s)$  in [4, 7]. Here we

also define “tilded” functions

$$\Delta \tilde{\Phi}_\Gamma(x; M^2) \equiv \frac{1}{2} M^4 \partial_{M^2} \Delta \Phi_\Gamma(x; M^2), \quad (17)$$

and the whole tensor NLC contribution

$$\Delta \tilde{\Phi}_T(x; M^2) \equiv \Delta \tilde{\Phi}_{T_1}(x; M^2) + \Delta \tilde{\Phi}_{T_2}(x; M^2) - \Delta \tilde{\Phi}_{T_3}(x; M^2). \quad (18)$$

The later noticeably differs from the case of longitudinally polarized  $\rho$ -meson due to the opposite sign of  $T_3$ -term, cf. [7]. The theoretical “condensate” part in (15)-(16) contains 5 elements obtained from the same  $\Delta \Phi_\Gamma(x; M^2)$  as for the  $\rho^L$ -meson case, whereas the self-dual four-quark contribution  $\Delta \tilde{\Phi}_S(x; M^2)$  is a new element of the analysis.

These two SRs reveal a considerably lower sensitivity to the gluon condensate contribution. To be concrete in the local limit  $\lambda_q^2 \rightarrow 0$ , the gluon part does not depend on the Borel parameter  $M^2$  at all, and its relative value is 6 times as low as that in the “mixed” SR. But the price one pays for this is high, the fidelity windows of the SRs are significantly reduced. For the  $\rho$ -meson case, fidelity windows of the Borel parameters  $M^2$  shrink to  $M^2 = 0.7 - 1.15$  GeV<sup>2</sup> (to be compared with  $M^2 = 0.75 - 2.25$  GeV<sup>2</sup> in “mixed” SR) and demand one to take into account the  $\rho'$ -meson explicitly. Here we cannot obtain the  $\rho'$ -meson mass from SR (15) because of the enhanced perturbative spectral density ( $\sim s$ ; this means that the differentiated SR has a spectral density  $\sim s^2$  and presumably, is not stable at all); instead, we use the  $\rho'$ -meson mass extracted in our previous paper on the longitudinally polarized  $\rho$ -meson DA [7],  $m_{\rho'} = 1496 \pm 37$  MeV, rather close to the Particle Data Group value  $m_{\rho'} = 1465 \pm 22$  MeV [21].

In the case of  $b_1$ -meson, one can analyze only the SR for the zeroth moment (decay constant  $f_{b_1}^T$ ) of the DA (see Fig.3), the SRs for higher moments appearing to be invalid.

## PROCESSING DIFFERENT SUM RULES AND COMPARISON OF THE RESULTS

We start with considering the results of processing both the types of SRs for  $f_\rho^T$ . Its dependence on the Borel parameter  $M^2$  obtained from the “mixed parity” NLC SR, Eq. (14), with  $s_0 = 2.9$  GeV<sup>2</sup> is shown in Fig. 1(a). Figure 1(b) shows  $f_\rho^T$  as a function of the Borel parameter  $M^2$  obtained from the “pure parity” NLC SR, Eq. (15), with  $s_0 = 2.8$  GeV<sup>2</sup>. Both kinds of SRs are rather sensitive to the  $\rho'$ -meson contribution and, for this reason, they were processed with taking it into account (see numerical results in Table 1). Solid lines correspond to the optimal thresholds  $s_0$ ; the dashed lines — to the curves with the 10-fold variation of  $\chi_{\min}^2$  (this corresponds approximately to the 5%-variation of  $s_0$ ; definition of  $\chi^2$ , see in Appendix C, Eq.(C.1)). So, one can conclude that both types of NLC SRs agree rather well about the value of  $f_\rho^T$ . Note that the presented  $f_\rho^T$  is rather close to the standard estimation  $f_\rho^T = 0.160(10)$  GeV [4] and to the lattice one  $f_{\rho}^{T, \text{Latt}}(4\text{GeV}^2) = 0.165(11)$  GeV [22], and differs significantly from the result  $f_\rho^T = 0.140$  GeV in [23].




Fig. 1:  $f_\rho^T$  as a function of the Borel parameter  $M^2$  obtained from: (a) the “mixed parity” NLC SR, Eq. (14), with  $s_0 = 2.9 \text{ GeV}^2$ ; (b) the “pure parity” NLC SR, Eq. (15), with  $s_0 = 2.8 \text{ GeV}^2$ . The fidelity windows for both figures coincide with the whole depicted range of  $M^2$ .

Type of SR	Table 1: The moments $\langle \xi^N \rangle_M(\mu^2)$ at $\mu^2 \sim 1 \text{ GeV}^2$ (errors are depicted in brackets in a standard manner)				
	$f_M(1\text{GeV}^2)$	$N = 2$	$N = 4$	$N = 6$	$N = 8$
Asympt. WF	1	0.2	0.086	0.047	0.030
NLC SR Eq.(15) : $\rho^T$	0.157(5)	0.296(20)	0.196(6)	0.132(5)	0.089(4)
NLC SR Eq.(14) : $\rho^T$	0.162(5)	0.329(11)	-	-	-
B&B SR : $\rho^T$	0.160(10)	0.304(40) <sup>4</sup>	does not work		
NLC SR Eq.(15) : $\rho^T$	0.140(10)	0.086(6)	0.010(1)	0.013(1)	0.022(2)
NLC SR Eq.(16) : $b_1^T$	0.184(5)	does not work			
NLC SR Eq.(14) : $b_1^T$	0.181(5)	0.144(15)	-	-	-
B&B SR : $b_1^T$	0.175(5)	does not work			

Now we consider the results of processing SRs for the second moment  $\langle \xi^2 \rangle_\rho^T$ . First, we demonstrate the results of the “standard” approach:  $\langle \xi^2 \rangle_\rho^T$  from Eq.(3.21) in [4] as a function of  $M^2$  is shown in Fig.2(b) by a long-dashed line. This curve is not stable in  $M^2$  at all, therefore the SR can provide merely a range of admissible values,  $0.27 \leq \langle \xi^2 \rangle_\rho^T \leq 0.4$ . As it is evident from Fig. 2, this wide window agrees reasonably with both the estimates from the “mixed” (a) and “pure” (b) NLC SRs.

Note, the authors of [4] dealt with the quantity  $a_2$ , the Gegenbauer coefficient in the expansion of DA. The second moment of DA is trivially connected with this coefficient,  $\langle \xi^2 \rangle = 0.2 + (12/35)a_2$ . Using the SR of [4] for  $a_2$ , we obtain the corresponding window,  $0.2 \leq a_2 \leq 0.4$ , that leads to the mean value  $\langle \xi^2 \rangle_\rho^T[\text{Stand}] = 0.30$  being surprisingly close to our estimate

<sup>4</sup>The estimate presented in this cell has been obtained by processing the “mixed parity” SR established in [4], whereas in the original paper [4] this value amounts to 0.27(4).




Fig. 2:  $\langle \xi^2 \rangle_\rho^T$  as a function of the Borel parameter  $M^2$  obtained from: (a) the “mixed parity” NLC SR, Eq. (14), with  $s_0 = 2.9 \text{ GeV}^2$ ; (b) the “pure parity” NLC SR, Eq. (15), with  $s_0 = 2.8 \text{ GeV}^2$ . Both kinds of SRs were processed with taking the  $\rho'$ -meson into account, the arrows show the fidelity window (for figure (a) the window coincides with the whole depicted range of  $M^2$ ). Solid lines correspond to the optimal thresholds  $s_0$ , the short-dashed lines on both the figures correspond to the curves with the 10%-variation of  $s_0$  (a) or of  $\chi_{\min}^2$  (b). The long-dashed line in figure (b) represents the SR of Ball-Braun [4].

from NLC SRs, (see Table 1). However Ball and Braum have obtained the erroneous estimate  $a_2 = 0.2 \pm 0.1$  producing, instead, the mean value  $\langle \xi^2 \rangle_\rho^T [\text{B\&B}] = 0.27$ .

The curves for the next higher moments, whose estimates are presented in Table 1, have the fidelity windows and the stability behavior similar to  $\langle \xi^2 \rangle_\rho^T(M^2)$  in Fig.2(b). Finally, in Fig.3, we demonstrate a very good correspondence between the values of  $f_{b_1}^T$  obtained in different NLC SRs.




Fig. 3: The curves  $f_{b_1}^T$  in  $M^2$  obtained from: (a) the “mixed parity” NLC SR (with taking the  $\rho'$ -meson into account with  $f_{\rho'}$ , defined from “pure parity” SR (15)); (b) the “pure parity” NLC SR (16). The arrows show the fidelity window (for the right figure, the window coincides with the whole depicted range of  $M^2$ ). Solid lines correspond to the optimal thresholds; the short-dashed lines on both the figures, to the curves with the 10-fold variation of  $\chi_{\min}^2$ ; the long-dashed line on the right figure corresponds to the real B&B curve.

## DA MODELS AND THEIR APPLICATION TO EXCLUSIVE PROCESSES

Possible models of DAs corresponding to the moments in Table 1 are of the form

$$\begin{aligned} \varphi_\rho^{T,mod}(x, \mu^2) &= 1.382 \{ \varphi^{as}(x) \}^2 \left( 1 + 0.927 C_2^{3/2}(\xi) + 0.729 C_4^{3/2}(\xi) \right) \\ &= \varphi^{as}(x) \left( 1 + 0.29 C_2^{3/2}(\xi) + 0.41 C_4^{3/2}(\xi) - 0.32 C_6^{3/2}(\xi) \right), \end{aligned} \quad (19)$$

$$\varphi_{\rho'}^{T,mod}(x, \mu^2) = \varphi^{as}(x) \left( 1 - 0.339C_2^{3/2}(\xi) + 0.003C_4^{3/2}(\xi) + 0.192C_6^{3/2}(\xi) \right), \quad (20)$$

$$\varphi_{b_1}^{mod}(x, \mu^2) = \varphi^{as}(x) \left( 1 - (0.175 \pm 0.05)C_2^{3/2}(\xi) \right). \quad (21)$$

where  $\xi \equiv 1 - 2x$ ,  $C_n^\nu(\xi)$  are the Gegenbauer polynomials (GP), and the norm  $\mu^2 \simeq 1 \text{ GeV}^2$  corresponds to a mean value of  $M^2$ . Recall again that the value of the important coefficient  $a_2 = 0.29$  in (19) is confirmed by 3 sources: “pure” NLC SR (15), “mixed” NLC SR (14), and a mean value from the “mixed” standard SR. Figures 4, 5(a) contain curves of DA corresponding




Fig. 4: The curves of  $\varphi_{\rho'}^{T,mod}(x, 1 \text{ GeV}^2)$ : (a) Solid lines correspond to the best fits for determined moments (see Table 1); the dashed line on the left figure corresponds to the B&B curve (which fits only  $\langle \xi^2 \rangle_{\rho}^T \simeq 0.27$ ). (b) The rhs of Eq.(15) SR $_{\rho'}^T(x, M^2)$  in  $x$ . Solid and dashed lines here correspond to different values of Borel  $M^2$ .

to  $\rho_{\perp}$ , eqs. (19), and  $\rho'_{\perp}$  (20). The arising 3-hump shape of DA for  $\rho_{\perp}$  drastically differs from that obtained in [4] and from the one obtained in chiral effective theory [23].

This difference mainly appears due to the higher moments,  $N = 4, 6, 8$ , involved into consideration. Nevertheless, the hump shape is not an artifact of the GP expansion series truncation. These models really contain only 3 first GPs, meanwhile, it is enough to reproduce all 4 moments up to  $N = 8$ . Moreover, an additionally smoothed<sup>5</sup> rhs of the NLC SR (15) demonstrates qualitatively the same behaviour in  $x$  (at admissible  $M^2$ ) as the model DA, compare Figs. 4(a) and (b).

Inverse moments of DAs often appear in perturbative QCD predictions for exclusive reactions. The estimates for important  $\langle x^{-1} \rangle_M$  moments obtained from the model DAs are presented here<sup>6</sup>

$$\langle x^{-1} \rangle_{\rho} \equiv \int_0^1 \frac{\varphi_{\rho}^T(x, 1 \text{ GeV}^2)}{x} dx = \begin{cases} 4.15_{-0.1}^{+0.4} & \text{(here)} \\ 3.6 & \text{(B\&B model)} \end{cases} \quad (22)$$

$$\langle x^{-1} \rangle_{\rho'} \equiv \int_0^1 \frac{\varphi_{\rho'}^T(x, 1 \text{ GeV}^2)}{x} dx = 2.57 \pm 0.20 \text{ (here)} \quad (23)$$

$$\langle x^{-1} \rangle_{b_1} \equiv \int_0^1 \frac{\varphi_{b_1}^T(x, 1 \text{ GeV}^2)}{x} dx = 2.48 \pm 0.20 \text{ (here)} \quad (24)$$

It is useful to construct **an independent** SR for these inverse moments to verify the DA models (19, 20, 21). Namely, the weighted sum  $C(M^2)$  of these moments

$$C(M^2) \equiv \langle x^{-1} \rangle_{\rho} + \langle x^{-1} \rangle_{\rho'} \left( \frac{f_{\rho'}^T}{f_{\rho}^T} \right)^2 e^{-(m_{\rho'}^2 - m_{\rho}^2)/M^2} + \langle x^{-1} \rangle_{b_1} \left( \frac{f_{b_1}^T}{f_{\rho}^T} \right)^2 e^{-(m_{b_1}^2 - m_{\rho}^2)/M^2} \quad (25)$$

<sup>5</sup>A certain smoothing of some  $\delta$ -functions in the rhs of the SR (see Appendix B) is not important.

<sup>6</sup>The upper error +0.4 in (22) corresponds to an overestimate  $\langle \xi^2 \rangle = 0.329$  from the “mixed” SR

can be obtained by integrating the rhs of the “mixed” NLC SR (14) with the weight  $1/x$ . A comparison of the function  $C(M^2)$  with the corresponding combination of model estimates (22, 23, 24) obtained in different kinds of NLC SRs (mainly from the “pure” ones) leads to an approximate equation

$$4.15 + 2.57 \left( \frac{f_{\rho'}^T}{f_{\rho}^T} \right)^2 e^{-(m_{\rho'}^2 - m_{\rho}^2)/M^2} + 2.48 \left( \frac{f_{b_1}^T}{f_{\rho}^T} \right)^2 e^{-(m_{b_1}^2 - m_{\rho}^2)/M^2} \approx C(M^2), \quad (26)$$

illustrated in Fig. 5(b).




Fig. 5: (a) The curve of  $\varphi_{\rho'}^{T,mod}(x, 1 \text{ GeV}^2)$  in  $x$ . (b)  $C(M^2)$  as a function of  $M^2$  (solid line) determined by Eq.(25) and integrating in  $x$  Eq.(14) in comparison with the lhs of Eq.(26) (long-dashed line). The dotted line corresponds to  $(x^{-1})_{\rho} = 4.15$ ; whereas the dashed line – to the lhs of Eq.(26) with upper values of corresponding moments.

As a result, one can conclude:

1. The “mixed” NLC SR is highly sensitive to  $b_1$ - and  $\rho'$ -meson contributions, the difference in the behavior of  $C(M^2)$  (solid line) and in the  $\rho$ -contribution alone (dotted line) illustrates this point.
2. The curve  $C(M^2)$  lies between mean and upper estimates for the lhs of (26), so it is in reasonable agreement with the estimates (22, 23, 24). It also demonstrates an overestimation of DA moments in the “mixed” SR as compared to that obtained from the “pure” one.

The new DA shapes result in different pQCD predictions for exclusive reactions with the  $\rho$ -meson. As an example, we re-estimate form factors  $V(t)$ ,  $A_{1,2}(t)$  of the process  $B \rightarrow \rho e \nu$ , in the framework of the light-cone SR approach. That was done earlier by Ball and Braun in [24], [25] on the base of DAs from [4]. Our form factors are slightly higher than those in [24] and possess a much better accuracy (compare the  $\chi^2$  in (28)). The source of the difference can be traced to the difference of the estimates like (22) for the simplest integrals. Below, the form factor values are written at a zero momentum transfer ( $t = 0$ ) as compared with B&B results:

$$\begin{aligned} V(0) &= \begin{cases} 0.37(1) & (\text{here } [s_0 = 50 \text{ GeV}^2], \chi^2 \approx 0.4) \\ 0.35(2) & ([24] [s_0 = 34 \text{ GeV}^2], \chi^2 \approx 3.4) \end{cases} \\ A_1(0) &= \begin{cases} 0.283(4) & (\text{here } [s_0 = 45 \text{ GeV}^2], \chi^2 \approx 0.1) \\ 0.27(1) & ([24] [s_0 = 34 \text{ GeV}^2], \chi^2 \approx 1.1) \end{cases} \\ A_2(0) &= \begin{cases} 0.30(1) & (\text{here } [s_0 = 50 \text{ GeV}^2], \chi^2 \approx 0.2) \\ 0.28(1) & ([24] [s_0 = 34 \text{ GeV}^2], \chi^2 \approx 1.1) \end{cases} \end{aligned} \quad (27)$$

# CONCLUSION

Let us summarize the main results of this paper:

1. We construct NLC SRs for DA for each P-parity channels, based on the properties of the duality transformation. The negative parity NLC SR for transversely polarized  $\rho$ -,  $\rho'$ -mesons works rather well and allows us to estimate the 2-nd, 4-th, 6-th, and 8-th moments of the leading twist DAs. The positive parity SR for the transversely polarized  $b_1$ -meson can provide only the value of the  $b_1$ -meson lepton decay constant,  $f_{b_1}^T$ . It should be emphasized that an analogous evaluation of the moments within the standard QCD SR approach is impossible.
2. Results of processing different NLC SRs of the “pure” (see Figs. 1b, 2b, 3b) and “mixed” (see Figs. 1a, 2a, 3a) parity are compared, and a reasonable agreement between them is found. The “mixed” SR in the standard version admits merely a window of possible values of the second moment  $\langle \xi^2 \rangle$  (see, *e.g.*, [4]); the position of the window is corrected here and, as a result, agrees with the NLC SR results presented in Table 1.
3. The models for the leading twist DAs of the  $\rho_\perp$ - and  $\rho'_\perp$ -mesons, (19,20), and of the  $b_1^\perp$ -meson, (21), are suggested. The shape of a new  $\rho_\perp$ -meson distribution (see Fig. 4a) drastically differs from that obtained by Ball and Braun [4] only on the basis of the value  $a_2 = 0.2$ . It should be emphasized that the Ball-Braun SR for the coefficient  $a_2$ , Eq.(3.20) of [4], produces the estimate  $a_2 = 0.3 \pm 0.1$ .
4. We estimate important integrals appearing in perturbative QCD predictions for different exclusive reactions,  $\langle x^{-1} \rangle_M \equiv \int_0^1 \frac{\varphi_M^T(x)}{x} dx$  in (22)-(24), based on our results for the DA shapes. We check the self-consistency of these results by comparing them with those obtained from an independent “mixed” QCD SR for the inverse moment  $\langle x^{-1} \rangle_M$  and find an agreement.
5. Form factors of the process  $B \rightarrow \rho e \nu$ ,  $V(0)$ ,  $A_{1,2}(0)$  at a zero momentum transfer are also re-estimated in the framework of the light-cone SR approach [24] on the basis of the new model for the  $\rho$ -meson DAs; the results are slightly higher and have uncertainties a few times as small as those obtained by Ball and Braun.

Finally, we can conclude that the nonlocal condensate QCD SR approach to distribution amplitudes is self-consistent and gives reliable results. An open problem of this approach is to determine well-established models of distribution functions  $f_T(\nu)$  from the theory of nonperturbative QCD vacuum. First direct attempts to calculate quark NLC have been done in lattice simulations in [18]. The “short distance” correlation length of NLC has also been extracted later in [19]; it turns out to be reasonably close to the value of  $1/\lambda_q$  and confirms the validity of our Gaussian NLC model.

## Acknowledgments

This work was partially supported by the Russian Foundation for Basic Research (grant N 00-02-1669) and Heisenberg-Landau Program. We are grateful to O. Teryaev, R. Ruskov, M. Polyakov, and N. G. Stefanis for fruitful discussions. We gratefully acknowledge the warm hospitality of Prof. K. Goeke and Dr. Dr. N. G. Stefanis at Bochum University, where this work was completed.

## Appendix

### A DECOMPOSITION OF RANK-4 TENSOR $\Pi_{(N)}^{\mu\nu;\alpha\beta}$

$$P_1^{\mu\nu;\alpha\beta} \equiv \frac{1}{2q^2} \left[ g^{\mu\alpha} q^\nu q^\beta - g^{\nu\alpha} q^\mu q^\beta - g^{\mu\beta} q^\nu q^\alpha + g^{\nu\beta} q^\mu q^\alpha \right]; \quad (\text{A.1})$$

$$P_2^{\mu\nu;\alpha\beta} \equiv \frac{1}{2} \left[ g^{\mu\alpha} g^{\nu\beta} - g^{\mu\beta} g^{\nu\alpha} \right] - P_1^{\mu\nu;\alpha\beta}; \quad (\text{A.2})$$

$$Q_1^{\mu\nu;\alpha\beta} \equiv \frac{1}{2(qz)} \left[ g^{\mu\alpha} q^\nu z^\beta + g^{\nu\beta} q^\mu z^\alpha - g^{\mu\beta} q^\nu z^\alpha - g^{\nu\alpha} q^\mu z^\beta \right]; \quad (\text{A.3})$$

$$Q_3^{\mu\nu;\alpha\beta} \equiv \frac{1}{2(qz)} \left[ g^{\mu\alpha} z^\nu q^\beta + g^{\nu\beta} z^\mu q^\alpha - g^{\mu\beta} z^\nu q^\alpha - g^{\nu\alpha} z^\mu q^\beta \right]; \quad (\text{A.4})$$

$$Q_z^{\mu\nu;\alpha\beta} \equiv \frac{q^2}{2(qz)^2} \left[ g^{\mu\alpha} z^\nu z^\beta + g^{\nu\beta} z^\mu z^\alpha - g^{\mu\beta} z^\nu z^\alpha - g^{\nu\alpha} z^\mu z^\beta \right]; \quad (\text{A.5})$$

$$Q_q^{\mu\nu;\alpha\beta} \equiv \frac{1}{2(qz)^2} \left( q^\alpha z^\beta - q^\beta z^\alpha \right) \left( q^\mu z^\nu - q^\nu z^\mu \right). \quad (\text{A.6})$$

$$g^{\mu\alpha} z^\nu z^\beta P_1^{\mu\nu;\alpha\beta} \equiv P_1^{\mu z;\mu z} = -P_2^{\mu z;\mu z} = \frac{(qz)^2}{q^2}; \quad Q_1^{\mu z;\mu z} = Q_3^{\mu z;\mu z} = Q_z^{\mu z;\mu z} = Q_q^{\mu z;\mu z} = 0; \quad (\text{A.7})$$

$$q^\mu q^\alpha z^\nu z^\beta P_1^{\mu\nu;\alpha\beta} \equiv P_1^{qz;qz} = Q_1^{qz;qz} = Q_3^{qz;qz} = -Q_q^{qz;qz} = -\frac{(qz)^2}{2}; \quad P_2^{qz;qz} = Q_z^{qz;qz} = 0. \quad (\text{A.8})$$

Let us write down the parameterization of matrix elements of a composite tensor current operator, see, e.g., [25]:

$$\begin{aligned} \langle 0 | \bar{d}(z) \sigma_{\mu\nu} u(0) | \rho_\perp(p, \lambda) \rangle \Big|_{z^2=0} &= i f_{\rho_\perp}^T \left[ (\varepsilon_\mu(p, \lambda) p_\nu - \varepsilon_\nu(p, \lambda) p_\mu) \int_0^1 dx \varphi_\rho^T(x) e^{ix(zp)} \right. \\ &\quad + (\varepsilon_\mu(p, \lambda) z_\nu - \varepsilon_\nu(p, \lambda) z_\mu) p^2 \int_0^1 dx V_1(x) e^{ix(zp)} \\ &\quad \left. + (p^\mu z_\nu - p^\nu z_\mu) (\varepsilon(p, \lambda) z) p^2 \int_0^1 dx V_2(x) e^{ix(zp)} \right] \quad (\text{A.9}) \end{aligned}$$

$$\begin{aligned} \langle 0 | \bar{d}(z) \sigma_{\mu\nu} u(0) | b_1(p, \lambda) \rangle \Big|_{z^2=0} &= f_{b_1}^T \left[ \epsilon_{\mu\nu\alpha\beta} \varepsilon^\alpha(p, \lambda) p^\beta \int_0^1 dx \varphi_{b_1}(x) e^{ix(zp)} \right. \\ &\quad + \epsilon_{\mu\nu\alpha\beta} \varepsilon^\alpha(p, \lambda) z^\beta p^2 \int_0^1 dx U_1(x) e^{ix(zp)} \\ &\quad \left. + \epsilon_{\mu\nu\alpha\beta} p^\alpha z^\beta (\varepsilon(p, \lambda) z) p^2 \int_0^1 dx U_2(x) e^{ix(zp)} \right] \quad (\text{A.10}) \end{aligned}$$

Here we decode our shorthand notation used in Section 2:

$$\begin{aligned} v_0 &\equiv \left| f_{\rho_\perp}^T \right|^2 \langle x^N \rangle_{\rho_\perp}; & v_1 &\equiv \left| f_{\rho_\perp}^T \right|^2 \langle -iN x^{N-1} \rangle_{v_1}; & v_2 &\equiv \left| f_{\rho_\perp}^T \right|^2 \langle -N(N-1)x^{N-2} \rangle_{v_2}; \\ u_0 &\equiv \left| f_{b_1}^T \right|^2 \langle x^N \rangle_{b_\perp}; & u_1 &\equiv \left| f_{b_1}^T \right|^2 \langle -iN x^{N-1} \rangle_{u_1}; & u_2 &\equiv \left| f_{b_1}^T \right|^2 \langle -N(N-1)x^{N-2} \rangle_{u_2}, \end{aligned}$$

(with  $\langle f(x) \rangle_U \equiv \int_0^1 dx f(x)U(x)$ ). In the general case, the whole system of equations for different twist DA contributions is of the following form

$$\frac{\Pi_-(q^2, qz)}{2(qz)^N q^2} = -v_0 + u_1 + u_2; \quad \frac{K_1(q^2, qz)}{2(qz)^N q^2} = -v_1 - u_2; \quad \frac{K_2(q^2, qz)}{2(qz)^N q^2} = +u_2; \quad (\text{A.11})$$

$$\frac{\Pi_+(q^2, qz)}{2(qz)^N q^2} = +u_0 + u_1 + u_2; \quad \frac{K_3(q^2, qz)}{2(qz)^N q^2} = -u_1 - u_2; \quad \frac{K_4(q^2, qz)}{2(qz)^N q^2} = v_2 - u_2. \quad (\text{A.12})$$

## B EXPRESSIONS FOR NONLOCAL CONTRIBUTIONS TO SUM RULES

For vacuum distribution functions  $f_\Gamma(\nu)$ , we use the set of the simplest ansatzes

$$f_S(\nu) = \delta(\nu - \lambda_q^2/2); \quad f_V(\nu) = \delta'(\nu - \lambda_q^2/2); \quad (\text{B.1})$$

$$f_{T_{1,2,3}}(\alpha_1, \alpha_2, \alpha_3) = \delta(\alpha_1 - \lambda_q^2/2) \delta(\alpha_2 - \lambda_q^2/2) \delta(\alpha_3 - \lambda_q^2/2). \quad (\text{B.2})$$

Their meaning and relation to initial NLC's have been discussed in [9, 6]. The contributions of NLC's  $\Delta\Phi_\Gamma(x, M^2)$  corresponding to these ansatzes are shown below; the limit of these expressions to the standard (local) contributions  $\varphi_\Gamma(x, M^2)$   $\lambda_q^2 \rightarrow 0$ ,  $\Delta\Phi_\Gamma(x, M^2) \rightarrow \Delta\varphi_\Gamma(x, M^2)$  are also written for comparison. Hereafter  $\Delta \equiv \lambda_q^2/(2M^2)$ ,  $\bar{\Delta} \equiv 1 - \Delta$ :

$$\Delta\Phi_S(x, M^2) = \frac{A_S}{M^4} \frac{18}{\bar{\Delta}\Delta^2} \{ \theta(\bar{x} > \Delta > x) \bar{x} [x + (\Delta - x) \ln(\bar{x})] + (\bar{x} \rightarrow x) + \theta(1 > \Delta) \theta(\Delta > x > \bar{\Delta}) [\bar{\Delta} + (\Delta - 2\bar{x}x) \ln(\Delta)] \}, \quad (\text{B.3})$$

$$\Delta\varphi_S(x, M^2) = \frac{A_S}{M^4} 9 (\delta(x) + (\bar{x} \rightarrow x));$$

$$\Delta\Phi_V(x, M^2) = \frac{A_S}{M^4} (x\delta'(\bar{x} - \Delta) + (\bar{x} \rightarrow x)), \quad (\text{B.4})$$

$$\Delta\varphi_V(x, M^2) = \frac{A_S}{M^4} (x\delta'(\bar{x}) + (\bar{x} \rightarrow x)); \quad (\text{B.5})$$

$$\Delta\Phi_{T_1}(x, M^2) = -\frac{3A_S}{M^4} \theta(1 > 2\Delta) \left\{ [\delta(x - 2\Delta) - \delta(x - \Delta)] \left( \frac{1}{\Delta} - 2 \right) + \theta(2\Delta > x) \cdot \theta(x > \Delta) \frac{\bar{x}}{\bar{\Delta}} \left[ \frac{x - 2\Delta}{\Delta\bar{\Delta}} \right] \right\} + (\bar{x} \rightarrow x), \quad (\text{B.6})$$

$$\Delta\varphi_{T_1}(x, M^2) = \frac{3A_S}{M^4} (\delta'(\bar{x}) + (\bar{x} \rightarrow x));$$

$$\Delta\Phi_{T_2}(x, M^2) = \frac{4A_S}{M^4} \bar{x} \theta(1 > 2\Delta) \left\{ \frac{\delta(x - 2\Delta)}{\Delta} - \theta(2\Delta > x) \theta(x > \Delta) \cdot \frac{1 + 2x - 4\Delta}{\bar{\Delta}\Delta^2} \right\} + (\bar{x} \rightarrow x), \quad (\text{B.7})$$

$$\Delta\varphi_{T_2}(x, M^2) = -\frac{2A_S}{M^4} (x\delta'(\bar{x}) + (\bar{x} \rightarrow x));$$

$$\Delta\Phi_{T_3}(x, M^2) = \frac{3A_S \bar{x}}{M^4 \bar{\Delta} \Delta} \left\{ \theta(2\Delta > x) \theta(x > \Delta) \theta(1 > 2\Delta) \left[ 2 - \frac{\bar{x}}{\Delta} - \frac{\Delta}{\bar{\Delta}} \right] \right\} + (\bar{x} \rightarrow x), \quad (\text{B.8})$$

$$\Delta\varphi_{T_3}(x, M^2) = \frac{3A_S}{M^4} (\delta(\bar{x}) + (\bar{x} \rightarrow x));$$

$$\Delta\Phi_G(x, M^2) = \frac{\langle \alpha_s GG \rangle}{24\pi M^2} (\delta(x - \Delta) + (\bar{x} \rightarrow x)), \quad (\text{B.9})$$

$$\begin{aligned}
\Delta\varphi_G(x, M^2) &= \frac{\langle\alpha_s GG\rangle}{24\pi M^2} (\delta(\bar{x}) + (\bar{x} \rightarrow x)); \\
\Delta\Phi'_G(x, M^2) &= \frac{\langle\alpha_s GG\rangle}{6\pi M^2} \frac{\theta(\Delta < x)\theta(x < 1 - \Delta)}{1 - 2\Delta}; \\
\Delta\varphi'_G(x, M^2) &= \frac{\langle\alpha_s GG\rangle}{6\pi M^2}.
\end{aligned}
\tag{B.10}$$

Here  $A_s = \frac{8\pi}{81} (\sqrt{\alpha_s} \bar{q}(0) q(0))^2$ ; for quark and gluon condensates, we use the standard estimates  $(\sqrt{\alpha_s} \bar{q}(0) q(0)) \approx (-0.238 \text{ GeV})^3$ ,  $\frac{\langle\alpha_s GG\rangle}{12\pi} \approx 0.001 \text{ GeV}^4$  [26] and  $\lambda_q^2 = \frac{\langle\bar{q}(i g \sigma_{\mu\nu} G^{\mu\nu}) q\rangle}{2\langle\bar{q}q\rangle} = 0.4 \pm 0.1 \text{ GeV}^2$  normalized at  $\mu^2 \approx 1 \text{ GeV}^2$ .

**Expressions for perturbative spectral density:** Radiative corrections reach 10 % of the Born result at  $s \sim 1 \text{ GeV}^2$ .

$$\rho_T^{\text{pert}}(x, s) = \frac{3}{2\pi^2} x \bar{x} \left\{ 1 + \frac{\alpha_s(\mu^2) C_F}{4\pi} \left( 2 \ln \left[ \frac{s}{\mu^2} \right] + 6 - \frac{\pi^2}{3} + \ln^2(\bar{x}/x) + \ln(x\bar{x}) \right) \right\} \tag{B.11}$$

Here  $\mu^2 \sim 1 \text{ GeV}^2$  corresponds to the average value of the Borel parameter  $M^2$  in the stability window;  $\alpha_s(1 \text{ GeV}^2) \approx 0.52$ . We also use the ‘mixed’ perturbative spectral density suggested in [27] in the ‘mixed’ SR:

$$\rho_T^{\text{mixed}}(x, s; s_\rho^T, s_b^T) \equiv \rho_T^{\text{pert}}(x; s) \frac{1}{2} \left[ \theta(s_\rho^T - s) + \theta(s_b^T - s) \right]. \tag{B.12}$$

## C ABOUT $\chi^2$ -DEFINITION IN SUM RULES

Let us discuss the definition of  $\chi^2$  for the SR case. We have here the function  $F(M^2, s)$ , and the problem is to find the best value  $s_0$ , such that  $F(M^2, s_0)$  is the most close to a constant value for  $M_-^2 \leq M^2 \leq M_+^2$  (values of  $M_\pm^2$  are known and fixed from standard constraints of QCD SR, see [26, 7]). We define the function  $\chi^2(s)$  for the curve  $F(M^2, s)$  with  $M^2 \in [M_-^2, M_+^2]$  in the following manner:

$$\chi^2(s) \equiv \frac{1}{(N-1)\epsilon^2} \sum_{k=0}^N \left[ F(M_-^2 + k\delta, s) - \frac{1}{N+1} \sum_{k=0}^N F(M_-^2 + k\delta, s) \right]^2, \tag{C.1}$$

where  $\delta = (M_+^2 - M_-^2)/N$ ,  $N \simeq 10$ , and  $\epsilon$  is of an order of the last decimal digit in  $F(M^2, s)$  we are interested in (in the case of decay constant  $f_\rho \approx 200 \text{ MeV}$ ,  $\epsilon \approx 1 \text{ MeV}$ ; in the case of the second moment  $\langle\xi^2\rangle_\rho \approx 0.25$ ,  $\epsilon \approx 0.01$ ). Then, if we obtain  $\chi^2(s_0) \approx 1$ , this tells us that the mean deviation of  $F(M^2, s_0)$  from a constant value in the region  $[M_-^2, M_+^2]$  is about  $\epsilon$ . To find the minimum value of  $\chi^2$  and the corresponding  $s_0$ , we used the code Mathematica.

## References

- [1] S. J. Brodsky, G. P. Lepage, *Phys. Lett.* **B87**, 359 (1979);  
A. V. Efremov, A. V. Radyushkin, *Phys. Lett.* **94B**, 245 (1980)
- [2] V. L. Chernyak, A. R. Zhitnitsky, *Nucl. Phys.* **B201** (1982) 492; **B214** (1983) 547(E);  
*Phys. Rept* **112** (1984) 173
- [3] J. Govaerts, L. J. Rubinstein, F. De Viron, J. Weyers, *Nucl. Phys.* **B283** (1987) 706.

- [4] Patricia Ball, V. M. Braun, *Phys. Rev.* **D54** (1996) 2182
- [5] Patricia Ball, V. M. Braun, Y. Koike, K. Tanaka, *Nucl. Phys.* **B529** (1998) 323
- [6] S. V. Mikhailov, A. V. Radyushkin, *Sov. J. Nucl. Phys.* **49** (1989) 494; *Phys. Rev.* **D45** (1992) 1754;  
A. P. Bakulev, A. V. Radyushkin, *Phys. Lett.* **B271** (1991) 223
- [7] A. P. Bakulev, S. V. Mikhailov, *Phys. Lett.* **B436** (1998) 351
- [8] A. V. Radyushkin, hep-ph/9707335
- [9] S. V. Mikhailov, A. V. Radyushkin, *JETP Lett.* **43** (1986) 712
- [10] V. M. Braun, I. E. Filyanov, *Z. Phys.* **C44** (1989) 157
- [11] N. G. Stefanis, W. Schroers, H. C. Kim, *Phys. Lett.* **B449** (1999) 299
- [12] A. Schmedding, O. Yakovlev, *ITP Prepr.* WUE-ITP-99-011, Wurzburg (1999), hep-ph/9905392
- [13] A. V. Radyushkin, *CEBAF Prepr.* TH-94-13, Newport News (1994), hep-ph/9406237;  
A. P. Bakulev, S. V. Mikhailov, *Z. Phys.* **C68** (1995) 451; *Mod. Phys. Lett.* **A11** (1996) 1611
- [14] A. V. Radyushkin, R. Ruskov, *Nucl. Phys.* **481** (1996) 625
- [15] A. P. Bakulev, S. V. Mikhailov, *Eur. Phys. J.* **C466** (2000) to be published;  
hep-ph/9908287
- [16] V. M. Belyaev, B. L. Ioffe, *ZhETF* **83** (1982) 876;  
A. A. Ovchinnikov, A. A. Pivovarov, *Yad. Fiz.* **48** (1988) 1135;  
A. A. Pivovarov, *Kratk. Soobshch. Fiz. (Bull. Lebedev Phys. Inst.)* **N<sup>o</sup>5** (1991) 3
- [17] M. V. Polyakov, C. Weiss, *Phys. Lett.* **B387** (1996) 841;  
A. E. Dorokhov, S. V. Esaybegyan, S. V. Mikhailov, *Phys. Rev.* **D56** (1997) 4062
- [18] M. D'Elia, A. Di Giacomo, E. Meggiolaro, *Phys. Rev.* **D59** (1999) 054503;  
E. Meggiolaro, hep-lat/9909068.
- [19] H. G. Dosch, M. Eidemüller, M. Jamin, E. Meggiolaro, hep-ph/0004040.
- [20] S. V. Mikhailov, *Sov. J. Nucl. Phys.* **56** (1993) 650
- [21] Particle Data Group Booklet, July 1998.
- [22] D. Becirevic et al., hep-lat/9808187; hep-lat/9808129.
- [23] M. V. Polyakov, *Nucl. Phys.* **B555** (1999) 231
- [24] Patricia Ball, V. M. Braun, *Phys. Rev.* **D55** (1997) 5561
- [25] Patricia Ball, V. M. Braun, *Phys. Rev.* **D58** (1998) 094016
- [26] M. A. Shifman, A. I. Vainshtein, V. I. Zakharov, *Nucl. Phys.* **B147** (1979) 385, 448.
- [27] V. M. Belyaev, A. Oganesyan, *Phys. Lett.* **B395** (1997) 307

Received by Publishing Department  
on July 28, 2000.

**The Publishing Department  
of the Joint Institute for Nuclear Research  
offers you to acquire the following books:**

Index	Title
E1,2-97-79	Proceedings of the XIII International Seminar on High Energy Physics Problems. Relativistic Nuclear Physics and Quantum Chromodynamics. Dubna, 1996 (2 volumes, 364 p. and 370 p., in English)
D5,11-97-112	Proceedings of the 9th International Conference «Computational Modelling and Computing in Physics». Dubna, 1996 (378 p., in English)
E2-97-213	Proceedings of the V International Seminar on Interaction of Neutron with Nuclei «Neutron Spectroscopy, Nuclear Structure, Related Topics». Dubna, 1997 (446 p., in English)
E2,4-97-263	Proceedings of the Third International Conference «Renormalization Group'96». Dubna, 1996 (436 p., in English)
E10-97-272	Proceedings of the Data Acquisition Systems of Neutron Experimental Facilities (DANEF'97). Dubna, 1997 (325 p., in English)
D19-97-284	Proceedings of the International Symposium «Problems of Biochemistry. Radiation and Space Biology». Dubna, 1997 (2 volumes 284 p. and 405 p., in Russian and English)
E2-97-413	Proceedings of the VII Workshop on High Energy Spin Physics (SPIN'97). Dubna, 1997 (398 p., in English)
E15-98-57	Proceedings of the III International Workshop «Hyperfine Structure and Nuclear Moments of Exotic Nuclei by Laser Spectroscopy». Poznan, 1997 (200 p., in English)
D13-98-66	Proceedings of the XVII International Symposium on Nuclear Electronics. Varna, 1997 (242 p., in English and Russian)
E1,2-98-154	Proceedings of the XIII International Seminar on High Energy Physics Problems «Relativistic Nuclear Physics and Quantum Chromodynamics». Dubna, 1996 (2 volumes 300 p. and 282 p., in English)
E3,14-98-168	Proceedings of the German–Russian User Meeting «Condensed Matter Physics with Neutrons at IBR-2». Dubna, 1998 (126 p., in English)
E3-98-202	Proceedings of the VI International Seminar on Interaction of Neutrons with Nuclei Neutron Spectroscopy, Nuclear Structure, Related Topics. Dubna, 1998 (352 p., in English)
D1,2-98-215	Proceedings of the International Workshop «Relativistic Nuclear Physics: from MeV to TeV». Dubna, 1998 (384 p., in English and Russian)
E2-98-254	Proceedings of the International Workshop Hadronic Atoms and Positronium in the Standard Model. Dubna, 1998 (260 p., in English)

Index	Title
D9,11-98-273	Proceedings of the 4th International Workshop «Beam Dynamics and Optimization». Dubna, 1997 (162 p., in English and Russian)
E17,19-98-305	Proceedings of the International Workshop on Deuteration of Biological Molecules for Structural and Dynamic Studies. Applications to Neutron Scattering and NMR. Dubna, 1998 (100 p., in English)
E1,2-98-307	Proceedings of the International School–Seminar «Actual Problems of Particle Physics». Gomel, 1997, Belarus (2 volumes 304 p. and 220 p., in English)
E2-98-372	Proceedings of the III International Workshop «Classical and Quantum Integrable Systems». Erevan, 1998 (200 p., in English)
E9-99-26	Proceedings of the International Conference on High Energy Accelerators. Dubna, 1998 (432 p., in English)
E2-99-35	Proceedings of the XI International Conference on Problems of Quantum Field Theory. Dubna, 1998 (508 p., in English)
E5-99-38	Proceedings of the International Workshop «Self-Similar Systems». Dubna, 1998 (404 p., in English)
E9-99-92	Proceedings of the XVII International Conference on High Energy Accelerators HEACC'98. Dubna, 1998 (435 p., in English)

Please apply to the Publishing Department of the Joint Institute for Nuclear Research for extra information. Our address is:

Publishing Department  
 Joint Institute for Nuclear Research  
 Dubna, Moscow Region  
 141980 Russia  
 E-mail: [publish@pds.jinr.dubna.su](mailto:publish@pds.jinr.dubna.su).

Бакулев А.П., Михайлов С.В.

E2-2000-176

Новые формы амплитуд распределений на световом конусе для поперечно-поляризованных  $\rho$ -мезонов

В рамках метода правил сумм КХД с нелокальными конденсатами заново анализируются амплитуды распределений ведущего твиста поперечно-поляризованных  $\rho$ -,  $\rho'$ - и  $b_1$ -мезонов. Для получения обоснованных оценок используются различные типы правил сумм; в результате получены предсказания для 2-, 4-, 6- и 8-го моментов амплитуд распределений поперечно-поляризованных  $\rho$ - и  $\rho'$ -мезонов и для тензорных распадных констант  $f_{\rho, \rho', b_1}^T$ . Подчеркивается, что результаты обработки стандартных правил сумм

КХД согласуются с полученной оценкой второго момента поперечно-поляризованного  $\rho$ -мезонного распределения. Построены новые модели амплитуд распределений на световом конусе для поперечно-поляризованных  $\rho$ - и  $\rho'$ -мезонов. Кратко обсуждаются феноменологические следствия предложенных моделей. Полученные результаты сравниваются с результатами Баль-Брауна (1996), в результате выявлена неполнота последних.

Работа выполнена в Лаборатории теоретической физики им. Н.Н.Боголюбова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2000

Bakulev A.P., Mikhailov S.V.

E2-2000-176

New Shapes of Light-Cone Distributions of the Transversely Polarized  $\rho$ -Mesons

The leading twist light-cone distributions for transversely polarized  $\rho$ -,  $\rho'$ - and  $b_1$ -mesons are reanalyzed in the framework of QCD sum rules with nonlocal condensates. Using different kinds of sum rules to obtain reliable predictions, we estimate the 2-, 4-, 6- and 8-th moments for transversely polarized  $\rho$ - and  $\rho'$ -meson distributions and reestimate tensor couplings  $f_{\rho, \rho', b_1}^T$ . We stress that the results of standard sum rules also

support our estimation of the second moment of the transversely polarized  $\rho$ -meson distribution. New models for light-cone distributions of these mesons are constructed. Phenomenological consequences from these distributions are briefly discussed. Our results are compared with those found by Ball and Braun (1996), and the latter is shown to be incomplete.

The investigation has been performed at the Bogoliubov Laboratory of Theoretical Physics, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 2000

Макет Т.Е.Попеко

Подписано в печать 11.09.2000  
Формат 60 × 90/16. Офсетная печать. Уч.-изд. листов 1,61  
Тираж 425. Заказ 52225. Цена 1 р. 94 к.

Издательский отдел Объединенного института ядерных исследований  
Дубна Московской области