



LABORATORI NAZIONALI DI FRASCATI
SIS-Pubblicazioni

LNF-00/002 (P)

11 Gennaio 2000

hep-ph/9911422

LEPTOGENESIS FROM R PARITY NON-CONSERVATION

Thomas Hambye¹, Ernest Ma², and Utpal Sarkar³

¹ *INFN, Laboratori Nazionali di Frascati, P.O. Box 13, I-00044 Frascati, Italy*

² *Department of Physics, University of California, Riverside, California 92521, USA*

³ *Physical Research Laboratory, Ahmedabad 380 009, India*

Abstract

It is known that realistic neutrino masses for neutrino oscillations may be obtained from R parity nonconserving supersymmetry. It is also known that such interactions would erase any preexisting lepton or baryon asymmetry of the Universe because of the inevitable intervention of the electroweak sphalerons. We now show how a crucial subset of these R parity nonconserving terms may in fact create its own successful leptogenesis.

PACS:12.60.Jv, 11.30.Fs, 98.80.Cq, 14.80.Ly

Submitted to Phys. Rev. Lett.

To obtain naturally small Majorana neutrino masses[1], the canonical approach is to have heavy singlet right-handed neutrinos in the guise of the famous seesaw mechanism[2]. An equally simple alternative is to use a heavy scalar Higgs triplet[3]. Both offer the important additional benefit of generating a lepton asymmetry[3,4] which gets converted into the present observed baryon asymmetry of the Universe by virtue of the electroweak sphalerons[5,6].

In the past several years, a third way has come under active consideration, namely that of R parity nonconserving supersymmetry. Whereas realistic neutrino masses for neutrino oscillations may be generated[7], the lepton-number nonconserving interactions involved would certainly erase any preexisting lepton asymmetry of the Universe[8]. This negative aspect of R parity nonconserving supersymmetry has prevented it from being considered as a truly competitive alternative to the canonical seesaw or the triplet Higgs mechanism for neutrino masses. To remedy this situation, we now show for the first time how a crucial subset of these R parity nonconserving terms may in fact create its own successful leptogenesis, through the suppressed decay of the lightest neutralino into a charged Higgs boson and a lepton.

In a supersymmetric extension of the minimal standard model of fundamental interactions, the R parity of a particle is conventionally defined as

$$R \equiv (-1)^{3B+L+2J}, \quad (1)$$

where B is its baryon number, L its lepton number, and J its spin angular momentum. Hence the standard-model particles have $R = +1$ and their supersymmetric partners have $R = -1$. Using the common notation where all chiral superfields are considered left-handed, the three families of leptons and quarks are given by

$$L_i = (\nu_i, e_i) \sim (1, 2, -1/2), \quad e_i^c \sim (1, 1, 1), \quad (2)$$

$$Q_i = (u_i, d_i) \sim (3, 2, 1/6), \quad (3)$$

$$u_i^c \sim (3^*, 1, -2/3), \quad d_i^c \sim (3^*, 1, 1/3), \quad (4)$$

where i is the family index, and the two Higgs doublets are given by

$$H_1 = (h_1^0, h_1^-) \sim (1, 2, -1/2), \quad (5)$$

$$H_2 = (h_2^+, h_2^0) \sim (1, 2, 1/2), \quad (6)$$

where the $SU(3)_C \times SU(2)_L \times U(1)_Y$ content of each superfield is also indicated. If R parity is conserved, the superpotential is restricted to have only the terms

$$W = \mu H_1 H_2 + f_{ij}^e H_1 L_i e_j^c + f_{ij}^d H_1 Q_i d_j^c + f_{ij}^u H_2 Q_i u_j^c. \quad (7)$$

If R parity is violated but not baryon number, then the superpotential contains the additional terms

$$W' = \mu_i L_i H_2 + \lambda_{ijk} L_i L_j e_k^c + \lambda'_{ijk} L_i Q_j d_k^c, \quad (8)$$

resulting in nonzero neutrino masses either from mixing with the neutralino mass matrix or in one-loop order[7].

Consider the simplified case of having only L_τ and τ^c as effective $L = 0$ superfields[9,10]. The neutralino mass matrix now spans five fields: the U(1) gaugino (\tilde{B}), the SU(2) gaugino (\tilde{W}_3), the two Higgsinos ($\tilde{h}_1^0, \tilde{h}_2^0$), and ν_τ .

$$\mathcal{M} = \begin{bmatrix} M_1 & 0 & -sm_3 & sm_4 & -sm_5 \\ 0 & M_2 & cm_3 & -cm_4 & cm_5 \\ -sm_3 & cm_3 & 0 & -\mu & 0 \\ sm_4 & -cm_4 & -\mu & 0 & -\mu_\tau \\ -sm_5 & cm_5 & 0 & -\mu_\tau & 0 \end{bmatrix}, \quad (9)$$

where $s = \sin \theta_W$, $c = \cos \theta_W$, $m_3 = M_Z \cos \beta \cos \theta_\tau$, $m_4 = M_Z \sin \beta$, $m_5 = M_Z \cos \beta \sin \theta_\tau$, with $\tan \beta = \langle h_2^0 \rangle / [\langle h_1^0 \rangle^2 + \langle \tilde{\nu}_\tau \rangle^2]^{1/2}$ and $\tan \theta_\tau = \langle \tilde{\nu}_\tau \rangle / \langle h_1^0 \rangle$.

To understand the structure of the above 5×5 mass matrix, let us assume that μ is the dominant term, then $\tilde{h}_{1,2}^0$ form a heavy Dirac particle of mass μ and the reduced 3×3 matrix in the basis ($\tilde{B}, \tilde{W}_3, \nu_\tau$) becomes

$$\mathcal{M} = \begin{bmatrix} M_1 - s^2 \delta & sc\delta & -s\epsilon \\ sc\delta & M_2 - c^2 \delta & c\epsilon \\ -s\epsilon & c\epsilon & 0 \end{bmatrix}, \quad (10)$$

where

$$\delta = 2m_3 m_4 / \mu = M_Z^2 \sin 2\beta \cos \theta_\tau / \mu, \quad (11)$$

$$\begin{aligned} \epsilon &= m_5 - m_3 \mu_\tau / \mu \\ &= M_Z \cos \beta \cos \theta_\tau (\tan \theta_\tau - \mu_\tau / \mu). \end{aligned} \quad (12)$$

From the above, ν_τ gets a seesaw mass, i.e.

$$m_{\nu_\tau} = -\epsilon^2 \left(\frac{s^2}{M_1} + \frac{c^2}{M_2} \right), \quad (13)$$

corresponding to the mass eigenstate

$$\nu'_\tau = \nu_\tau + \frac{s\epsilon}{M_1} \tilde{B} - \frac{c\epsilon}{M_2} \tilde{W}_3, \quad (14)$$

whereas the other two mass eigenstates are

$$\tilde{B}' = \tilde{B} + \frac{sc\delta}{M_1 - M_2} \tilde{W}_3 - \frac{s\epsilon}{M_1} \nu_\tau, \quad (15)$$

$$\tilde{W}'_3 = \tilde{W}_3 - \frac{sc\delta}{M_1 - M_2} \tilde{B} + \frac{c\epsilon}{M_2} \nu_\tau. \quad (16)$$

Because \tilde{B}' and \tilde{W}'_3 contain ν_τ , they are not stable, but would decay into $\tau_\pi^m W^\pm$ pairs. This may generate a lepton asymmetry, but it is several orders of magnitude too small because it has to be much less than $(\epsilon/M_{1,2})^2$, which is of order $m_{\nu_\tau}/M_{1,2}$, i.e. $< 5 \times 10^{-13}$ if $m_{\nu_\tau} < 0.05$ eV and $M_{1,2} > 100$ GeV.

Consider now the couplings of \tilde{B} and \tilde{W}_3 . The former couples to both $\tilde{\tau}_L \tilde{\tau}_L$ and $\tilde{\tau}_L^c \tilde{\tau}_L^c$, but the latter only to $\tilde{\tau}_L \tilde{\tau}_L$, because τ_L^c is a singlet under $SU(2)_L$. Since R parity is violated, both $\tilde{\tau}_L$ and $\tilde{\tau}_L^c$ mix with the charged Higgs boson of the supersymmetric standard model: $h^\pm = h_2^\pm \cos \beta + h_1^\pm \sin \beta$. Hence \tilde{B}' and \tilde{W}'_3 may decay into $\tau_\pi^m h^\pm$. We now assume that the $\tilde{\tau}_L$ mixing is negligible, so that the only relevant coupling is that of \tilde{B} to $\tilde{\tau}_L^c h^+$. Hence \tilde{W}'_3 decay (into $\tau_\pi^m h^\pm$) is suppressed because it may only do so through the small component of \tilde{B} that it contains, assuming of course that $\tilde{\tau}_L$ and $\tilde{\tau}_L^c$ are heavier than \tilde{B} or \tilde{W}_3 .

We now envisage the following leptogenesis scenario. At temperatures well above $T = M_{SUSY}$, the presence of sphalerons and R parity nonconserving interactions together ensure that there is no L or B asymmetry. Note that W' of Eq. (8) violates L_i as well as $B - 3L_i$ for $i = e, \mu, \tau$ [8]. As the Universe cools down to a temperature below the masses of all supersymmetric particles except \tilde{B}' and \tilde{W}'_3 , we need only consider their interactions. In Figure 1 we show the lepton number violating processes (a) $\tilde{B}' \leftrightarrow \tau_R^\pm h_\pi^m$, where we have adopted the more conventional notation of an outgoing τ_R in place of an incoming τ_L^c . These are still certainly fast and there can be no L asymmetry. Let us assume now that $M_1 > M_2$, then for $T < M_1$, the \tilde{B}' interactions are suppressed and we need only consider \tilde{W}'_3 . However, as we have explained in the previous paragraph, the lepton-number violating processes (b) $\tilde{W}'_3 \leftrightarrow \tau_R^\pm h_\pi^m$ are slow and can satisfy the out-of-equilibrium condition for generating a lepton asymmetry of the Universe, resulting from the interference of this tree-level diagram with the one-loop (c) self-energy and (d) vertex diagrams. Since the unsuppressed lepton number violating couplings of \tilde{B}' are involved, a realistic lepton asymmetry may be generated. It is then converted by the still active sphalerons into the present observed baryon asymmetry of the Universe.

We start with the well-known interaction of \tilde{B} with τ and $\tilde{\tau}_R$ given by[11]

$$-\frac{e\sqrt{2}}{\cos \theta_W} \left[\tilde{\tau} \left(\frac{1 - \gamma_5}{2} \right) \tilde{B} \tilde{\tau}_R + H.c. \right]. \quad (17)$$

We then allow $\tilde{\tau}_R$ to mix with h^- , and \tilde{B} to mix with \tilde{W}_3 , so that the interaction of the physical state \tilde{W}'_3 of Eq. (16) with τ and h^\pm is given by

$$\left(\frac{sc\xi}{M_1 - M_2} \right) \left(\frac{e\sqrt{2}}{\cos \theta_W} \right) \left[\delta\tilde{\tau} \left(\frac{1 - \gamma_5}{2} \right) \tilde{W}'_3 h^- + H.c. \right], \quad (18)$$

where ξ represents the $\tilde{\tau}_R - h^-$ mixing and is assumed real, but the parameter δ of Eq. (10) is now assumed complex. The origin of this nontrivial CP phase is from the 2×2 Majorana mass matrix spanning \tilde{B} and \tilde{W}_3 , with complex M_1 and M_2 . It is independent of the phase of μ and contributes negligibly to the neutron electric dipole moment because the magnitude of δ is very small. [Note that the usual assumption of CP violation in supersymmetric models is that M_1 and M_2 have a common phase, in which case the phase of δ would be equal to the phase of μ .]

To satisfy the out-of-equilibrium condition, we require[12]

$$\Gamma(\tilde{W}'_3 \rightarrow \tau^\pm h_\pi^m) < H = 1.7\sqrt{g_*}(T^2/M_{Pl}) \quad (19)$$

at the temperature $T \sim M_2$, where H is the Hubble expansion rate of the Universe with g_* the effective number of massless degrees of freedom and M_{Pl} the Planck mass. This implies

$$\left(\frac{\xi|\delta|r}{M_1 - M_2}\right)^2 \frac{(M_2^2 - m_h^2)^2}{M_2^5} < 1.9 \times 10^{-14} \text{ GeV}^{-1}, \quad (20)$$

where we have used $g_* = 10^2$ and $M_{Pl} = 10^{18}$ GeV, and $r = (1 + M_2/\mu \sin 2\beta)/(1 - M_2^2/\mu^2)$ is a correction factor for finite M_2/μ . To make sure that at $T \sim M_2$, the lepton number violating processes $\tau^\pm h_\pi^m \leftrightarrow \tau_\pi^m h^\pm$ through \tilde{B}' exchange do not erase the lepton asymmetry created by the decay of \tilde{W}'_3 , we require

$$\left(\frac{2e^2\xi^2}{\cos^2\theta_W}\right)^2 \frac{1}{M_1^2} \frac{T^3}{32\pi} \frac{f(x)}{(1-x)^2} < H \quad (21)$$

at $T \sim M_2$, where $x = M_2^2/M_1^2$ and

$$f(x) = 1 + \frac{2(1-x)}{x^2} [(1+3x)\ln(1+x) - x(1+x)], \quad (22)$$

which implies

$$\frac{\xi^4}{M_2} \frac{xf(x)}{(1-x)^2} < 2.6 \times 10^{-14} \text{ GeV}^{-1}. \quad (23)$$

Both conditions may be satisfied for example with the following choice of parameters:

$$\mu = 5 \text{ TeV}, \quad \sin 2\beta = 0.5, \quad \xi = 2 \times 10^{-3}, \quad (24)$$

$$M_1 = 3 \text{ TeV}, \quad M_2 = 2 \text{ TeV}, \quad m_h = 200 \text{ GeV}, \quad (25)$$

for which the left-hand sides of Eqs. (20) and (23) are $0.6 \times 10^{-14} \text{ GeV}^{-1}$ and $2.6 \times 10^{-14} \text{ GeV}^{-1}$ respectively. Hence the usual parameter K defined as the ratio of the left-hand side of Eq. (19) to the right-hand side at $T = M_2$ is 0.3 in this case. From a numerical calculation of the Boltzmann equation, taking into account the inverse decay

contribution, we find[13] the asymptotic suppression factor to be about 2/3, which is reached at a temperature of about 0.4 TeV.

The interference between the tree-level and self-energy + vertex diagrams results in the following lepton asymmetry in the decay of \tilde{W}'_3 :

$$\epsilon = \frac{\alpha\xi^2}{3\cos^2\theta_W} \frac{Im\delta^2}{|\delta|^2} \left(1 - \frac{m_h^2}{M_2^2}\right)^2 \frac{x^{1/2}g(x)}{(1-x)}, \quad (26)$$

where the factor 2/3 has been inserted and

$$g(x) = 1 + \frac{2(1-x)}{x} \left[\left(\frac{1+x}{x}\right) \ln(1+x) - 1 \right], \quad (27)$$

which yields 2.4×10^{-8} if $Im\delta^2/|\delta|^2 = 1$. Dividing by $3g_*$, we then obtain a realistic baryon-to-photon ratio of the order 10^{-10} .

In the above, we have assumed $M_{SUSY} \sim \text{few TeV}$. This value is naturally suited for our scenario because (1) it allows ϵ in Eq. (26) to be large enough without contradicting the condition given by Eq. (21), and (2) it allows the lepton asymmetry to be converted at a temperature when the sphalerons are still very active[14,15], into the present observed baryon asymmetry of the Universe. We have also assumed in our scenario that \tilde{W}'_3 only couples to $\tau^\pm h_\pi^m$ through its \tilde{B} content. This means that $\tilde{\tau}_R - h^-$ mixing is significant, but $\tilde{\tau}_L - h^-$ mixing is negligible. The latter is related to the mixing of the neutrino with the neutralinos as given by Eq. (9) which is indeed small and may even be set to zero, because realistic neutrino masses may be obtained through R parity nonconserving trilinear couplings instead[7]. The former is usually taken[16] from the soft supersymmetry breaking term $h_1^- \tilde{\nu}_\tau \tilde{\tau}_L^c$ and the $\tilde{\tau}_L h_2^+$ term through $\tilde{\tau}_L^* - \tilde{\tau}_L^c$ mixing. In that case, the $\tilde{\tau}_R - h^-$ mixing would be proportional to a linear combination of $\langle \tilde{\nu}_\tau \rangle$ and μ_τ , and be constrained to be very small[13]. For our scenario to be successful, we need to introduce the new term

$$H_2^\dagger H_1 \tilde{\tau}_L^c = (\bar{h}_2^+ h_1^0 + \bar{h}_2^0 h_1^-) \tilde{\tau}_L^c, \quad (28)$$

which is unconstrained and may have the desired mixing of $\xi = 2 \times 10^{-3}$ shown. Such a soft supersymmetry breaking term is nonholomorphic[17] and whereas this class of couplings has been studied previously[18], we are the first to make use of this particular term. We should also mention that although we have used only τ to represent a lepton, replacing it with a linear combination of e , μ , and τ is just as appropriate.

In conclusion, we have shown in this paper how R parity nonconservation may create its own successful leptogenesis through the suppressed decay of \tilde{W}'_3 into $l^\pm h_\pi^m$. We require $M_{SUSY} \sim \text{few TeV}$ and the existence of the nonholomorphic soft supersymmetry breaking term $H_2^\dagger H_1 \tilde{l}_L^c$. Whereas this mechanism depends on R parity nonconservation, it

comes from a sector distinct from that responsible for nonzero neutrino masses[19]. We also require the soft supersymmetry breaking gaugino masses M_1 and M_2 to be a few TeV with $M_1 > M_2$.

Acknowledgements

E.M. thanks S. Pakvasa and X. Tata for discussions, and the support of the U.S. Department of Energy under Grant No. DE-FG03-94ER40837.

References

- [1] For a recent overview, see for example E. Ma, Phys. Rev. Lett. **81**, 1171 (1998).
- [2] M. Gell-Mann, P. Ramond and R. Slansky, in *Supergravity*, edited by P. van Nieuwenhuizen and D. Freedman, (North-Holland, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe*, edited by O. Sawada and A. Sugamoto (KEK Report No. 79-18, Tsukuba, 1979), p. 95; R. N. Mohapatra and G. Senjanović, Phys. Rev. Lett. **44**, 912 (1980).
- [3] E. Ma and U. Sarkar, Phys. Rev. Lett. **80**, 5716 (1998).
- [4] M. Fukugita and T. Yanagida, Phys. Lett. **174B**, 45 (1986).
- [5] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. **155B**, 36 (1985).
- [6] M. Fukugita and T. Yanagida, Phys. Rev. **D42**, 1285 (1990); J. A. Harvey and M. S. Turner, Phys. Rev. **D42**, 3344 (1990); A. E. Nelson and S. M. Barr, Phys. Lett. **B246**, 141 (1990); W. Buchmüller and T. Yanagida, Phys. Lett. **B302**, 240 (1993).
- [7] R. Hempfling, Nucl. Phys. **B478**, 3 (1996); F. M. Borzumati, Y. Grossman, E. Nardi, and Y. Nir, Phys. Lett. **B384**, 123 (1996); E. Nardi, Phys. Rev. **D55**, 5772 (1997); M. A. Diaz, J. C. Romao, and J. W. F. Valle, Nucl. Phys. **B524**, 23 (1998); M. Drees, S. Pakvasa, X. Tata, and T. ter Veldhuis, Phys. Rev. **D57**, R5335 (1998); B. Mukhopadhyaya, S. Roy, and F. Vissani, Phys. Lett. **B443**, 191 (1998); E. J. Chun, S. K. Kang, and C. W. Kim, Nucl. Phys. **B544**, 89 (1999); R. Adhikari and G. Omanovic, Phys. Rev. **D59**, 073003 (1999); O. C. W. Kong, Mod. Phys. Lett. **A14**, 903 (1999); S. Rakshit, G. Bhattacharyya, and A. Raychaudhuri, Phys. Rev. **D59**, 091701 (1999); K. Choi, E. J. Chun, and K. Hwang, Phys. Rev. **D60**, 031301 (1999); S. Y. Choi, E. J. Chun, S. K. Kang, and J. S. Lee, Phys. Rev. **D60**, 075002 (1999); L. Clavelli and P. H. Frampton, hep-ph/9811326; D. E. Kaplan and A. E. Nelson, J. High Energy

- Phys. **0001**, 033 (2000); A. S. Joshipura and S. Vempati, Phys. Rev. **D60**, 111303 (1999); Y. Grossman and H. E. Haber, hep-ph/9906310; G. Bhattacharyya, H. V. Klapdor-Kleingrothaus, and H. Pas, Phys. Lett. **B463**, 77 (1999); A. Abada and M. Losada, hep-ph/9908352; O. Haug, J. D. Vergados, A. Faessler, and S. Kovalenko, hep-ph/9909318; E. J. Chun and S. K. Kang, hep-ph/9909429; F. Takayama and M. Yamaguchi, hep-ph/9910320.
- [8] B. A. Campbell, S. Davidson, J. E. Ellis, and K. Olive, Phys. Lett. **B256**, 457 (1991); W. Fischler, G. F. Giudice, R. G. Leigh, and S. Paban, Phys. Lett. **B258**, 45 (1991); H. Dreiner and G. G. Ross, Nucl. Phys. **B410**, 188 (1993); J. M. Cline, K. Kainulainen, and K. A. Olive, Phys. Rev. **D49**, 6394 (1994); E. Ma, M. Raidal, and U. Sarkar, Phys. Lett. **B460**, 359 (1999).
- [9] If only ν_τ and τ mix with the neutralinos, then a consistent view is to assign $L = 0$ to all such fields, but keep $L = 1$ for e and μ , as discussed in E. Ma and P. Roy, Phys. Rev. **D41**, 988 (1990).
- [10] C.-H. Chang and T.-F. Feng, Eur. Phys. J. **C12**, 137 (2000); see also M. A. Diaz, J. C. Romao, and J. W. F. Valle, Ref.[7].
- [11] H. E. Haber and G. L. Kane, Phys. Rep. **117**, 75 (1985).
- [12] E. W. Kolb and M. S. Turner, *The Early Universe* (Addison-Wesley, Reading, MA, 1990).
- [13] This and other details will be discussed elsewhere.
- [14] See for example A. Pilaftsis, Phys. Rev. **D56**, 543 (1997); Int. J. Mod. Phys. **A14**, 1811 (1999).
- [15] Note also that we do not require the electroweak phase transition to be first-order for satisfying the out-of-equilibrium condition. See for example A. Riotto, hep-ph/9807454; A. Riotto and M. Trodden, hep-ph/9901362.
- [16] A. Akeroyd, M. A. Diaz, J. Ferrandis, M. A. Garcia-Jareno, and J. W. F. Valle, Nucl. Phys. **B529**, 3 (1998); C.-H. Chang and T.-F. Feng, hep-ph/9908295.
- [17] L. J. Hall and L. Randall, Phys. Rev. Lett. **65**, 2939 (1990).
- [18] F. Borzumati, G. R. Farrar, N. Polonsky, and S. Thomas, Nucl. Phys. **B555**, 53 (1999).

[19] This is not unlike the case of neutrinoless double beta decay, where the dominant contribution may not be directly related to nonzero Majorana neutrino masses.