




the
abdus salam
international
centre
for theoretical
physics



XA0203945



**THE DAS-POPOWICZ MOYAL MOMENTUM
ALGEBRA**

A. Boulahoual

and

M.B. Sedra

preprint

34 / 04

United Nations Educational Scientific and Cultural Organization
and
International Atomic Energy Agency
THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

THE DAS-POPOWICZ MOYAL MOMENTUM ALGEBRA

A. Boulahoual and M.B. Sedra¹
*Laboratoire de Physique Théorique et Appliquée LPTA,
Université Ibn Tofail, Faculté des Sciences, Département de Physique,
B.P. 133, Kenitra, Morocco*
and
The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

Abstract

We introduce in this short note some aspects of the Moyal momentum algebra that we call the Das-Popowicz Mm algebra. Our interest on this algebra is motivated by the central role that it can play in the formulation of integrable models and in higher conformal spin theories.

MIRAMARE – TRIESTE

August 2002

¹Regular Associate of the Abdus Salam ICTP.

*This work is dedicated to the memory of
my father who recently passed away...*

M.B.S.

1 Introduction

In the past few years, there has been a growth in the interest in non-commutative geometry (NCG), which appears in string theory in several ways [1]. Much attention has been paid also to field theories on NC spaces and more specifically Moyal deformed space-time, because of the appearance of such theories as certain limits of string, D-brane and M-theory [2]. One of the strong points of the NCG framework is its richness and also the fact that we can recover all the well known standard results just by requiring the vanishing of the deformed parameter which means also the vanishing of noncomutativity. Note that the passage from commutative to noncommutative space time is achieved by replacing the ordinary commutative product, in the space of smooth functions on R^2 with coordinates x, t , by the noncommutative associative \star - product [3]. Moyal deformation is also applied to Lax equations and supersymmetric KdV hierarchies [4,5].

The aim of this letter is to introduce some aspects of the Moyal momentum algebra that we call the Das-Popowicz Mm algebra [4] shown to play, in the NCG framework, an important role in formulating integrable models and higher conformal spin theories in a systematic way [6].

2 The Das-Popowicz Mm-algebra

Using our convention notations [7], we denote this algebra by $\widehat{\Sigma}(\theta)$. This is a non-commutative space based on arbitrary momentum Lax operators and which decomposes as:

$$\widehat{\Sigma}(\theta) = \bigoplus_{r \leq s} \bigoplus_{m \in Z} \widehat{\Sigma}_m^{(r,s)}, \quad (1)$$

where $\widehat{\Sigma}_m^{(r,s)}(\theta)$ is the space of θ -Lax operators of fixed conformal spin m and degrees (r,s) given by

$$\mathcal{L}_m^{(r,s)}(u) = \sum_{i=r}^s u_{m-i}(x) \star p^i. \quad (2)$$

These are θ -differentials whose operator character is inherited from the star product law defined as follows:

$$f(x, p) \star g(x, p) = \sum_{s=0}^{\infty} \sum_{i=0}^s \frac{\theta^s}{s!} (-)^i c_s^i (\partial_x^i \partial_p^{s-i} f) (\partial_x^{s-i} \partial_p^i g), \quad (3)$$

with $c_s^i = \frac{s!}{i!(s-i)!}$ and $f(x, p)$ are arbitrary functions on the two-dimensional phase space.

Now, it is important to precise how the momentum operators act on arbitrary functions $f(x, p)$ via the star product. Performing computations based on relation (3), we find the following θ -Leibniz rules:

$$p^n \star f(x, p) = \sum_{s=0}^n \theta^s c_n^s f^{(s)}(x, p) p^{n-s}, \quad (4)$$

and

$$p^{-n} \star f(x, p) = \sum_{s=0}^{\infty} (-)^s \theta^s c_{n+s-1}^s f^{(s)}(x, p) p^{-n-s}, \quad (5)$$

with $n \in \mathbb{N}$ and $f^{(s)} = (\partial_x^s f)$ is the prime derivative.

We also find that the Moyal bracket defined as [8]:

$$\{f(x, p), g(x, p)\}_\theta = \frac{f \star g - g \star f}{2\theta} \quad (6)$$

is subject to the following expressions

$$\begin{aligned} \{p^n, f\}_\theta &= \sum_{s=0}^n \theta^{s-1} c_n^s \left\{ \frac{1-(-)^s}{2} \right\} f^s p^{n-s}, \\ \{p^{-n}, f\}_\theta &= \sum_{s=0}^{\infty} \theta^{s-1} c_{s+n-1}^s \left\{ \frac{(-)^s - 1}{2} \right\} f^s p^{-n-s}. \end{aligned} \quad (7)$$

Having defined the Das-Popowicz algebra, we present here below, some remarkable properties.

P1: We give here below the conformal dimensions of objects used in this study

$$[x] = -1, [p] = [\partial_x] = 1, [\theta] = 0, [u_m] = m, [\widehat{Res}] = [Res] = 1$$

P2: The momentum operators p^i satisfy

$$p^n \star p^m = p^{n+m}. \quad (8)$$

ensuring the suspected rule

$$p^n \star (p^{-n} \star f(x, p)) = f(x, p). \quad (9)$$

P3: We should note that formulas (4-5,7) are computed for integers values $n = 0, 1, 2, 3, \dots$. Now, what happens if half integers $n = \pm\frac{1}{2}, \pm\frac{3}{2}, \pm\frac{5}{2}, \dots$ or arbitrary fractional powers were allowed? Using the general formula (3) and performing algebraic computations we find the following formulas:

$$p^{\frac{a}{2}} \star f(x, p) = \sum_{s=0}^{\infty} \Pi_{j=0}^{s-1} \left(\frac{a}{2} - j \right) \frac{\theta^s}{s!} f^{(s)}(x, p) p^{\frac{a}{2}-s}, \quad (10)$$

describing the \star -product action of half integer powers of the momentum operators on the phase space with

$$\Pi_{j=0}^{s-1} \left(\frac{a}{2} - j \right) = \left(\frac{a}{2} \right) \left(\frac{a}{2} - 1 \right) \left(\frac{a}{2} - 2 \right) \dots \left(\frac{a}{2} - s + 1 \right). \quad (11)$$

Moreover, from eq.(10) one can recover eqs.(4-5) once even values $a = 2n$, (for n positive or negative) are considered. We can then conclude that these equations, as well as eq.(8), are special cases of (10). The more general situation consists in considering fractional objects type $p^{\frac{a}{b}}$, we have

$$p^{\frac{a}{b}} \star f(x, p) = \sum_{s=0}^{\infty} \Pi_{j=0}^{s-1} \left(\frac{a}{b} - j \right) \frac{\theta^s}{s!} f^{(s)}(x, p) p^{\frac{a}{b}-s}, \quad (12)$$

Few remarks are in order. If eqs.(4-5) are associated to 'bosonic' behavior of the Das-Popowicz algebra $\widehat{\Sigma}$, the half conformal spin object $p^{\frac{1}{2}} \star u_0 \in \widehat{\Sigma}_{\frac{1}{2}}^{\left(\frac{1}{2}, \frac{1}{2}\right)}$, for example, may lead us

to interpret naively this object as being of fermionic character. The same judgment is made for the fractional object $p^{\frac{a}{b}} \star u_0 \in \widehat{\Sigma}_{\frac{a}{b}}^{(\frac{a}{b}, \frac{a}{b})}$ having fractional conformal spin.

The obviousness of this correspondence with statistics is only an impression. In fact, for example, for objects like $p^{\frac{1}{2}} \star u_0$, even though they present half conformal spins, they are not fermionic objects because $p^{\frac{1}{2}} \star p^{\frac{1}{2}} = p$ and a fermionic object should be nilpotent. In order to attribute a fermionic behavior to our momentum operators, one has to consider a graded phase space parametrized by the set of variables $(x, p, \eta$ and $p_\eta)$ where η is the Grassmann variable and p_η the corresponding nilpotent conjugate momenta given by $p_\eta^2 = 0$ [5].

Nevertheless, it would be very interesting to look for the contribution and the meaning of these fractional powers of momenta in the framework of KdV hierarchy equations and also in the building of the GD second Hamiltonian structure.

P4: The space $\widehat{\Sigma}(\theta)$ may decompose into the underlying sub-algebras as

$$\begin{aligned}\widehat{\Sigma}(\theta) &= \bigoplus_{r \leq s} \widehat{\Sigma}^{(r,s)} \\ &= \bigoplus_{r \leq s} \bigoplus_{m \in Z} \widehat{\Sigma}_m^{(r,s)}(\theta) \\ &= \bigoplus_{r \leq s} \bigoplus_{m \in Z} \bigoplus_{k=r}^s \widehat{\Sigma}_m^{(k,k)}(\theta)\end{aligned}\tag{13}$$

where $\widehat{\Sigma}_m^{(k,k)}(\theta)$ is generated by elements of type $u_{m-k} \star p^k$ or $p^k \star u_{m-k}$.

P5: Using the θ -Leibniz rule, we can write, for fixed value of k :

$$\widehat{\Sigma}_m^{(k,k)} \equiv \Sigma_m^{(k,k)} \oplus \theta \Sigma_m^{(k-1,k-1)} \oplus \theta^2 \Sigma_m^{(k-2,k-2)} \oplus \dots\tag{14}$$

where $\Sigma_m^{(k,k)}$ is the standard one dimensional sub-space, containing the prototype objects $u_{m-k} p^k$ that we consider as the $(\theta = 0)$ -limit of $\widehat{\Sigma}_m^{(k,k)}$.

P6: The space $\widehat{\Sigma}_m^{(0,0)} \equiv \Sigma_m^{(0,0)}$ is nothing but the ring of analytic fields u_m of conformal spin $m \in Z$. With respect to this definition, the subspace $\widehat{\Sigma}_m^{(k,k)}$ can be written formally as:

$$\widehat{\Sigma}_m^{(k,k)} \equiv p^k \star \Sigma_{m-k}^{(0,0)}.\tag{15}$$

P7: We can easily check that, in general, the space $\widehat{\Sigma}_m^{(k,k)}$ is not closed under the action of Moyal bracket (6) since we have:

$$\{.,.\}_\theta : \widehat{\Sigma}_m^{(r,s)} \star \widehat{\Sigma}_m^{(r,s)} \rightarrow \widehat{\Sigma}_{2m}^{(r,2s-1)}\tag{16}$$

Imposing the closure, one gets strong constraints on the integers m , r and s namely

$$m = 0, r \leq s \leq 1\tag{17}$$

Under these constraint equations, the sub-spaces $\widehat{\Sigma}_m^{(r,s)}$ exhibit then a Lie algebra structure since the \star -product is associative.

P8: The sub-space $\widehat{\Sigma}_m^{(r,s)}$ is characterized by the existence of a residue operation that we denote as \widehat{Res} and which acts as follows

$$\begin{aligned}\widehat{Res}(u_k \star p^{-k}) &= (u_k \star p^{-k}) \delta_{k-1,0} \\ &= u_1 \delta_{k-1,0}\end{aligned}\tag{18}$$

This result coincides with the standard residue: Res , acting on the sub-space $\Sigma_m^{(r,s)}$:

$$Res(u_1.p^{-1}) = u_1 \quad (19)$$

We thus have two types of residues \widehat{Res} and Res acting on two different spaces $\widehat{\Sigma}_m^{(r,s)}$ and $\Sigma_m^{(r,s)}$ but with value on the same ring $\Sigma_{m+1}^{(0,0)}$. This property is summarized as follows:

$$\begin{array}{ccc} \widehat{\Sigma}_m^{(r,s)} & \xrightarrow{\theta=0} & \Sigma_m^{(r,s)} \\ \widehat{Res} \searrow & & \swarrow Res \\ & \Sigma_{m+1}^{(0,0)} & \end{array} \quad (20)$$

We learn from this diagram that the residue operation exhibits a conformal spin quantum number equal to 1.

P9: With respect to the previous residue operation, we define on $\widehat{\Sigma}$ the following degrees pairing

$$(\cdot, \cdot) : \widehat{\Sigma}_m^{(r,s)} \star \widehat{\Sigma}_n^{(-s-1, -r-1)} \rightarrow \Sigma_{m+n+1}^{(0,0)} \quad (21)$$

such that

$$(\mathcal{L}_m^{(r,s)}(u), \tilde{\mathcal{L}}_n^{(\alpha,\beta)}(v)) = \delta_{\alpha+s+1,0} \delta_{\beta+r+1,0} Res(\mathcal{L}_m^{(r,s)}(u) \star \tilde{\mathcal{L}}_n^{(\alpha,\beta)}(v)), \quad (22)$$

showing that the spaces $\widehat{\Sigma}_m^{(r,s)}$ and $\widehat{\Sigma}_n^{(-s-1, -r-1)}$ are \widehat{Res} -dual as $\Sigma_m^{(r,s)}$ and $\Sigma_n^{(-s-1, -r-1)}$ are dual with respect to the Res -operation.

Other important properties with possible applications will be considered in a forthcoming work.

Acknowledgements

We would like to thank the Abdus Salam International Centre for Theoretical Physics (ICTP) for hospitality and acknowledge the considerable help of the High Energy Section. We would also like to thank the Associateship office and the Young Collaborator Program. This work was done within the framework of the Associateship Scheme of the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy. We are also grateful to Ashok Das and Ziemowit Popowicz for their interest in this study. A. B. would like to thank the UFR-Faible Radiocativite, Physique Mathematique et Environnement, Departement de Physique, Kenitra.

References

- [1] A. Connes, Noncommutative geometry, Academic Press (1994),
A. Connes, M.R. Douglas, A. Schwarz, Noncommutative geometry and matrix theory: Compactification on tori, JHEP 02(1998) 003, [hep-th/9711162] and references there in.
- [2] N. Seiberg and E. Witten, String theory and non-commutative geometry, JHEP 09(1999) 032.
- [3] M. Kontsevitch, Deformation quantization of Poisson manifolds I, [q-alg/9709040]
D.B. Fairlie, Moyal Brackets, Star Products and the Generalized Wigner Function, [hep-th/9806198]
D.B. Fairlie, Moyal Brackets in M-Theory, Mod.Phys.Lett. A13 (1998) 263-274, [hep-th/9707190],
C. Zachos, A Survey of Star Product Geometry, [hep-th/0008010],
C. Zachos, Geometrical Evaluation of Star Products, J.Math.Phys. 41 (2000) 5129-5134, [hep-th/9912238],
C. Zachos, T. Curtright, Phase-space Quantization of Field Theory, Prog.Theor. Phys. Suppl. 135 (1999) 244-258, [hep-th/9903254].
- [4] A. Das and Z. Popowicz, Properties of Moyal-Lax Representation, Phys.Lett. B510 (2001) 264-270, [hep-th/0103063].
- [5] A. Das and Z. Popowicz, Supersymmetric Moyal-Lax Representations, J. Phys. A, Math. and Gene. 34(2001) 6105 and hep-th/0104191.
- [6] Ming-Hsien Tu, W_n^θ algebra associated with the Moyal KdV hierarchy, Phys.Lett. B508 (2001) 173-183,
Ming-Hsien Tu, Niann-Chern Lee, Yu-Tung Chen, Conformal Covariantization of Moyal-Lax Operators, J.Phys. A35 (2002) 4375.
- [7] E.H.Saidi and M.B.Sedra, On the Gelfand Dickey algebra $GD(sl_n)$ and the W_n -symmetry, J.Math.Phys.35(6), June 1994,
M.B.Sedra, On The huge Lie superalgebra of pseudo superdifferential operators and super KP Hierarchies, J.Math.Phys.37(1996)3483.
- [8] H. Groenewold, Physica 12(1946)405,
J.E. Moyal, Proc. Cambridge Phil. Soc. 45(1949)90.