



the

**abdus salam**  
international  
centre  
for theoretical  
physics



XA0302100

IC/2002/153

**SEPARATION METHOD IN THE PROBLEM OF  
A BEAM-PLASMA INTERACTION IN BOUNDED  
WARM PLASMA UNDER THE EFFECT OF  
HF ELECTRIC FIELD**

**Kh.H. El-Shorbagy**

...34/12

preprint

United Nations Educational Scientific and Cultural Organization  
and  
International Atomic Energy Agency

THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**SEPARATION METHOD IN THE PROBLEM OF A BEAM-PLASMA INTERACTION IN  
BOUNDED WARM PLASMA UNDER THE EFFECT OF HF ELECTRIC FIELD**

Kh.H. El-Shorbagy

*Plasma Physics and Nuclear Fusion Department, Nuclear Research Center,  
Atomic Energy Authority, P.O. Box 13759, Inshass, Cairo, Egypt*

*and*

*The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.*

**Abstract**

The stabilization effect of a strong HF electric field on beam-plasma instability in a cylindrical warm plasma waveguide is discussed. A new mathematical technique "separation method" which has been applied to the two-fluid plasma model to separate the equations, which describe the system, into two parts, temporal and space parts. Plasma electrons are considered to have a thermal velocity. It is shown that a HF electric field has no essential influence on dispersion characteristics of unstable surface waves excited in a warm plasma waveguide by a low-density electron beam. The region of instability only slightly narrowing and the growth rate decreases by a small parameter and this result has been reduced compared to cold plasma. Also, it is found that the plasma electrons have not affected the solution of the space part of the problem.

MIRAMARE – TRIESTE

November 2002

## Introduction

A great number of publications [1-4], and references cited therein, are devoted to the theoretical study of beam-plasma instability.

The investigation of beam-plasma instabilities evokes interest not only from the point of view of understanding this particular type of instability but also as a method of modeling the instabilities caused by resonant interactions in more complex cases.

In particular, in that of modeling the transition from the laminar to the turbulent state, the turbulence resulting from instabilities, and the control of instabilities.

It is well known that (e.g. [1,5]) propagation of charged particle beams through plasma leads to the development of a great variety of instabilities. Sometimes such instabilities could be used as a basis for the construction of an effective generators of electromagnetic radiation, for heating of magnetically confined plasmas, and for particle acceleration. But often beam instabilities are undesirable. This is true particularly for the experiments on magnetic fusion research where instabilities may cause an appreciable enhancement in the transport of particles and heat across the confining magnetic field. Therefore stabilization of instabilities excited by a beam of charged particles in plasmas is an important part of the more general problem of controlling instabilities in magnetic traps.

The stabilization effect of a uniform HF electric field on a two – stream (Buneman) instability in uniform unbounded plasma has been investigated for the first time in [6]. The dispersion equation for characteristic frequencies of electrostatic oscillations excited by relative motion of electrons and ions in a HF electric field has been obtained and analyzed. The presence of a pump wave strongly modifies the dispersion equation of Buneman instability. As a consequence the growth rate of instability reduces in comparison with the growth rate at vanishing external field amplitude.

The extremely interesting properties of plasma located in a strong high frequency (HF) electric field have stimulated a broad range of theoretical investigations in this promising field of physics.

A broad class of instabilities was predicted to accompany the processes taking place in plasma placed in a HF field. On the other hand, for a number of cases the stabilizing effect of a HF field on unstable plasma states was pointed out. A detailed review of the theory of the plasma HF field interaction was given in ref. [7].

Parametric interaction of external HF electric field with an electrostatic surface wave in an isotropic nonuniform plasma has been investigated in ref. [8] using a special method based on the separation of variables.

Different from previous works on beam-plasma instability [9-11], we study the effects of both the effects of both plasma inhomogeneity and the thermal electron motion on the quenching of the beam-plasma instability. Investigation of beam-plasma interaction presents a great interest for the development

of effective methods via plasma stability, amplification and generation of electromagnetic waves, acceleration of charged particles in plasma, high frequency heating of plasma, and so on [12-14].

It has been shown that [15] dispersion equations describing parametric excitation of surface waves at the boundary of isotropic plasma - vacuum to within the eigen frequency renormalization coincide with the equations that determine the parametric excitation of volumetric waves in uniform unbounded plasma. Proceeding from this conclusion the method for investigation of parametric interaction of external HF electrical field with electrostatic oscillations in isotropic bounded nonuniform plasma has been proposed [8]. The method makes it possible to separate the problem into two parts. The "dynamical" part describes the parametric build up of oscillations and corresponding equations within the renormalization of eigen frequencies coinciding with equations for the parametrically unstable waves in an infinite uniform plasma. Natural frequencies of oscillations and spatial distribution of the amplitude of the self - consistent electrical field are determined from the solution of a boundary - value problem ("space" part) taking into account specific spatial distribution of the plasma density. The proposed approach ("separation method") is significantly simpler than the method ordinarily employed in the theory of parametric resonance in nonuniform plasma [10,15,16]. Therefore it is of special interest to apply the separation method to the solution of different problems involving parametric excitation of electrostatic waves in bounded nonuniform plasma.

In the present work we investigate the suppression of HF electric field on beam-plasma interaction in the cylindrical warm plasma waveguide.

## **2. The influence of HF electric field on the instability of a low - density electron beam passing through plasma waveguide**

We assume that a uniform cold electron beam propagating along radially nonuniform cylindrical plasma waveguide. The radius of the beam  $R$  is supposed to coincide with the radius of the plasma cylinder.

Let us now assume that an electron beam of low density ( $\epsilon_b = n_{ob} / n_0 \ll 1$ ) is passing through a quasineutral plasma with the velocity  $\vec{u}_{b_0}$ . We shall also suppose that both plasma components are at rest ( $\vec{u}_{e_0} = \vec{u}_{i_0} = 0$ ). We take the vector of the external HF field  $\vec{E}_p = \vec{E}_0 \sin(\omega_0 t)$  to be oriented along the axis of the plasma cylinder. Equilibrium density of plasma  $n_{\alpha_0}$  varies along the radius.

The "separation method" has been described [6,17] in application to the problem of parametric excitation of surface waves in a cold isotropic plasma. Here we shall follow this paper. Representing the perturbations of velocity, density and electrical potential in the form  $\delta \vec{V}_\alpha, \delta n_\alpha, \Phi \sim \exp i(m\Psi + kz)$  the linearized set of hydrodynamical equations together with Poisson equation can be reduced to the form

$$\frac{\partial^2 v_{\alpha_1}}{\partial t^2} + \eta v_{\alpha_1} = -\frac{m_e p^2}{e^2} e^{iA_\alpha} \sum_{\beta=b,e,i} v_{\beta_1} \frac{e_\beta^2}{m_\beta} e^{-iA_\beta}, \quad (1)$$

where  $\eta = k^2 V_{th}^2$ ,  $\alpha = e, i$  or  $b$  and  $p$  is a separation constant, and  $v_{\alpha_1}$  is the temporal part of the density perturbations ( $v_\alpha = v_{\alpha_1}(t)v_{\alpha_2}(r)$ ),

$$A_\alpha = (ku_{\alpha_0})t - a_\alpha \sin(\omega_0 t), \text{ and } a_\alpha \equiv \frac{e_\alpha k E_0}{m_\alpha \omega_0^2} \approx a_e. \quad (2)$$

Assuming that the ions are at rest ( $u_{i_0} \equiv 0$ ) and that the frequency of the HF field is much larger than the eigen frequencies of the excited surface waves ( $\omega_0 \gg \omega_{SW} \sim \omega_p$ ), we may use the method of averaging.

Introducing the explicit form of  $A_\alpha$  ( $a_b = a_e \gg a_i$ ), and a new variable  $w_i = \frac{m_e \alpha_i}{m_i \alpha_e} v_{i_1}$ , we find the

final form of the equations, which describes the dynamical (temporal) part of our problem

$$\left. \begin{aligned} \frac{d^2 v_{b_1}}{dt^2} + \varepsilon_b p^2 (v_{b_1} + v_{e_1} e^{ik(u_{b_0} - u_{e_0})t} + w_i e^{ik(u_{b_0} - u_{i_0})t - ia \sin(\omega_0 t)}) &= 0 \\ \frac{d^2 v_{e_1}}{dt^2} + \eta v_{e_1} + p^2 (v_{e_1} + v_{b_1} e^{ik(u_{b_0} - u_{e_0})t} + w_i e^{ik(u_{b_0} - u_{i_0})t - ia \sin(\omega_0 t)}) &= 0 \\ \frac{d^2 w_i}{dt^2} + p^2 \frac{m_e}{m_i} (w_i + e^{ia \sin(\omega_0 t)} (v_{e_1} e^{-ik(u_{e_0} - u_{i_0})t} + v_{b_1} e^{-ik(u_{b_0} - u_{i_0})t})) &= 0 \end{aligned} \right\} \quad (3)$$

where  $\varepsilon_b = n_{0b} / n_0$ .

Here we shall confine our analysis for the influence of the HF electric field on the dispersion characteristics of unstable surface waves excited in a plasma waveguides by an electron beam.

Using the Jacobi-Anger formula  $e^{\pm ia \sin(\omega_0 t)} = \sum_{m=-\infty}^{\infty} J_m(a) e^{\pm im\omega_0 t}$ ,

where,  $J_m(a)$  are the Bessel functions, we obtain from equations (3):

$$\left. \begin{aligned} \frac{d^2 \langle v_{b_1} \rangle}{dt^2} + \varepsilon_b p^2 (\langle v_{b_1} \rangle + (\langle v_{e_1} \rangle + \langle w_i \rangle J_0(a)) e^{i(ku_0)t}) &= 0 \\ \frac{d^2 \langle v_{e_1} \rangle}{dt^2} + \eta \langle v_{e_1} \rangle + p^2 (\langle v_{e_1} \rangle + \langle v_{b_1} \rangle e^{-i(ku_0)t} + \langle w_i \rangle J_0(a)) &= 0 \\ \frac{d^2 \langle w_i \rangle}{dt^2} + p^2 \frac{m_e}{m_i} (\langle w_i \rangle + (\langle v_{e_1} \rangle e^{-i(ku_0)t} + \langle v_{b_1} \rangle) J_0(a)) &= 0 \end{aligned} \right\} \quad (4)$$

where  $\langle v_{\alpha_i} \rangle, \langle w_i \rangle = \frac{\omega_0}{2\pi} \int_0^{2\pi/\omega_0} (v_{\alpha_i}, w_i) dt$ .

The system of equations (4) coincides (if  $V_{th} = 0, n_{ob} = 0, \omega_{P_e}^2 \rightarrow p^2, \omega_{P_i}^2 \rightarrow (m_e/m_i)p^2$ ) with the system describing the HF stabilization of the two-stream instability in a uniform (or nonuniform) unbounded (or bounded) plasma [8] or [17].

If there is no externally injected beam ( $\varepsilon_b = (n_{ob}/n_0) = 0$ ) then the system of equations (4) coincides with the system describing the HF stabilization of the Buneman instability in a nonuniform bounded relativistic cold plasma [18].

### 3. Solution of the "temporal" (time - dependent) equations

According to equations (4), plasma oscillations are then described by the dispersion equation

$$(\omega^2 - \omega_{LF}^2)[(\omega - ku_b)^2(\omega^2 - \omega_{HF}^2 - \eta) - \varepsilon_b p^2(\omega^2 - \eta)] = 0, \quad (5)$$

where

$$\omega_{LF}^2(p) = \frac{m_e}{m_i} p^2 (1 - J_0^2(a)), \quad \omega_{HF}^2(p) = p^2 (1 + \frac{m_e}{m_i} J_0^2(a)) \quad (6)$$

In the case of vanishing electric field amplitude ( $\vec{E}_0 = 0$ ), equation (5) agrees with the dispersion equation which describes the unstable oscillations that excited in a uniform unbounded (bounded) plasma by a low - density electron beam [8,17]. We shall analyze equation (5) (as in [8,17]) in two cases:

#### (a) Non - resonant case ( $ku_b \approx \omega_{HF}$ )

We have from (5)

$$\omega = ku_b \pm \frac{\sqrt{\varepsilon_b} p [ku_b (ku_b - \eta)]^{1/2}}{[(ku_b)^2 - \omega_{HF}^2(p) - \eta]^{1/2}}. \quad (7)$$

Under the conditions

$$p^2 (1 + \frac{m_e}{m_i} J_0^2(a)) > ku_b (ku_b - \eta) > 0, \quad (8)$$

the roots of equation (7) are complex and one of them corresponds to an instability with the growth rate

$$\gamma_{NR} = \frac{\sqrt{\varepsilon_b} p [(ku_b)^2 - \eta]^{1/2}}{[\omega_{HF}^2(p) - (ku_b)^2 - \eta]^{1/2}}. \quad (9)$$

At  $V_{th} = 0$  the result agrees with the result for cold plasma [17].

**(b) Resonant case ( $ku_b \approx \omega_{HF}$ )**

The frequency of unstable oscillations can be represented in the form  $\omega = \omega_{HF}(p) + \Delta\omega$ , where

$$\text{Re } \Delta\omega = - \frac{\varepsilon_b^{1/3} p (1 + \frac{m_e}{m_i} J_0^2(a))^{1/6}}{2^{4/3} [1 + \frac{2\eta}{p^2 (1 + \frac{m_e}{m_i} J_0^2(a))^{1/2}}]^{1/3}}, \quad (10)$$

$$\gamma_R = \text{Im } \Delta\omega = \frac{\sqrt{3}}{2^{4/3}} \frac{\varepsilon_b^{1/3} p (1 + \frac{m_e}{m_i} J_0^2(a))^{1/6}}{[1 + \frac{2\eta}{p^2 (1 + \frac{m_e}{m_i} J_0^2(a))^{1/2}}]^{1/3}}, \quad (11)$$

where  $p$  is determined by the equation of the space part of the problem.

It follows from expressions (8) - (11) that the HF electric field has no essential influence on the dispersion characteristics of unstable surface waves excited in a plasma waveguides by a low - density electron beam. The region of instability only slightly narrows and the growth rate decreases by a small parameter.

The results obtained are in full agreement with the conclusion that an external HF field may have a stabilizing effect on the electron beam-plasma interaction in uniform (or nonuniform) plasma [8,17]. The warm plasma reduced the growth rate.

We conclude that the growth rate of the electron beam-plasma interaction decreases more in a warm plasma than in a cold plasma [17].

**4. Solution of the space (spatial) part of the problem**

The main feature of expressions (6), (7) and (9) consists in an existence of a separation constant  $p$ , which enables us to consider the plasma boundaries.

To find an explicit expression for the constant  $p$  it is necessary to solve the following differential equation (for details see [8]).

$$\frac{1}{r} \frac{d}{dr} \left[ r \varepsilon(p, r) \frac{d\Phi_2(r)}{dr} \right] - \left( \frac{m^2}{r^2} + k^2 \right) \varepsilon(p, r) \Phi_2(r) = 0, \quad (12)$$

where  $\varepsilon(p, r) = 1 - \omega_{pe}^2(r) / p^2$ . If the radial profile of the plasma density and boundary conditions are specified, the solution of equation (12) gives us the desired value of the separation constant  $p$ . The feature of equation (12) is that neither the amplitude of HF electric field nor electron beam parameters enters into it. Therefore equation (12) coincides with equation describing propagation of natural (free of external influence) electrostatic surface waves in a nonuniform plasma cylinder (see e.g., [19] and references therein). Supposing that plasma density is uniform and the interface between plasma and vacuum is sharp. In such case the relations determine the solution of equation (12)

$$\Phi_2(r < R) = C_1 I_m(kr), \Phi_2(r > R) = C_2 K_m(kr), \quad (13)$$

where  $I_m(kr)$  and  $K_m(kr)$  are the modified Bessel functions and  $C_{1,2}$  are the constants. Using the continuity condition of  $\Phi_2$  and  $\varepsilon d\Phi_2/dr$  at the boundary, the following equation is found (see e.g. [19] and references therein)

$$\varepsilon_0(p) + \eta_m(kR) = 0, \quad (14)$$

with

$$\varepsilon_0(p) = 1 - \frac{\omega_{pe}^2}{p^2}, \quad \eta_m(kR) = - \frac{I_m(kR) K_m'(kR)}{K_m(kR) I_m'(kR)}, \quad (15)$$

where the stroke means differentiation with respect to the argument. Equation (14) gives us the relation between the separation constant  $p$ , electron plasma frequency  $\omega_{pe}$  and the axial wave number  $k$ :

$$p = \omega_{pe}(kR)^{1/2} (K_m(kR) I_m'(kR))^{1/2}. \quad (16)$$

In the limiting case of small radius of plasma waveguide ( $kR \ll 1$ ), from (16) we find

$$p_{m=0} = \omega_{pe}(kR) \left[ \frac{1}{2} \ln \left( \frac{kR}{2} \right) \right]^{1/2}, \quad p_{m \neq 0} \approx \frac{\omega_{pe}}{\sqrt{2}}.$$

For  $kR \gg m$  ("thick" waveguide) always  $\eta_m(kR) \approx 1$  and we have  $p = \omega_{pe} / \sqrt{2}$ .

This equation is the same equation in cold plasma waveguide [8,17] i.e.; the warm plasma waveguide has no effect on the space part of the problem.



## 5. Conclusions

The paper deals with parametric excitation of the potential surface waves in bounded nonuniform warm plasma by monochromatic HF electrical field. It is shown that the problem can be reduced to the solution of the "temporal" (parametric) and "stationary" (spatial) parts. The "temporal" part determining frequencies and growth rates of unstable oscillations coincides with accuracy to the redefinition of natural frequencies with equations describing parametric resonance in homogeneous plasma. Natural frequencies of oscillations and spatial distribution of the amplitude of the self-consistent electrical field are determined from the solution of a boundary-value problem ("space" part) taking into account specific spatial distribution of the plasma density. The method described is used to solve the effect of HF field on the excitation of surface waves by an electron beam under the development of instability of low-density beam passing through a cylindrical warm plasma waveguide.

The method was used for the solution of the stabilization effect of a strong HF electric field on beam-plasma interaction in a cylindrical warm plasma waveguide. We solved the "temporal" (time-dependent) equations and obtained the corresponding dispersion equation (5) in a cylindrical geometry, which was analyzed for two cases: nonresonance instability ( $ku_b \approx \omega_{HF}$ ), and resonance one ( $ku_b \approx \omega_{HF}$ ). In both cases the frequency growth rates of the oscillations are obtained (relations (9) and (11)). The separation constant  $p$  is obtained from relation (16), the results are compared with the case when the external electric field is absent ( $\vec{E}_0 = 0$ ) and a cold plasma ( $V_{th} = 0$ ).

In conclusion it is shown that a HF electric field has no essential influence on dispersion characteristics of unstable surface waves excited in a warm plasma waveguide by a low-density electron beam. The region of instability only slightly narrowing and the growth rate decreases by a small parameter and this result has been reduced compared to cold plasma. Also, it is found that the plasma electrons have not affected the solution of the space part of the problem.

## Acknowledgments

The author thanks the Abdus Salam International Centre for Theoretical Physics, Trieste, Italy, for hospitality.

## References

- [1] M. V. Nezlin, Dynamics of Beams in Plasmas, Springer Verlag, Heideleberg, 1991.
- [2] L. S. Bogdankevich and A. A. Rukhadze, Sov. Phys. JETP, V. 62, 1972.
- [3] A. F. Alexanddrov, M. V. Kuzelev and A. N. Khalilov, Sov. Phys. JETP, V. 66, 1987.
- [4] W. H. Amein and Y. A. Sayed, Physica Scripta, V. 50, 1994.

- [5] A. B. Mikhailovskii, Theory of Plasma Instabilities, V. 1,2 Consultants Bureau, NY, 1974.
- [6] Yu. M. Aliev and V. P. Silin, Sov. Phys. JETP, V. 21, 1965.
- [7] V. P. Silin, A Survey of Phenomena in Ionized Gases, IAEA, Vienna, 1968.
- [8] V. V. Demchenko and A. Ya. Omelchenko, Sov. Phys. Radiophysics and Quanta. Electronics, V. 19, 1976.
- [9] H. Bohmer, E. A. Jacson and M. Rather, Phys. Fluids, V. 16, 1973.
- [10] W. H. Amein, V. V. Dolgoplov, A. M. Hussien and K. E. Zayed, Physica, V. 79C, 1975.
- [11] W. H. Amein, V. V. Dolgoplov, A. M. Hussien and K. E. Zayed, Plasma physics, V. 17, 1975.
- [12] P. Kaw, W. Kruer, C. Liu and K. Nishikawa, Advances in Plasma Physics, A. Simon and W. Thompson, (eds), V. 6, J. Wiley, NY, 1976.
- [13] M. V. Kuzelev and A.A. Rukhadze, "Electrodynamics of Dense Electron Beam in Plasma", Mosco, "Nauka", 1990.
- [14] M. V. Kuzelev and A. A. Rukhadze, English Completed Edition, " Plasma Free Electron Lasers", Edition Frontier, Paris, 1995.
- [15] Yu. M. Aliev and E. Ferlengi, Sov. Phys. JETP, V. 30, 1970.
- [16] Yu. M. Aliev and O.M. Gradov, Sov. Phys.-Tech. Phys., V. 17, 1973.
- [17] V. V. Demchenko, Kh. H. El-Shorbagy, Sh. M. Khalil and N. G. Zaki, IEEE Catalog Number 98EX109 [15<sup>th</sup> National Radio Science Conference, 24-26 feb. 1998, Helwan Univ.,  $H_2$ , Cairo, Egypt].
- [18] Kh. H. El-Shorbagy, Physica Scripta, V. 62, 2000.
- [19] V. V. Demchenko and K. E. Zayed, Physica, V. 59, 1972.