

## Particle Simulation of Pedestal Buildup and Study of Pedestal Scaling Law in a Quiescent Plasma Edge

C. S. Chang,<sup>a,b</sup> Seunghoe Ku,<sup>b</sup> H. Weitzner,<sup>a</sup>

<sup>a</sup>*New York University, New York, NY 10012, USA*

<sup>b</sup>*Korea Advanced Institute of Science and Technology, Daejeon 305-710, Korea*

Email: cschang@cims.nyu.edu

R. Groebner, T. Osborne

*General Atomics, San Diego, CA 92186-5608, USA*

### Abstract

A discrete guiding-center particle code XGC (X-point included Guiding Center code) is used to study pedestal buildup and sheared  $E_r$  formation in a quiescent plasma edge of a diverted tokamak. A neoclassical pedestal scaling law has been deduced, which shows that the density pedestal width is proportional to  $T_i^{1/2} M^{1/2} / B_t$  where  $T_i$  is the ion temperature,  $M$  is ion mass and  $B_t$  is the toroidal magnetic field. Dependence on the pedestal density or the poloidal magnetic field is found to be much weaker. Ion temperature pedestal is not as well defined as the density pedestal. Neoclassical electron transport rate, including the collisional heat exchange rate with ions, is too slow to be considered in the time scale of simulation ( $\sim 10$  ms).

### 1. Introduction

Understanding and predicting the edge pedestal behavior in H-mode is one of the most critical issues in ITER Physics. If the H-mode layer is quiescent, the buildup of a pedestal will be governed by the neoclassical physics. We need to know how much of the pedestal physics is neoclassical. However, there is no analytic neoclassical theory to be used for such a study in the edge pedestal region where the gradient scale length is less than an ion banana width, where an unconventional orbit loss due to X-point (X-transport [1]) dominates the conventional neoclassical banana transport, and where the neutral collisions can be as important as the Coulomb collisions. The pedestal physics is closely tied to the orbit squeezing and X-transport in the self-consistent radial electric field ( $E_r$ ) shear. Since most of these effects are highly kinetic, a pedestal buildup must be studied in a fully kinetic plasma-neutral system with a self-consistent radial electric field evaluation in a realistic flux-surface and first-wall geometry. There has been no known prior kinetic code in the literature for studying such an edge pedestal buildup by neutral ionization. A recent Monte Carlo kinetic code work reported by Heikkinen et al [2] simply evaluated an edge radial electric field profile from a prescribed plasma profile without the neutral penetration physics; hence, it was not able to study the pedestal formation physics.

In the IAEA FEC 2002 meeting, we reported the discovery of X-transport as a significant mechanism to help create a sharp negative radial electric field in a thin quiescent layer just inside the separatrix surface [1]. We also reported the launch of a large numerical project to study the kinetic neoclassical pedestal physics. The code XGC (X-point included Guiding-center Code), based upon first principles physics, has recently been completed on a state-of-the-art physical and mathematical platform, and been producing a rather complete and self-consistent picture of the edge-pedestal buildup and other related physics in a quiescent diverted edge [3]. More importantly, XGC has recently been producing a neoclassical pedestal scaling law.

XGC is a massively parallel (typically, using over 2,000 processors) 3D ion kinetic code for a quiescent plasma. It is a full-f, Hamiltonian discrete-particle guiding-center-ion code in cylindrical coordinates, being five dimensional (three in the real and two in the velocity space) in a flux surface and limiter geometry provided by EFIT. It has an advanced momentum and energy conserving Monte Carlo Coulomb collision scheme [4]. The background plasma evolves in time, following the evolution of the radial marker particle distribution. An axisymmetric 2D kinetic Monte Carlo neutral atom diffusion module is built into XGC for a more accurate accounting of the neutral penetration dynamics. Plasma and neutral particles experience charge exchange, ionization, and elastic collisions with each other self-consistently. A continuous heat outflow from the core is included.  $E_r$  is evaluated dynamically from the flux-surface averaged Poisson's equation, consistently with the classical and neoclassical ion polarization currents, in the time-evolving plasma pedestal profile. In the current version, the electrostatic potential is assumed to be a flux function.

## II. Pedestal buildup

A pedestal buildup is simulated after mimicking an L-H transition by turning off an artificial anomalous diffusion coefficient  $D_A$ .  $D_A$  is incorporated into XGC by a random shuffling of each radial particle positions by a random-walk step size smaller than the density gradient scale length, and the shuffling frequency is determined by requiring the anomalously affected density slope being milder than a typical pedestal slope.  $D_A = 1 \text{ m}^2/\text{s}$  is found to be adequate for a DIII-D edge plasma. The neutral atom density at a distance outside the separatrix can be kept constant after the L-H transition or made to be automatically adjusted according to the particle recycling. The poloidal neutral density distribution is prescribed analytically, to model a divertor dominant recycling, limiter recycling, or gas puffing at a specific poloidal location. The plots shown in this report are the simplest case, which uses a time invariant neutral density during the buildup with a uniform poloidal distribution. Effect of the spatiotemporally non-uniform neutral density will be reported elsewhere. However, those

results are not too much different from the simple neutral case presented here.




Fig. 1

Figure 1 shows buildup of a hydrogen ( $M=1$ ) plasma density pedestal by neutral ionization in a purely neoclassical stage after the anomalous diffusion is turned off. Neutral density of  $5 \times 10^{16} \text{ m}^{-3}$  at  $\psi=1.04$  is used. Total elapsed time is 40 toroidal transit time of a 200 eV passing ion. The lines are drawn at one toroidal transit time interval. The green line is the density profile at 40 toroidal transit times. The density pedestal continues to build up. If the simulation is left to go on for a long

enough time, the pedestal buildup makes the edge density profile hollow. The radial location of the inner edge of the density pedestal does not change much as the pedestal density grows (seems to show a very weak inward propagation with rising density). This observation does not agree with a simple minded neutral penetration model by Mahdavi et al in Ref. [5], which reported that the pedestal width is basically the radial neutral mean free path, thus decreases

with pedestal density. The ionized plasma ions must obey the neoclassical radial orbit dynamics. The density pedestal cannot be any sharper than what is shown in Fig. 1.

Figure 2 shows profile evolution of the electrostatic potential, ion temperature, ExB speed, and the parallel flow speed corresponding to Fig. 1. The electrostatic potential basically shows wobbling motions, while the average potential becomes deeper very slowly, ending up to a little deeper than 2 keV for the ion pedestal temperature of 0.5 keV. The same trend is seen in the ExB speed. The fact that the electric field varies slowly assures us that we are in quasi-steady neoclassical state with a vanishing net radial current profile. There is a GAM type of oscillation observed, which decays completely within the simulation time. We continuously add 0.5 MW of heating power to the ions at the inner most radial zone in order to prevent the ion temperature from falling. The average ion kinetic energy  $K_i$ , though, shows some decrease at the inner most zone, indicating that the heat input was not enough. The ion energy pedestal shows a much broader slope than the density pedestal. The broader slope in  $K_i$  at smaller minor radii is due to the usual neoclassical ion thermal conduction, while the steeper slope around the pedestal region is due to the X-transport phenomenon. It should be noted here that  $K_i$  does not drop to a much smaller value on the separatrix, like the density pedestal does. This is the radial ion orbit effect. It should also be noted that the  $K_i$  profile shows non-Maxwellian tail behavior at outside the separatrix. A close examination shows that the ion distribution function is anisotropic: the parallel component of  $K_i$  is greater than the perpendicular component. We find that the orbit squeezing effect causes this phenomenon. The fat banana ions cannot penetrate the high ExB shear layer, while the fat passing orbits can. This non-Maxwellian feature becomes more distinctive at lower ion collisionality. At low ion temperature is low, and/or the scrape density is high, then the ion distribution function becomes almost Maxwellian.




Fig. 2

We note here that the density profiles shown in Fig. 1 accurately defines the inner boundary of a neoclassical density pedestal in a quiescent plasma. However, the outer boundary may not be accurately described in the figure. It is due to the facts that we do not include in the figure the distortion of the gyro orbits in a strong radial electric field (the static classical polarization density), even though we include the polarization current in the time dependent


electric field evolution, and that the gyroviscosity is neglected. Actual outside boundary of a density pedestal may be somewhat broader.

The density and temperature pedestal structures show different responses. The density pedestal is raised by neutral ionization while the ion temperature is lowered by neutral cooling, while both obeying the orbit dynamics of the newly ionized ions in a sheared radial electric field (orbit squeezing and expansion, and X-transport).

### III. Pedestal scaling law

There are two ways to obtain a pedestal scaling law from the present simulation. The first one is to obtain data from numerous pedestal buildup simulations as shown in the previous section. The second method is to find a neoclassical profile solution at the steepest pedestal profile (find the maximal pedestal root) under a given set of pedestal temperature and density, using the observation that a plasma pedestal builds up along the maximal pedestal root allowed by radial orbit dynamics. Getting a data set from the first method is like getting a data set from experiments. It may require much more number of expensive computational runs than the second method. On the other hand, the second method can be extremely systematic, for we can set the density and temperature pedestal heights to the values we want. In the present report, we summarize the result from the second method.

To obtain the steepest density profile solution, we start with a step function density and temperature profile. The neoclassical orbital motions of the individual ions will immediately spread the ions in radius, reaching to an equilibrium profile solution eventually, where the radial ion current vanishes. It is not a real physical process, but only a mathematical process to find the steepest profile state. A proper amount of heat should be added to the ions at the inner most radial location to keep the ion temperature from falling, since the ions will lose energy by neoclassical thermal conduction.



**Fig. 3 Neoclassical edge density pedestal profile with a Tanh fit**

Figure 3 shows a maximal neoclassical density pedestal profile obtained from XGC (dots) and a Tanh-fit (red line). A DIII-D equilibrium is used with the axis-magnetic field 1 T, pedestal ion temperature 1 keV and pedestal density  $5 \times 10^{13} \text{ cm}^{-3}$ . The horizontal axis is the normalized poloidal flux  $\psi$ . Agreement with the Tanh-fit is remarkably good, as was in the experimental pedestal data fittings. The pedestal width obtained here is about 1 cm, which is somewhat narrower than, but in the right ball park with the experimentally measured values. The Tanh-fit defines a consistent pedestal width for scaling studies.

Figure 4 shows scaling of a hydrogen density pedestal width (in  $\psi$ ) against the square root of the pedestal ion temperature, obtained from the Tanh-fit to the maximal pedestal shapes for various ion temperature values. Deuteron plasma yields wider width by  $M^{1/2}$ . It shows an offset linear behavior in  $T_i^{0.5}$  with the offset value some where between  $T_i = 40 - 100 \text{ eV}$ . As mentioned before, the ion temperature pedestal shape is not well defined. Thus, the definition of the ion temperature pedestal value is ambiguous. The ion pedestal temperature in the present study is measured at  $\psi = 0.9$ , which can be different from what is used in the real experimental analysis.

A similar scaling study has been performed by varying the axis magnetic field, which shows a rather surprising result (Fig. 5).. It shows a strong  $1/B_T$  dependence, but only a very weak  $1/B_P$  dependence. The density width dependence on the pedestal density has also been examined. The maximal pedestal scaling shows that the density width does not vary much as the pedestal density is varied, confirming the observation made in the actual buildup process shown in Sec. II (Fig. 1). Thus, the neoclassical density pedestal width  $\Delta_n$  does not increase with the poloidal ion gyroradius, nor it decreases with density, but scales approximately as  $\Delta_n \propto T_i^{1/2} B_T^{-1} M^{1/2}$ . This scaling seems to agree with a preliminary experimental result on C-MOD [6] (but not conclusive yet) and a new regression study of DIII-D data [7].




Fig. 4. Density pedestal width vs SQRT of pedestal  $T_i$




Fig. 5. Density pedestal width v.s.  $1/B$ (axis, Tesla)

Another interesting finding from this study is that there is often a positive toroidal momentum source in a strong pedestal edge, supporting a recent observation on C-MOD [John Rice, 2003 APS-DPP, Invited Paper]. Without a toroidal momentum source, a conventional neoclassical theory and XGC would predict a negative toroidal flow generation in a steep pedestal. However, a positive toroidal rotation is often observed from XGC in a strong pedestal region at a separatrix edge. It is also found that the ion distribution function at the foot of the pedestal (in the scrape-off layer) can be highly anisotropic, in agreement with a recent observation on DIII-D [Keith Burrell, 2003 APS], which suggests that an anisotropic radial force balance equation should be used to deduce an  $E_r$  profile from the experimentally observed pressure gradient and plasma flows.

#### IV. Conclusion and discussions

Assuming that the plasma edge is turbulence free, we have performed a 3D kinetic neoclassical simulation of plasma ions in a realistic flux surface and first wall geometry using the XGC code. For a more reliable simulation of the neutral penetration physics, an axisymmetric kinetic Monte Carlo neutral particle transport module is built into the code. In this purely neoclassical system, the plasma electron transport is too slow to evolve within the simulation time scale (about 10 ms). The electron-ion coupling is also too weak to affect evolution of electron temperature profile within the simulation time. Thus, buildup of electron temperature pedestal at a similar rate as the density pedestal buildup is beyond neoclassical analysis. An independent electron transport mechanism (a residual electron turbulence) may be necessary to model an electron temperature pedestal buildup.

Pedestal scaling law has been deduced, which shows that the neoclassical pedestal width is proportional to  $T_i^{1/2}/B_t$  where  $T_i$  is the ion temperature and  $B_t$  is the toroidal magnetic field. Density pedestal width does not show much dependence on  $B_p$  or density, in disagreement with some recent reports. A residual turbulence may modify this result. Comparison with experimental data will reveal how much of the experimentally observed pedestal is from the neoclassical process. A preliminary study shows that a significant portion is neoclassical.

### Acknowledgement

The first author wishes to acknowledge helpful discussions with K. Burrell (GA) and R. Maingi (ORNL).

### References

- [1] 2002 IAEA FEC, IAEA-CN-94/THP3-01; Chang, C. S., Kue, Seunghoe, and Weitzner, H., "X-Transport: A baseline non-ambipolar transport in a diverted tokamak plasma edge," Phys. Plasmas **9** (2002) 3884;
- [2] Heikkinen, J. A., Kiviniemi, T. P., Peeters, A. H., Phys. Rev. Lett. **84** (2000) 487.
- [3] Chang, C.S., Kue, Seunghoe, and Weitzner H., Phys. Plasmas **11**, 2649 (2004).
- [4] W. X. Wang, N. Nakajima, et al., Plasma Phys. Cont. Fusion **41**, (1999) 1091.
- [5] M. A. Mahdavi, R. Maingi, R. J. Groebner, A. W. Leonard, T. H. Osborne, and Gary Porter, Phys. Plasmas **10**, (2003) 3984.
- [6] J. W. Hughes, D. A. Mossessian, A. E. Hubbard, et al, Phys. Plasmas **9**, 3019 (2002).
- [7] C. S. Chang and R. Groebner, ITPA, Lisbon, Portugal