

LTH788
HU-EP-08/11
DESY 08-041
SFB/CPP-08-23

The η' meson from lattice QCD.



K. Jansen^(a), C. Michael^(b), C. Urbach^(c)

^(a) DESY, Zeuthen, Platanenallee 6, D-15738 Zeuthen, Germany

^(b) Theoretical Physics Division, Dept. of Mathematical Sciences,
University of Liverpool, Liverpool L69 7ZL, UK

^(c) Institut für Elementarteilchenphysik, Fachbereich Physik,
Humbolt Universität zu Berlin, D-12489, Berlin, Germany

Abstract

We study the flavour singlet pseudoscalar mesons from first principles using lattice QCD. With $N_f = 2$ flavours of light quark, this is the so-called η_2 meson and we discuss the phenomenological status of this. Using maximally twisted-mass lattice QCD, we extract the mass of the η_2 meson at two values of the lattice spacing for lighter quarks than previously discussed in the literature. We are able to estimate the mass value in the limit of light quarks with their physical masses.

1 Introduction

There is considerable interest in understanding hadronic decays involving η and η' in the final state. The phenomenological study of hadronic processes involving flavour singlet pseudoscalar mesons makes assumptions about their composition. Here we address the issue of the nature of the η and η' from QCD directly, making use of lattice techniques.

Lattice QCD directly provides a bridge between the underlying quark description and the non-perturbative hadrons observed in experiment. The amplitudes to create a given meson from the vacuum with a particular operator made from quark fields are measurable, an example being the determination of f_π . It also allows a quantitative study of the disconnected quark contributions that arise in the flavour singlet sector. The lattice approach provides other information such as that obtained by varying the number of quark flavours and their masses.

The disconnected diagram responsible for giving the flavour-singlet pseudoscalar meson a mass (in the chiral limit) is closely related to the fluctuations in the topological charge. This link will be discussed shortly in this paper and in more detail in a later publication [1].

From the chiral perturbation theory description (for a review, see ref. [2]), one expects the mixing of η and η' to be most simply described in a quark model basis. In the flavour singlet sector, for pseudoscalar mesons, we then have contributions to the mass squared matrix with quark model content $(u\bar{u} + d\bar{d})/\sqrt{2}$ and $s\bar{s}$ (which we label as nn and ss respectively):

$$\begin{pmatrix} m_{nn}^2 + 2x_{nn} & \sqrt{2}x_{ns} \\ \sqrt{2}x_{ns} & m_{ss}^2 + x_{ss} \end{pmatrix}. \quad (1)$$

Here m corresponds to the mass of the flavour non-singlet eigenstate and is the contribution to the mass coming from connected fermion diagrams while x corresponds to the contribution from disconnected fermion diagrams. Thus m_{nn} is the pion mass. Because of mixing, as will be discussed, m_{ss} does not correspond to any specific meson, but its mass can be estimated assuming that the non-singlet pseudoscalar mass-squared is linear in the quark mass. So $m_{ss}^2 = 2m_{ns}^2 - m_{nn}^2$, that is $2m_K^2 - m_\pi^2$, leading to $m_{ss} = 0.687$ GeV. For a discussion from lattice results of small corrections to this assumption, see ref. [3].

The mixing between the nn and ss flavour singlet channels must produce the experimental $\eta(548)$ and $\eta'(958)$. One approximation used historically is that of SU(3) flavour symmetry for which the two physical states will be a flavour octet ($nn - \sqrt{2}ss$) and a flavour singlet ($\sqrt{2}nn + ss$). The η is then identified as primarily the flavour octet while the η' is the flavour singlet.

Using as input m_{nn} and m_{ss} and requiring that the output masses (m_η and $m_{\eta'}$) are correctly reproduced, the three mixing parameters x cannot be fully

determined. It is usual to express the resulting one parameter freedom in terms of a mixing angle, here defined by

$$\eta = \eta_{nn} \cos \phi - \eta_{ss} \sin \phi, \quad \eta' = \eta_{nn} \sin \phi + \eta_{ss} \cos \phi. \quad (2)$$

In our lattice study, we have two degenerate light quarks in the sea but no explicit strange quark. Since we do not consider partially quenched QCD here, we only study the top left corner of the mixing matrix. The flavour singlet pseudoscalar meson (called η_2 in this case) will then have mass-squared $m_{nn}^2 + 2x_{nn}$.

From the phenomenological analysis of the full mixing matrix, one can then estimate the mass of the η_2 meson. One such phenomenological analysis, motivated by lattice input (basically the magnitude of x for strange quarks), gave [3] values $x_{nn} = 0.292$, $x_{ns} = 0.218$, $x_{ss} = 0.13$ GeV². This assignment corresponds to a mixing angle ϕ close to 45° and to a mass of the η_2 meson of 0.776 GeV. This mass value is plausible since it is close to the (mass-squared weighted) average mass of the η and η' . Moreover, it is somewhat lighter than the η' meson, which has an additional contribution to its mass from strange quark loops (x_{ss}).

Note that the mass of the η_2 meson comes from two components: m_{nn} which decreases with decreasing quark mass and x_{nn} which, in the phenomenological study, increases with decreasing quark mass. It is thus important to study the η_2 meson at light quark masses to explore this. For instance, with the parameters of the phenomenological model described above, the η_2 meson made of light quarks of mass equal to the strange quark mass would have mass-squared $m_{ss}^2 + 2x_{ss}$ giving mass 0.862 GeV. Thus we can expect a rather flat dependence of the η_2 mass on the underlying quark mass.

Lattice QCD is able to access flavour-singlet states, but at a cost which has limited these explorations in practice. Since the correlator of a flavour singlet meson created at x and annihilated at y will have two contributions: a connected contribution where both quark and antiquark propagate from x to y and a disconnected contribution where there is a quark loop at x and another at y . This latter contribution needs to be measured at many values of x and y , so it is optimal to use a stochastic method to achieve this. One then evaluates the disconnected contributions (loops) at all x , and from combining them pairwise, one can evaluate the disconnected contribution to the meson correlator. The stochastic method introduces noise, which can be reduced by appropriate variance reduction techniques. The goal is to reduce the noise to a level lower than the inherent noise coming from the underlying gauge-field variation. We shall show that the formalism known as twisted mass lattice QCD allows a very effective variance reduction method to be applied.

We present details of our lattice methods and explain the stochastic method used to evaluate the flavour singlet pseudoscalar meson correlators (both connected and disconnected). Because the signal from the disconnected correlator is relatively noisy, we discuss several strategies for reducing this error. From a

combination of methods, we are able to present quite precise results, which enable us to discuss the continuum and chiral limit of the flavour singlet pseudoscalar mass. We compare with previous determinations and summarise.

2 Twisted mass lattice QCD action

In the gauge sector we employ the so-called tree-level Symanzik improved gauge action (tlSym) [4], viz.

$$S_g = \frac{\beta}{3} \sum_x \left(b_0 \sum_{\substack{\mu, \nu=1 \\ 1 \leq \mu < \nu}}^4 \{1 - \text{Re Tr}(U_{x, \mu, \nu}^{1 \times 1})\} + b_1 \sum_{\substack{\mu, \nu=1 \\ \mu \neq \nu}}^4 \{1 - \text{Re Tr}(U_{x, \mu, \nu}^{1 \times 2})\} \right),$$

with the bare inverse gauge coupling β , $b_1 = -1/12$ and $b_0 = 1 - 8b_1$. The fermionic action for two flavours of maximally twisted, mass degenerate quarks in the so called twisted basis [5, 6] reads

$$S_{\text{tm}} = a^4 \sum_x \{ \bar{\chi}(x) [D_W[U] + m_0 + i\mu_q \gamma_5 \tau^3] \chi(x) \}, \quad (3)$$

where m_0 is the untwisted bare quark mass, μ_q is the bare twisted quark mass, τ^3 is the third Pauli matrix acting in flavour space and

$$D_W[U] = \frac{1}{2} [\gamma_\mu (\nabla_\mu + \nabla_\mu^*) - a \nabla_\mu^* \nabla_\mu]$$

is the mass-less Wilson-Dirac operator. ∇_μ and ∇_μ^* are the forward and backward covariant difference operators, respectively.

The mass term m_0 is tuned to maximal twist (as described in ref. [7]) at our lightest μ_q parameter. This guarantees $\mathcal{O}(a)$ improvement [6]. This was shown to work excellently in the quenched approximation [8, 9], and there are good indications that this holds true also with dynamical fermions [10, 11]. Moreover, this formalism has been found to give a very effective way to reach light quark masses [7, 12, 13]. As such, it provides an attractive route to evaluate the properties of the flavour-singlet pseudoscalar mesons.

There is one complication, however, namely that twisted mass lattice formalism breaks flavour and parity symmetries at finite value of the lattice spacing a . These symmetries are restored in the continuum limit and the theory is well defined (so providing a valid regularisation) at finite lattice spacing. For our present purpose, this flavour-breaking causes the π^+ and π^0 mesons to have a mass splitting (of order a^2). Moreover the π^0 has contributions from disconnected diagrams [14, 7]. The analysis of the flavour-singlet pseudoscalar meson is not significantly affected by this and it can be studied in the same way as was

Ensemble	$L^3 \times T$	β	$a\mu_q$	κ	r_0/a
B_1	$24^3 \times 48$	3.9	0.0040	0.160856	5.22(2)
B_3			0.0085		
B_6	$32^3 \times 64$	3.9	0.0040	0.160856	
C_1	$32^3 \times 64$	4.05	0.003	0.157010	6.61(3)
C_2			0.006		

Table 1: Summary of ensembles produced by the ETM collaboration used in this work. We give the lattice volume $L^3 \times T$ and the values of the inverse coupling β , the twisted mass parameter $a\mu_q$, the hopping parameter $\kappa = (2am_0 + 8)^{-1}$ and the Sommer parameter r_0/a in the chiral limit from ref. [7, 10].

used previously in Wilson-based lattice formalisms [3, 15]. For a recent review see ref. [16].

For later convenience we introduce the following notation

$$D_{u,d} = D_W + m_0 \pm i\mu_q \gamma_5 \quad (4)$$

for the fermion matrix of the two degenerate quark flavours (here labelled u and d) separately.

The results presented in this paper are based on gauge configurations as produced by the European Twisted Mass collaboration (ETMC). The details of the ensembles are described in ref. [10]. In particular, we concentrated for this paper on the ensembles labelled B_1, B_3, B_6 and C_1 and C_2 . We have compiled the details for those ensembles in table 1. The B -ensembles correspond to a value of the lattice spacing of about $a \sim 0.09$ fm and the C -ensembles to $a \sim 0.07$ fm. The spatial lattice size is of about $L \sim 2.2$ fm for all ensembles used here, apart from B_6 , which has identical parameters to B_1 but $L \sim 2.7$ fm.

3 Neutral particles in twisted mass QCD

In quenched or partially-quenched lattice QCD, the disconnected contribution to the flavour singlet meson does not combine properly with the connected contribution to give a physical state. To avoid this problem, it is mandatory to study flavour singlet states in full QCD - with sea quarks having the same properties as valence quarks. Then the spectrum of flavour singlet states is well defined and can be extracted from the t -dependence of the full correlator. Here we focus on the case where there are two degenerate light quarks (called $N_f = 2$) which is a consistent theory in which to study the flavour singlet mesons.

One promising way to study light quarks is using twisted mass QCD at maximal twist. In the case of twisted mass fermions, the bilinear operator appropriate to create the η_2 state is $\bar{\psi}\gamma_5\psi$ which on transformation into the twisted basis (used in lattice evaluation) will become $\bar{\chi}\tau_3\chi$, where τ_3 acts in the (u, d) flavour space.

This amounts to evaluating, in the lattice basis, the difference of the disconnected loop between u and d quarks. As already reported [17, 18], this enables a very efficient variance reduction technique to be used to evaluate the disconnected diagram relevant to the η_2 .

The key observation is the following relation for the inverse D_u and D_d :

$$1/D_u - 1/D_d = -2i\mu_q(1/D_d)\gamma_5(1/D_u) . \quad (5)$$

Consider now the disconnected loop $\sum X(1/D_u - 1/D_d)$ where X is some γ -matrix (here I) and/or colour-matrix and the sum is over space. The conventional method involves solving $\phi_r = (1/D_u)\xi_r$ with stochastic volume sources ξ_r . The desired result is then obtained from

$$\sum X/D_u = \sum \langle \xi^* X \phi \rangle_r ,$$

where the average is over noise samples (labelled r). However, the case mentioned above can be evaluated efficiently using the ‘one-end-trick’ [19, 20]. Then the required disconnected loop is given by

$$\sum X(1/D_u - 1/D_d) = -2i\mu_q \sum \langle \phi^* X \gamma_5 \phi \rangle_r .$$

This has signal/noise $V/\sqrt{V^2} = 1$ which is much more favourable than the conventional method with signal/noise $1/\sqrt{V}$ (here $V = L^3T$). For ensemble B_1 , we find the zero-momentum disconnected loop has a standard deviation of $\sigma = 18$ for the variation with gauge configuration and time-slice (in the limit of no stochastic error) whereas the stochastic noise on this loop has standard deviation of $\sigma = 87$ from 24 samples of volume source (conventional method) but only $\sigma = 7.5$ from the above method (with 12 samples). So 12 inversions give the disconnected correlator from all t to all t' with no significant increase in errors from the stochastic evaluation.

We also use a non-local source/sink for the meson, constructed using ‘fuzzed’ links of length $6a$ as described in refs. [21, 18], which can be evaluated by replacing X by the corresponding fuzzed gauge links. Thus twisted mass lattice QCD allows a very efficient way to evaluate flavour-singlet pseudoscalar disconnected loops, and combining them, the required disconnected correlator $D(t)$.

Evaluating the connected correlator $C(t)$ for the neutral pseudoscalar meson as described elsewhere [18], then we have the full information $C(t) - 2D(t)$ needed to explore the spectrum in the η_2 channel. We have a 2×2 matrix of such correlators available (local or non-local at sink/source). We could also explore pseudoscalar correlators obtained by creating the η_2 state with $\bar{\psi}\gamma_0\gamma_5\psi$ which on twisting becomes $\bar{\chi}\gamma_0\gamma_5\chi$. This does not allow our very efficient variance reduction to be applied, so the disconnected contributions are evaluated using a hopping parameter variance reduction [22, 18]. Even so, they are sufficiently

β	$a\mu_q$	L/a	No.	$am(\eta_2)$	χ^2/dof
3.90	0.0040	24	888	0.47(8)	3/(24 - 6)
3.90	0.0040	32	184	0.37(6)	10/(24 - 6)
3.90	0.0085	24	245	0.41(7)	8/(24 - 6)
4.05	0.0030	32	198	0.34(7)	10/(24 - 6)
4.05	0.0060	32	188	0.37(5)	11/(24 - 6)

Table 2: Results for the flavour singlet pseudoscalar meson from zero momentum correlators. The number of configurations analysed for neutral correlators (connected and disconnected) is shown (No.). The fits are to t -range 3-10 with 2 states and a 2×2 matrix of correlators (local and fuzzed at source and sink). The excited state mass is around $am' = 1.3$ at $\beta = 3.9$.

noisy that we have been unable to use them to constrain the fits or to determine the η_2 decay constant.

We measure the auto-correlation versus trajectory number for our correlators. The largest auto-correlation is found for the disconnected contribution. To explore this more fully, we investigated the auto-correlation for the time-slice sum (i.e. zero momentum) of the quark loop at a given time versus trajectory, finding comparable auto-correlation times to that of the plaquette [10]. This is not unexpected, since the disconnected contribution is related to the topological charge density and thus both it and the plaquette are sensitive to the vacuum structure encoded in the gauge configurations. In order to cope with this measured auto-correlation, we block the data into sufficiently large blocks (more than 80 trajectories) to remove any effect on our final error estimates. Note that the gauge configurations we use for our measurements are usually well separated in units of trajectories, such that even after blocking 80 trajectories we are left with a sufficiently large number of (then independent) measurements.

To explore the signal, we plot the effective mass versus t for ensemble B_1 in fig. 1, where we have used the variational basis from t -values 3 and 4 to optimise the ground state contribution. The zero momentum results (open symbols) show a plateau, although statistical errors are large, as will be discussed later. The results for the two lattice spatial volumes available in this case agree well.

In order to explore options to reduce the errors, we also evaluate the full η_2 correlator for momentum 1 (in units of $2\pi/L$). Since we evaluate the disconnected correlator for momentum 1 in each of the 3 spatial directions, we obtain an improved estimate of D (for example with relative error reduced to 11% at $t = 10$ compared to error of 14% for zero-momentum). Since the symmetry classification of mesonic states is less sharp at non-zero momentum, the interpretation of the lattice spectrum is thus less straightforward. In practice, we find that the small reduction in error on D does not translate into a significant reduction in the error in determining the underlying mass of the η_2 state.

We make a fit to the combined 2×2 matrix of zero-momentum correlators




Figure 1: η_2 effective mass (from a variational basis) versus t . Results are from ensemble B_1 ($L = 24$) and B_6 ($L = 32$) (both $a\mu_q = 0.004$). The results from momentum $k = 1$ (in units of $2\pi/L$) have been plotted as mass values assuming $E^2 = m^2 + k^2$. The results are slightly displaced for better readability.

for a t -range of 3-10 and with two states. The results from an uncorrelated fit to blocked data are presented in table 2. These fits are stable to varying the t -range.

Before analysing these results further, we now discuss why the errors are so large for the η_2 disconnected correlator, despite the fact that we measure all t and t' , we use many gauge configurations and our stochastic errors are small. As described above, we have investigated the autocorrelation (versus trajectory) of the disconnected contribution and find no statistically significant evidence of any autocorrelation beyond 40 trajectories. We block our data into larger blocks (typically covering 80 trajectories) to avoid any increase in error from this source.

The origin of the large error is that the signal for the disconnected part of the correlator comes from only a small part of the total data sample. Here we concentrate on the zero-momentum method to illustrate this. For instance (for ensemble B_1 from 48 t -values for 888 gauge configurations) with $|t' - t| = 10$, 2.1% of the data contributes 26% of the signal. Thus the statistical impact of the data set is smaller than expected since parts of the data have big fluctuations (in a fermionic loop related to topological charge density).

Another way to illustrate this error problem is that from the first 444 gauge configurations (covering 4440 trajectories) we find $D/C = .96(29)$ at $|t' - t| = 16$, while for the second 444 gauge configurations we find $D/C = 0.47(23)$. Positivity requires $D/C < 0.5$. So the error estimate for certain quantities from even such a big ensemble as 5000 trajectories can be underestimated compared to having

10000 trajectories, while other quantities do not show such a problem on the same set of gauge configurations.

So even more configurations, than we have here, would be needed to get reliable and small errors in the case of disconnected contributions. This same conclusion has been obtained before, most strongly in a study involving staggered fermions [23].

4 Error reduction strategies

Because the statistical error on the η_2 mass is still large, we now discuss various strategies to reduce it while retaining the same set of gauge configurations. Some of these strategies do indeed reduce the statistical error, but at the expense of introducing a systematic error that has to be discussed.

4.1 Excited state removal

We now consider replacing the connected neutral pseudoscalar correlator C by just the ground state contribution. This has been considered previously [24]. The basic idea is that the disconnected contributions (D) are only big for the lightest flavour-singlet pseudoscalar (the η_2) and not for excited states. In other words, π and η_2 are split but π' and η_2' are almost degenerate. This might happen because the topological charge fluctuations in the vacuum, which give the flavour singlet states a different mass, are more strongly coupled to the ground state η_2 than to its excited states.

In order to have no significant excited state contributions in the appropriate flavour-singlet correlator $C - 2D$, one should remove them from C . Then neither C nor D will have excited state contributions, and a one state fit to $C - 2D$ should be possible down to quite low t -values, as indeed we shall find. The extension of this argument to twisted mass lattice QCD is not trivial, since there are also disconnected contributions to the neutral pion itself. Near the continuum limit, however, the argument goes through as before.

In detail, we use fits of the form $c \cosh(-m(T/2 - t))$ to the zero momentum data for the connected neutral correlator in the t -range 10-23 to determine the mass m and the coupling c . Note that this mass m corresponds to a neutral pion only if it is interpreted as a (partially-quenched or mixed-action) Osterwalder-Seiler state [14]; we shall call it the ‘connected pion’. We then use our best fit parameters of m and c to construct the connected pion correlator, and we assign it with an error using the bootstrap method. This constructed connected pion correlator is finally used together with the disconnected contribution to build the full η_2 correlator, which we use even below $t = 10$. As shown in fig. 2, the effective mass is now essentially flat. Hence our result is consistent with dominance by

β	$a\mu_q$	L/a	$am(\eta_2)$	χ^2/dof
3.90	0.0040	24	0.37(5)	0.2/(27 - 3)
3.90	0.0040	32	0.30(4)	2.4/(27 - 3)
3.90	0.0085	24	0.36(5)	3.0/(27 - 3)
4.05	0.0030	32	0.26(5)	1.8/(27 - 3)
4.05	0.0060	32	0.26(3)	0.8/(27 - 3)

Table 3: Results for the flavour singlet pseudoscalar mass from zero-momentum correlators. The connected correlator is taken from the fitted ground state component. The fits are to t -range 2-10 with 2 states and a 2×2 matrix of correlators (local and fuzzed at source and sink).

a single η_2 state down to surprisingly small t -values, where the data have small statistical errors.

We can then make one state fits to the resulting 2×2 matrix of correlators with t -range 2-10 and the results are reported in table 3. This shows that we achieve a significant reduction in the error on the η_2 mass.

4.2 Point to point correlators

We discussed above the possibility of using correlators at smaller t to reduce errors. Another approach, which we now discuss, is to reduce errors at all t -values by focussing on a different quantity. The method we describe has been used recently [25] in a study of the disconnected contributions to the η_c meson.

It is mandatory to use an all-to-all method to estimate the disconnected loops $d(\mathbf{x}, t)$ at all \mathbf{x} and t . The disconnected correlator of interest reads

$$D(\mathbf{x} - \mathbf{y}, t - t') = \langle d(\mathbf{x}, t)d(\mathbf{y}, t') \rangle,$$

assuming that $d(\mathbf{x}, t)$ has no vacuum expectation value, as is the case for the flavour-singlet pseudoscalar meson because of the combined flavour-parity symmetry of TMQCD. The approach we have considered above is to sum D over the spatial coordinates \mathbf{x} and \mathbf{y} . This has the advantage that the zero momentum correlator is studied which has the simplest theoretical interpretation: as a sum of exponentials for each state. Now D is actually peaked when $|\mathbf{x} - \mathbf{y}| \approx 0$ and is small at large $|\mathbf{x} - \mathbf{y}|$. Moreover we find that the absolute error is largely independent of $|\mathbf{x} - \mathbf{y}|$, so the relative error is thus smallest when $|\mathbf{x} - \mathbf{y}| \approx 0$. We illustrate the dependence of D on $r = |\mathbf{x} - \mathbf{y}|$ in fig. 3. The zero momentum contribution is a sum over this r -distribution with weight approximately r^2 and this enhances the larger r region which has larger relative statistical errors. Basically the zero-momentum sum picks up (inherent gauge-time variation) noise by summing over large r where the signal is unimportant but the noise is still significant.




Figure 2: Effective mass for zero-momentum η_2 correlator for ensemble B_1 ($a\mu_q = 0.004$) taking as the connected contribution the ground state only. The horizontal line is the value given by a fit to the correlator data in the t -range 2-10. Also shown are results with point-to-point correlators with $r = 2a$, as described in the text. The results are slightly displaced for improved readability.

For example, we find for ensemble B_1 with $|t - t'| = 10$ that the relative error on D when $r = 0$ ($x = y$) is 50% of the relative error on D summed over both \mathbf{x} and \mathbf{y} . Thus the peak signal to noise is twice as big as that summed over relative spatial separation. It might be thought that taking $r \neq 0$ would give an even smaller relative error, since 6 spatial directions are averaged (for spatial separations on axis). Although we find less error reduction than $1/\sqrt{6}$ for $r \neq 0$, we do make use of a further small error reduction and we select $r = 2a$ (on axis).

We now discuss how to extract the conventional physics (i.e. mass values) from correlators at fixed r (rather than summed over all r). Consider a correlator corresponding to a meson of mass m . In the region of t where the ground state contribution dominates, we expect the correlator to be dependent explicitly on the meson mass when point operators are used at source and sink. Indeed one way to extract this behaviour is by evaluating the 4-dimensional lattice Fourier transform:

$$C(t, \mathbf{r}) = \frac{1}{L^3 T} \sum_{p_0, \mathbf{p}} \frac{c e^{-ip_0 t - i\mathbf{p} \cdot \mathbf{r}}}{\hat{p}^2 + m^2}, \quad (6)$$

to model the data, where, to respect the periodic boundary conditions (for a meson), $p_i = 2\pi n_i/L$ and $p_0 = 2\pi n_0/T$. In eq. 6, we take $\hat{p}_i = 2 \sin(p_i/2)$ which corresponds to the most local lattice realisation of the derivative in the Klein Gordon action. Other implementations of the derivative would shift the




Figure 3: Disconnected contribution to the η_2 correlator at time t and spatial separation r from point source and sink. Results are from ensemble B_1 ($a\mu_q = 0.004$) from 319 gauge configurations. The curves are from the model fit described in the text.

mass by corrections of order $(ma)^2$ as discussed long ago [26]. We have checked that this expression correctly reproduces the r -dependence for the charged pion correlator in the t -region where the ground state dominates, using the mass value previously determined from fitting the zero momentum correlator. Thus, though less familiar, one can use the expression of eq. 6 to extract the mass value from the t -dependence at fixed r , in place of the usual expression $c(e^{-mt} + e^{-m(T-t)})/(2m)$ used at zero momentum.

To exploit this error reduction for the η_2 , we combine the disconnected contribution (with $r = 2a$ as discussed above) with the neutral pion connected contribution also evaluated with $r = 2a$. This we evaluate using point sources. However, it is optimal, as discussed in the preceding section, to use only the ground state contribution to the connected neutral pion correlator, as we now discuss.

For this ground-state connected component, we use eq. 6 with as input the mass m and coupling c determined from the zero-momentum fit to the neutral (connected) pion. The dependence of the resulting η_2 correlator on t (at fixed $r = 2a$) can then be parametrised again by eq. 6. One way to illustrate this, as shown in fig. 2, is by plotting an ‘effective mass’ (here defined as the mass parameter that solves the dependence given by eq. 6 for two adjacent t -values). This shows that we again have a rather constant ‘effective mass’ even from low t -values, consistent with a description by one state only. Moreover, the errors

β	$a\mu_q$	L/a	$am(\eta_2)$	χ^2/dof
3.90	0.0040	24	0.403(21)	2.0/(9 - 2)
3.90	0.0040	32	0.365(19)	0.1/(9 - 2)
3.90	0.0085	24	0.380(18)	0.7/(9 - 2)
4.05	0.0030	32	0.302(24)	0.0/(9 - 2)
4.05	0.0060	32	0.308(18)	0.0/(9 - 2)

Table 4: Results for the flavour singlet pseudoscalar mass using point-to-point correlators for the disconnected part and extracting the connected part from the fitted ground state component.

on the ‘effective mass’ are smaller than from the fixed momentum methods used above.

To determine the η_2 mass by this method, we then fit the resulting combined η_2 correlator to a two parameter expression (free mass and coupling) according to eq. 6. Since this approach is less familiar, we illustrate in fig. 4 the fit versus t and also in fig. 3 the r -dependence of the disconnected contribution compared to the measured data. In detail, we use fits to the zero momentum data for the connected neutral correlator for the t -range 10-23 to determine the connected pion mass m and the coupling c , as above. From the bootstrap ensemble of values of m and c , we determine, at each t -value, the point-to-point connected pion correlator at fixed $r = 2a$ with its associated error. This is then used together with the measured point-to-point disconnected contribution to construct the full η_2 correlator. The result is then fitted with one state using eq. 6 and a t -range 2-10 to obtain the η_2 mass value given in table 4.

Using different t -ranges gives the same mass value, within errors. Compared to using the zero-momentum approach described previously, the statistical error on the mass is much smaller. Here we have used local source and sink operators (since in this case the fixed r behaviour is given without extra parameters). What is especially helpful, however, is that the resultant data is well fitted by only one contribution for $t > a$.

We have presented this new and powerful method for one ensemble (B_1) but we also apply it (using $r = 2a$) to the other cases considered and the results are collected in table 4.

This point-to-point approach provides a useful reduction in the statistical error, though at the potential cost of introducing a source of systematic error in relating the quantity measured to that actually required. For local meson creation and destruction operators, the t -dependence at fixed r is given by the same parameters as the conventional zero-momentum case, up to possible non rotation-invariant contributions of order a^2 which we expect to be insignificant at large $t^2 + r^2$. We also find strong evidence of ground state dominance, which enables us to use the more precise data from smaller t -values.




Figure 4: Singlet pseudoscalar correlator for ensemble B_1 ($\mu_q = 0.004$) with local point sink and source. The correlator is illustrated for spatial separation $r = 2a$ versus time t . The upper solid line is the connected contribution from the ground state neutral meson while the circles show the disconnected contribution. The squares are the net contribution to the η_2 and the behaviour from a one state fit to them is also shown. Note that the one state fit is not exponential, as discussed in the text.

5 Summary of η_2 mass

In order to compare the error-reduced results from different lattice spacings, we plot them using the r_0/a -values given in table 1 to create dimensionless quantities. We see in fig. 5(a) that the results are consistent with each other and are also consistent with the results presented above using the standard fixed-momentum analysis technique. This increases the statistical impact of our results.

We also plot previous $N_f = 2$ results [27, 15, 28] that are in the quark mass and lattice spacing region we are exploring in fig. 5(b). Different groups using different lattice formulations obtain consistent results. Results from our two lattice spacings are consistent with each other and we are able to evaluate the flavour singlet mass closer to the continuum limit (and in an order a^2 improved formalism) than hitherto.

An extrapolation to the chiral limit of our results gives a value of the η_2 mass of $r_0 m(\eta_2) = 1.99(15)$ assuming a linear dependence. We actually see no statistically significant evidence for any slope, and if we assume a constant behaviour, we would obtain $r_0 m(\eta_2) = 2.00(5)$. Using $r_0 = 0.454(7)$ fm obtained [7] from the ETMC evaluation of f_π , we obtain an η_2 mass of 0.865(65) GeV. This updates the phenomenological estimate of 0.776 GeV discussed above, which was in turn




Figure 5: Comparison of values for the η_2 mass versus quark mass in units of $r_0 \approx 0.45$ fm. Values of the lattice spacing used are shown by listing r_0/a . Some of the points have been displaced horizontally for legibility. The ETMC results shown in (a) are those from the 2-state fits to zero-momentum correlators (first method in text, open symbols) and from the error-reduction obtained by using point-to-point correlators and projecting to the ground state of the neutral pseudoscalar connected contribution (filled symbols). In (b) we compare our results with reduced errors (filled symbols) to other $N_f = 2$ results from other collaborations [15, 27, 28] (open symbols) with relatively light pions and small lattice spacing ($a < 0.1$ fm, $r_0/a > 4.5$). The strange quark corresponds to $(r_0 m_\pi)^2 \approx 2.5$.

mainly motivated by lattice results at heavier quark masses than those we now have available. We have additional systematic errors which are not fully under control: from the chiral extrapolation, the continuum extrapolation and any possible volume dependence. We do not see any sign that these systematic errors are required to our current statistical precision, but, as a conservative estimate, we consider these systematic errors to be comparable to the statistical error. So we quote an η_2 mass of $0.865(65)(65)$ GeV.

One of the main conclusions of our study is that we find strong evidence that the flavour singlet mass remains finite as the quark mass is reduced to the chiral limit. This has implications for the topological charge: it implies that the topological charge susceptibility must decrease as m_π^2 in the chiral limit [29]. We will discuss our results for topological charge more fully elsewhere [1].

Acknowledgments

We thank Jaume Carbonell and Mariane Brinet for assistance with making point source propagators available for testing point to point propagation. We thank all members of ETMC for useful comments, help and a fruitful collaboration. One of the authors (CM) acknowledges a useful discussion with Paul Rakow.

The computer time for this project was made available to us by the John von Neumann-Institute for Computing on the JUMP and Jubl systems in Jülich and apeNEXT system in Zeuthen, by UKQCD on the QCDOC machine at Edinburgh, by INFN on the apeNEXT systems in Rome, by BSC on MareNostrum in Barcelona (www.bsc.es), by the NW Development cluster at Liverpool and by the Leibniz Computer centre in Munich on the Altix system. We thank these computer centres and their staff for all technical advice and help. On QCDOC we have made use of Chroma [30] and BAGEL [31] software and we thank members of UKQCD for assistance.

This work has been supported in part by the DFG Sonderforschungsbereich/Transregio SFB/TR9-03 and the EU Integrated Infrastructure Initiative Hadron Physics (I3HP) under contract RII3-CT-2004-506078. We also thank the DEISA Consortium (co-funded by the EU, FP6 project 508830), for support within the DEISA Extreme Computing Initiative (www.deisa.org).

References

- [1] ETMC, in preparation (2007).
- [2] T. Feldmann, *Int. J. Mod. Phys.* **A15**, 159 (2000), [[hep-ph/9907491](http://arxiv.org/abs/hep-ph/9907491)].
- [3] UKQCD, C. McNeile and C. Michael, *Phys. Lett.* **B491**, 123 (2000), [[hep-lat/0006020](http://arxiv.org/abs/hep-lat/0006020)].
- [4] P. Weisz, *Nucl. Phys.* **B212**, 1 (1983).
- [5] ALPHA, R. Frezzotti, P. A. Grassi, S. Sint and P. Weisz, *JHEP* **08**, 058 (2001), [[hep-lat/0101001](http://arxiv.org/abs/hep-lat/0101001)].
- [6] R. Frezzotti and G. C. Rossi, *JHEP* **08**, 007 (2004), [[hep-lat/0306014](http://arxiv.org/abs/hep-lat/0306014)].
- [7] ETMC, P. Boucaud *et al.*, *Phys. Lett.* **B650**, 304 (2007), [[hep-lat/0701012](http://arxiv.org/abs/hep-lat/0701012)].
- [8] $\chi_{\mathbb{F}}^{\mathbb{F}}$, K. Jansen, A. Shindler, C. Urbach and I. Wetzorke, *Phys. Lett.* **B586**, 432 (2004), [[hep-lat/0312013](http://arxiv.org/abs/hep-lat/0312013)].
- [9] $\chi_{\mathbb{F}}^{\mathbb{F}}$, K. Jansen, M. Papinutto, A. Shindler, C. Urbach and I. Wetzorke, *JHEP* **09**, 071 (2005), [[hep-lat/0507010](http://arxiv.org/abs/hep-lat/0507010)].
- [10] ETMC, C. Urbach, *PoS LAT2007*, 022 (2006), [[0710.1517](http://arxiv.org/abs/0710.1517)].
- [11] ETMC, P. Dimopoulos, R. Frezzotti, G. Herdoiza, C. Urbach and U. Wenger, *PoS LAT2007*, 102 (2007), [[0710.2498](http://arxiv.org/abs/0710.2498)].
- [12] ETMC, B. Blossier *et al.*, *JHEP* **04**, 020 (2008), [[0709.4574](http://arxiv.org/abs/0709.4574)].

- [13] ETMC, C. Alexandrou *et al.*, arXiv:0803.3190 [hep-lat].
- [14] $\chi_{\text{F}}^{\text{F}}$, K. Jansen *et al.*, Phys. Lett. **B624**, 334 (2005), [hep-lat/0507032].
- [15] UKQCD, C. R. Allton *et al.*, Phys. Rev. **D70**, 014501 (2004), [hep-lat/0403007].
- [16] A. Shindler, arXiv:0707.4093 [hep-lat].
- [17] ETMC, C. Michael and C. Urbach, PoS **LAT2007**, 122 (2007), [0709.4564].
- [18] ETMC, P. Boucaud *et al.*, arXiv:0803.0224 [hep-lat].
- [19] UKQCD, M. Foster and C. Michael, Phys. Rev. **D59**, 074503 (1999), [hep-lat/9810021].
- [20] UKQCD, C. McNeile and C. Michael, Phys. Rev. **D73**, 074506 (2006), [hep-lat/0603007].
- [21] UKQCD, P. Lacey, A. McKerrell, C. Michael, I. M. Stopher and P. W. Stephenson, Phys. Rev. **D51**, 6403 (1995), [hep-lat/9412079].
- [22] UKQCD, C. McNeile and C. Michael, Phys. Rev. **D63**, 114503 (2001), [hep-lat/0010019].
- [23] E. B. Gregory, A. C. Irving, C. M. Richards and C. McNeile, Phys. Rev. **D77**, 065019 (2008), [0709.4224].
- [24] H. Neff, N. Eicker, T. Lippert, J. W. Negele and K. Schilling, Phys. Rev. **D64**, 114509 (2001), [hep-lat/0106016].
- [25] C. DeTar and L. Levkova, PoS **LAT2007**, 116 (2007), [arXiv:0710.1322 [hep-lat]].
- [26] C. Michael and I. Teasdale, Nucl. Phys. **B215**, 433 (1983).
- [27] CP-PACS, V. I. Lesk *et al.*, Phys. Rev. **D67**, 074503 (2003), [hep-lat/0211040].
- [28] K. Hashimoto and T. Izubuchi, arXiv:0803.0186 [hep-lat].
- [29] H. Leutwyler and A. Smilga, Phys. Rev. **D46**, 5607 (1992).
- [30] SciDAC, R. G. Edwards and B. Joo, Nucl. Phys. Proc. Suppl. **140**, 832 (2005), [hep-lat/0409003].
- [31] P. Boyle, <http://www.ph.ed.ac.uk/~paboyle/bagel/Bagel.html>.