

Determination of Low-Energy Constants of Wilson Chiral Perturbation Theory



Gregorio Herdoíza ^{a,b}, Karl Jansen ^{c,d}, Chris Michael ^e,
Konstantin Ottnad ^f, Carsten Urbach ^f

^a*PRISMA Cluster of Excellence, Institut für Kernphysik
Johannes Gutenberg-Universität, D-55099 Mainz, Germany*

^b*Departamento de Física Teórica and Instituto de Física Teórica UAM/CSIC,
Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain*

^c*NIC, DESY, Platanenallee 6, D-15738 Zeuthen, Germany*

^d*Department of Physics, University of Cyprus
P.O.Box 20537, 1678 Nicosia, Cyprus*

^e*Theoretical Physics Division, Department of Mathematical Sciences
The University of Liverpool, Liverpool L69 3BX, UK*

^f*Helmholtz-Institut für Strahlen- und Kernphysik (Theorie) and Bethe Center for
Theoretical Physics, Universität Bonn, 53115 Bonn, Germany*

Abstract

By matching Wilson twisted mass lattice QCD determinations of pseudoscalar meson masses to Wilson Chiral Perturbation Theory we determine the low-energy constants W'_6 , W'_8 and their linear combination c_2 . We explore the dependence of these low-energy constants on the choice of the lattice action and on the number of dynamical flavours.

Key words: Lattice QCD, Wilson Fermions, Chiral Perturbation Theory.

PACS: 12.38.Gc, 12.39Fe

Preprint-No: DESY 13-043, FTUAM-13-127, IFT-UAM/CSIC-13-015,
MITP/13-015, SFB/CPP-13-18

1 Introduction

Lattice QCD simulations employing a discretisation of the Dirac operator based on the original proposal by Wilson [1] are currently being performed with light dynamical fermions [2–9]. When decreasing the light quark mass at a fixed value of the lattice spacing, a subtle interplay between mass and discretisation effects can take place due to the explicit breaking of chiral symmetry by the Wilson term. In simulations with light values of the quark mass it is, therefore, vital to understand and monitor the discretisation effects and to obtain a quantitative measure of their size.

Close to the continuum limit, a useful way to determine the discretisation effects in the regime of light quark masses is provided by Wilson chiral perturbation theory ($W\chi$ PT), an extension of the continuum chiral effective theory including additional terms proportional to powers of the lattice spacing [10]. Depending on the order of the expansion, additional low energy constants (LECs) appear, whose values are not known a priori: they depend on the lattice action and can only be computed from a simulation.

Knowing the values of the LECs of $W\chi$ PT is of particular interest, because $W\chi$ PT predicts a non-trivial phase structure for Wilson type fermions in the lattice spacing and quark mass plane [10–12]. Depending on the sign of a particular combination of Wilson LECs – commonly denoted as $c_2 \propto -(2W'_6 + W'_8)$ – either the Aoki-scenario [11] or a first order, so called Sharpe-Singleton [10] scenario is realised. Numerical evidence for both scenarios has been observed in lattice QCD simulations and dedicated studies of the associated phase diagrams have been performed by several groups [13–24].

The Aoki scenario with a positive c_2 was found to be realised in quenched simulations. In dynamical simulations the Sharpe-Singleton scenario with negative c_2 was observed when using Wilson twisted mass fermions at maximal twist [20–23]. This manifests itself in the fact that the neutral pion mass M_{π^0} is lighter than the charged one, M_{π^\pm} , where the splitting of the squared masses is proportional to c_2 . In turn, a measurement of the pion mass-splitting in Wilson twisted mass lattice QCD provides a way to measure c_2 and, hence, the LECs of the corresponding chiral effective theory. However, this way of computing c_2 is challenging since for the neutral pion mass disconnected contributions need to be evaluated.

There are also alternative ways to determine the LECs of $W\chi$ PT. In Refs. [25–28] they have been studied by matching the analytical predictions [27, 29–34] for the spectrum of the Wilson Dirac operator – with fixed index in a finite volume – to lattice data.¹ Determinations of the Wilson LECs have

¹ For a recent review, we refer to [35].

also been carried out via the spectral density of the Hermitian Wilson-Dirac operator [36–38]. Lattice determinations of the pion scattering lengths have been used in Refs. [39, 40], an approach that was extended to a partially quenched setup in Ref. [41]. In a mixed action with Wilson-type sea fermions and chirally invariant valence quarks, a mixed action chiral Lagrangian [42, 43] can be constructed. The corresponding LECs – in particular W'_8 – have been recently determined in the case of overlap valence quarks on a Wilson twisted mass sea [44].

In this paper we are going to determine the LECs using a method recently proposed in Ref. [45]. It relies on the measurement of pseudoscalar meson masses involving Wilson twisted mass fermions. In this approach, the LEC W'_8 is related to the mass-splitting between the charged pion mass M_{π^\pm} and the “connected neutral pion” mass $M_{\pi(0,c)}$. The latter is determined from the quark-connected correlation, which contributes to the neutral pion in twisted mass QCD. The “connected neutral pion” correlation function thus differs from the complete correlation function needed to determine the neutral pion by the absence of disconnected diagrams. This has the advantage that in numerical studies the “connected pion mass-splitting” should be accessible with good statistical precision. In addition, the mass-splitting, $M_{\pi(0,c)}^2 - M_{\pi^0}^2$, between the connected and full neutral pion mass, provides an estimate of W'_6 . We are also going to study the mass and lattice spacing dependence of these splittings arising at higher order in the $W\chi$ PT expansion.

The paper is structured as follows: in the next section we collect the $W\chi$ PT expressions relating the pion mass-splittings to the Wilson LECs. In Section 3 we present the lattice actions used in our study. The determination of the LECs from two of those lattice setups is described in Section 4. Finally, a qualitative comparison of the values of these LECs from different choices of the lattice action is reported in Section 5.

2 Wilson Chiral Perturbation Theory ($W\chi$ PT)

In this section, we briefly discuss Wilson chiral perturbation theory and introduce the expressions used in our study. For a recent review on the applications of χ PT to lattice QCD, we refer to Ref. [46].

Our study will be based on the computation of pseudoscalar meson masses involving Wilson twisted mass fermions. We are therefore interested in a chiral Lagrangian involving a mass matrix $M = m_0^R + i\mu_\ell^R \tau_3$, where m_0^R and μ_ℓ^R are the renormalised untwisted and twisted quark masses, respectively. The masses m_0 and μ_ℓ appear in the Wilson twisted mass action of eq. (12).

At leading order (LO) in the power counting, $m_0 \sim \mu_\ell \sim a^2 \Lambda_{\text{QCD}}^3$, and after a shift of the quark mass to remove a term of $\mathcal{O}(a)$, the partially quenched chiral Lagrangian reads [10, 43]

$$\begin{aligned} \mathcal{L}_\chi = & \frac{f^2}{8} \text{Str} \left(\partial_\mu \Sigma \partial_\mu \Sigma^\dagger \right) - \frac{f^2 B_0}{4} \text{Str} \left(M^\dagger \Sigma + \Sigma^\dagger M \right) \\ & - \hat{a}^2 W'_6 \left[\text{Str} \left(\Sigma + \Sigma^\dagger \right) \right]^2 - \hat{a}^2 W'_7 \left[\text{Str} \left(\Sigma - \Sigma^\dagger \right) \right]^2 \\ & - \hat{a}^2 W'_8 \text{Str} \left(\Sigma^2 + [\Sigma^\dagger]^2 \right), \end{aligned} \quad (1)$$

where Σ parametrises the vacuum manifold and thus characterises the Nambu-Goldstone bosons arising from the spontaneous breaking of chiral symmetry. The traces over the flavour indices are denoted by Str and $\hat{a} = 2W_0 a$. In addition to the continuum LECs B_0 and f (defined with the convention giving $f_\pi \approx 130$ MeV), the chiral Lagrangian also includes W_0 , W'_6 , W'_7 and W'_8 , which are Wilson LECs describing discretisation effects.

In this work we are interested in the determination of the Wilson LECs by matching lattice QCD calculations of pseudoscalar meson masses to their PQW χ PT expressions. As already mentioned, we consider Wilson twisted mass fermions at maximal twist. This is achieved in the chiral Lagrangian by setting $m_0 = 0$. At non vanishing values of the lattice spacing, the breaking of flavour symmetry by the twisted mass term in eq. (12) implies that the mass of the charged pion M_{π^\pm} differs from that of the neutral pion M_{π^0} by $\mathcal{O}(a^2)$ effects. A similar pattern holds for the mass of the ‘‘connected neutral pion’’ mass, $M_{\pi(0,c)}$. The PQW χ PT expressions for these three meson masses at LO read [45, 47–49],

$$M_{\pi^\pm}^2 = 2B_0 \mu_\ell, \quad (2)$$

$$M_{\pi^0}^2 = 2B_0 \mu_\ell - 8a^2 (2w'_6 + w'_8), \quad (3)$$

$$M_{\pi(0,c)}^2 = 2B_0 \mu_\ell - 8a^2 w'_8, \quad (4)$$

where w'_k is related to the Wilson LEC W'_k by

$$w'_k = \frac{16W_0^2 W'_k}{f^2} \quad (k = 6, 8). \quad (5)$$

To determine the individual values of the LECs it is, therefore, possible to consider the following mass-splittings:

$$M_{\pi^\pm}^2 - M_{\pi(0,c)}^2 = 8a^2 w'_8, \quad (6)$$

$$\frac{1}{2} \left(M_{\pi(0,c)}^2 - M_{\pi^0}^2 \right) = 8a^2 w'_6. \quad (7)$$

From eq. (3), it appears that the linear combination of LECs which controls the mass-splitting between charged and neutral pions is given by

$$c_2 = -\frac{32W_0^2}{f^2}(2W'_6 + W'_8). \quad (8)$$

This can also be re-expressed as

$$c_2 = \frac{1}{4a^2} (M_{\pi^0}^2 - M_{\pi^\pm}^2), \quad (9)$$

$$= -2(2w'_6 + w'_8). \quad (10)$$

In the $W\chi$ PT expressions presented above, two light mass-degenerate flavours were assumed to be present in the sea sector. When considering also other dynamical flavours, such as the strange and the charm quarks, the same expressions hold when assuming that these heavier flavours sufficiently decouple from the light quark sector. In this case, the values of the Wilson LECs will have a further residual dependence on the heavier quark masses.

Before closing this section, we mention that $W\chi$ PT calculations at NLO have been carried out for the pion mass and decay constant with $N_f = 2$ [50–52] and $N_f = 2 + 1 + 1$ [53] flavours of twisted mass fermions.

3 Lattice actions

The complete lattice action can be written as

$$S = S_f + S_g, \quad (11)$$

where S_f is the fermionic action and S_g is the pure gauge action. As we shall see, in this work we will consider a few alternatives for both the fermionic and the gauge actions in order to explore the dependence of the Wilson LECs on the details of the lattice action. As discussed below, we will use for S_F a few variants of Wilson twisted mass fermions.

3.1 Wilson Twisted Mass Fermions

The Wilson twisted mass (Wtm) lattice action for the mass degenerate light doublet (u, d) in the so-called twisted basis reads [54, 55],

$$S_l = a^4 \sum_x \{ \bar{\chi}_l(x) [D[U] + m_{0,l} + i\mu_\ell \gamma_5 \tau_3] \chi_l(x) \}, \quad (12)$$

where $m_{0,l}$ is the untwisted bare quark mass, μ_ℓ is the bare twisted light quark mass, τ_3 is the third Pauli matrix acting in flavour space and

$$D[U] = \frac{1}{2} \left[\gamma_\mu \left(\nabla_\mu + \nabla_\mu^* \right) - a \nabla_\mu^* \nabla_\mu \right], \quad (13)$$

is the massless Wilson-Dirac operator. ∇_μ and ∇_μ^* are the forward and backward gauge covariant difference operators, respectively. Twisted mass light fermions are said to be at maximal twist if the bare untwisted mass $m_{0,l}$ is tuned to its critical value, m_{crit} . The quark doublet $\chi_l = (\chi_u, \chi_d)$ in the twisted basis is related by a chiral rotation to the quark doublet in the physical basis

$$\psi_l^{\text{phys}} = e^{\frac{i}{2}\omega_l\gamma_5\tau_3} \chi_l, \quad \bar{\psi}_l^{\text{phys}} = \bar{\chi}_l e^{\frac{i}{2}\omega_l\gamma_5\tau_3}, \quad (14)$$

where ω_l is the twist angle.

The twisted mass parameter μ_ℓ provides an infrared regulator avoiding the presence of accidental zero-modes in the Wilson-Dirac operator. An important property of Wtm fermions is that at maximal twist physical observables are $\mathcal{O}(a)$ improved [55]. In numerical simulations, maximal twist is achieved by tuning the value of the hopping parameter $\kappa = 1/(2m_{0,l} + 8)$ to its critical value κ_{crit} by tuning the PCAC quark mass m_{PCAC} to zero. The expected $\mathcal{O}(a^2)$ scaling of physical observables when performing the continuum limit extrapolation has been confirmed in the quenched approximation [56–59] and with $N_f = 2$ [60–63] and $N_f = 2 + 1 + 1$ [7, 64, 65] dynamical quarks.

A peculiar lattice artifact can appear in observables made out of Wtm quarks due to the breaking of isospin and parity by the twisted mass term in eq. (12). This effect, which is expected to vanish in physical quantities at a rate of $\mathcal{O}(a^2)$ when approaching the continuum limit, has been observed to be numerically small in most of the observables which have been analysed [7, 60–64]

An exception to this observed small isospin breaking effects is found in the case of the neutral pseudoscalar mass. Indeed, isospin breaking induces a mass-splitting between charged and neutral pion masses. Dedicated numerical studies indicate that while in the charged pion mass only very mild cutoff effects are present, the neutral pion mass is instead affected by significant $\mathcal{O}(a^2)$ effects [7, 60, 63, 65, 66]. An analysis based on the Symanzik expansion indicates that isospin breaking affects only a limited set of observables in a sizeable way, namely the neutral pion mass and kinematically related quantities [67, 68]. This analysis is complementary to that based on $W\chi\text{PT}$ where, as previously mentioned, the mass-splitting between charged and neutral pions is parametrised by the combination of LECs appearing in c_2 defined in eq. (8).

The determination of the neutral pion mass M_{π^0} involves both connected and disconnected contributions. The computation of quark-disconnected diagrams

is challenging and requires the employment of specific techniques in order to achieve a statistically significant determination of M_{π^0} [69, 70].

$N_f = 2 + 1 + 1$ Wtm fermions

In addition to a doublet of mass-degenerate light quarks (u, d), a heavier doublet with strange and charm quarks – (s, c) – can be incorporated in lattice QCD studied with Wilson twisted mass fermions [71, 72]. Also in this $N_f = 2 + 1 + 1$ setup, a tuning to maximal twist by imposing $m_{\text{PCAC}} = 0$, allows to achieve the automatic $\mathcal{O}(a)$ improvement of physical observables. We refer to Ref. [7] for a complete description of the lattice action for the heavier quark doublet and for further details on the lattice setup.

Mixed action with Osterwalder-Seiler valence quarks

Osterwalder-Seiler (OS) valence quarks [73] can be viewed as the building blocks of Wilson twisted mass fermions at maximal twist. The OS action for an *individual* quark flavour χ_f reads

$$S_f^{\text{OS}} = a^4 \sum_x \{ \bar{\chi}_f(x) [D[U] + m_{\text{crit}} + i\mu_f \gamma_5 r_f] \chi_f(x) \} , \quad (15)$$

where r_f (here $|r_f| = 1$) is the Wilson parameter and m_{crit} the critical mass. By combining two flavours of OS quarks with opposite signs of r_f , e.g. $r_2 = -r_1$, the action of a doublet of maximally Wtm fermions of mass $\mu_f = \mu_1 = \mu_2$ can be recovered. The benefits of the OS action are that $\mathcal{O}(a)$ improved physical observables [71] can be obtained by using the same estimates of m_{crit} as in the Wtm case and, thus, avoiding further tuning effort. OS and Wtm fermions coincide with Wilson fermions in the massless limit and consequently share the same renormalisation factors. This simplifies the matching of sea and valence quark masses in the context of a mixed-action with Wtm sea and OS valence quarks.

The pseudoscalar correlation function obtained when considering only the connected contribution to the neutral pion correlation function (*i.e.* when ignoring disconnected diagrams), is precisely the pion correlator with OS fermions. The mass-splittings in eqs.(6)-(7) can hence be interpreted as involving sea and valence quarks in a mixed action setup.

In this work, we aim at determining the Wilson LECs in a lattice theory with $N_f = 2 + 1 + 1$ Wtm fermions. A particular effort will be devoted to addressing the main systematic effects present in these determinations. We furthermore aim at exploring the qualitative change on the values of these LECs when varying the details of the lattice action. Below, we briefly summarise the alternative lattice setups used in this work. We will consider variants of the

action differing by the presence of smearing of the gauge links in the covariant derivative, the inclusion of the Sheikholeslami-Wohlert term or by a change in the number of dynamical flavours.

Stout Smearing.

A smearing procedure can be applied to the gauge links entering in the covariant derivatives in eq. (13). The stout smearing [74] procedure is analytic in the un-smear link variables and hence well suited for simulations with the HMC algorithm. The smearing can be iterated several times, with the price of extending the coupling of fermions to the gauge links over a larger region.

Sheikholeslami-Wohlert term.

In our comparison of the values of the mass-splittings in eqs. (6)-(7) from different lattice setups, we will also consider results available in the literature from quenched lattice simulations with Wtm quarks including the Sheikholeslami-Wohlert term [75].

3.2 Gauge action

The lattice gauge actions considered in this work have a generic form which includes a plaquette term $U_{x,\mu,\nu}^{1\times 1}$ and rectangular (1×2) Wilson loops $U_{x,\mu,\nu}^{1\times 2}$,

$$S_g = \frac{\beta}{3} \sum_x \left(b_0 \sum_{\substack{\mu,\nu=1 \\ 1 \leq \mu < \nu}}^4 \{1 - \text{Re Tr}(U_{x,\mu,\nu}^{1\times 1})\} + b_1 \sum_{\substack{\mu,\nu=1 \\ \mu \neq \nu}}^4 \{1 - \text{Re Tr}(U_{x,\mu,\nu}^{1\times 2})\} \right), \quad (16)$$

with $\beta = 6/g_0^2$ the bare inverse coupling and the normalisation condition $b_0 = 1 - 8b_1$. We will consider the case of the Wilson plaquette [1] action ($b_1 = 0$), the tree-level Symanzik improved [76, 77] action ($b_1 = -1/12$) and the Iwasaki [78–80] action ($b_1 = -0.331$). In Wtm simulations, the strength of the phase transition has been found [21, 22] to depend on the value of the parameter b_1 in eq. (16).

4 Numerical Studies

4.1 $N_f = 2 + 1 + 1$ Wtm fermions with Iwasaki gauge action

The purpose of this study is to determine the mass-splittings in eqs. (6) and (7), which are directly related to the Wilson LECs W'_8 and W'_6 , respectively. The lattice action is composed of the Iwasaki gauge action and $N_f = 2 + 1 + 1$ flavours of Wilson twisted mass fermions.

The simulations [7] were performed at three values of the lattice gauge coupling, $\beta = 1.90$, 1.95 and $\beta = 2.10$, corresponding to values of the lattice spacing $a \approx 0.09$ fm, 0.08 fm and 0.06 fm, respectively. The charged pion mass M_{π^\pm} approximately ranges from 230 MeV to 510 MeV. Simulated volumes correspond to values of $M_{\pi^\pm}L$ larger than 3.3. Physical spatial volumes range from $(1.9 \text{ fm})^3$ to $(2.8 \text{ fm})^3$.

The values of the pseudoscalar meson masses for the $N_f = 2 + 1 + 1$ ensembles are collected in Table 1. The mass-splittings in eqs. (6)-(7), which are directly proportional to W'_8 and W'_6 respectively, are illustrated in Fig. 1. We observe that the conditions $W'_8 < 0$ and $W'_6 > 0$ are fulfilled by the lattice data, in agreement with the bounds derived in Refs. [29, 34, 45]. In order to quote the values of the Wilson LECs $W'_{6,8}$, the systematic effects from quark-mass dependence, residual lattice artifacts and finite volume effects have to be addressed.

In the ‘‘large cut-off effects’’ power counting and at LO in the $W\chi$ PT chiral Lagrangian, the mass-splittings in eqs. (6)-(7) are expected to be independent of the lattice spacing and the light-quark mass. The possible presence of such effects might thus signal effects entering at higher orders in the $W\chi$ PT chiral expansion. Since the NLO expressions for these mass-splittings is currently not available in the literature, we rely in our systematic error analysis on a separate study of (i) the continuum-limit of the mass-splittings at a reference mass and (ii) the comparison of a constant and a linear chiral extrapolation in $M_{\pi^\pm}^2$.

Starting with point (i), we show in Fig. 1 the mass splittings eq. (6) and (7) relevant for W'_8 and W'_6 , respectively, as a function of $M_{\pi^\pm}^2$. We first observe that data points with similar values of $(M_{\pi^\pm}r_0)^2$, but coming from different lattice spacings, tend to be compatible with each other, in particular when considering the larger lattice sizes, represented by the filled symbols. The lattice spacing dependence of the two aforementioned mass splittings at a reference mass $(M_{\pi^\pm}r_0)^2 \approx 0.55$ is illustrated in Fig.2(a). Note that only the largest lattice sizes L are considered in this figure. Although the lattice size slightly varies when changing β , the lattice data fulfils $L \gtrsim 2.5$ fm and $M_{\pi^\pm}L \gtrsim$

Ens.	β	L/a	$a\mu_\ell$	aM_{π^\pm}	$aM_{\pi(0,c)}$	aM_{π^0}	r_0/a
A30.32	1.90	32	0.0030	0.1234(03)	0.2111(33)	0.0611(036)	5.23(4)
A40.32			0.0040	0.1415(04)	0.2274(31)	0.0811(050)	
A40.24		24	0.0040	0.1445(06)	0.2375(25)	0.0694(065)	
A60.24			0.0060	0.1727(06)	0.2544(26)	0.1009(113)	
A80.24			0.0080	0.1987(06)	0.2659(25)	0.1222(157)	
A100.24			0.0100	0.2215(04)	0.2883(14)	0.1570(178)	
A80.24s	1.90	24	0.0080	0.1982(04)	0.2649(16)	0.1512(115)	
A100.24s			0.0100	0.2215(04)	0.2841(16)	0.1863(141)	
B25.32	1.95	32	0.0025	0.1064(07)	0.1836(21)	0.0605(036)	5.71(4)
B35.32			0.0035	0.1249(07)	0.1919(17)	0.0710(061)	
B55.32			0.0055	0.1540(04)	0.2177(19)	0.1323(080)	
B75.32			0.0075	0.1808(05)	0.2360(12)	0.1557(126)	
B85.24		24	0.0085	0.1931(08)	0.2480(11)	0.1879(180)	
D15.48	2.10	48	0.0015	0.0695(03)	0.1124(15)	0.0561(031)	7.46(6)
D20.48			0.0020	0.0797(05)	0.1170(16)	0.0651(042)	
D30.48			0.0030	0.0978(04)	0.1296(15)	0.0860(046)	
D45.32sc	2.10	32	0.0045	0.1198(05)	0.1480(09)	0.0886(095)	

Table 1

Determination of the charged M_{π^\pm} , neutral connected $M_{\pi(0,c)}$ and neutral M_{π^0} pseudoscalar meson masses from simulations with $N_f = 2 + 1 + 1$ flavours of Wilson twisted mass fermions at maximal twist and the Iwasaki gauge action. The value of the Sommer scale r_0 is determined in the chiral limit. The ensemble names containing an “s” or a “c” refer to a change of the strange or the charm quark mass, respectively. The values for M_{π^\pm} agree within errors with the once published already in Ref. [7].

4 and, therefore, we do not expect a large effect from a small mismatch in the physical volume. Fig.2(a) suggests that the residual lattice spacing effects are small. We remind that these lattice artifacts appear beyond the leading $O(a^2)$ effects. As an estimate of these effects, we include in our systematic error analysis the difference between the values of the mass-splittings from the two finer lattice spacings.

Concerning the light-quark mass dependence in point (ii), we can expect that the mass terms appearing at NLO can contain a linear term in $M_{\pi^\pm}^2$ but




Fig. 1. Determination of the mass-splittings (a) $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ in eq. (6) and (b) $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ in eq. (7) as a function of $M_{\pi^\pm}^2$. These quantities are in units of the chirally extrapolated Sommer scale r_0 . The mass-splittings are directly related to the Wilson LECs W'_8 and W'_6 . The lattice setup with the Iwasaki gauge action and $N_f = 2 + 1 + 1$ flavours of Wilson twisted mass fermions is considered. Filled and empty symbols signal a change in the lattice size. The label “(s,c)” in the legend indicates the effect of changing the values of the strange and charm quark masses in the sea. The mass-splittings illustrated in these figures assess the size of $O(a^2)$ discretisation effects. Hence, any lattice spacing dependence in these quantities points to residual higher order discretisation effects. The results of a chiral extrapolation from a constant and a linear fit in $M_{\pi^\pm}^2$ are shown. The deviation between the two extrapolated values is included in the systematic error analysis.

also a term of the form $M_{\pi^\pm}^2 \log(M_{\pi^\pm}^2)$. Indeed, such terms are present in the $W\chi$ PT expression of the mass splitting between the charged and neutral pion masses at NLO [51]. Since the precise form of these logarithmic terms is yet unknown for the mass-splittings considered here, we limit ourselves to a linear chiral extrapolation in $M_{\pi^\pm}^2$. Note that our data is not precise enough to disentangle possible logarithmic contributions. We take as our central values the linearly extrapolated mass-splittings and use the difference with respect to the constant fit as an estimate of the systematic error from the light-quark mass dependence.

Finite volume effects are taken into account by adding to the systematic error the difference between the values of the mass splittings from two ensembles – A40.24 and A40.32 – differing only by a change of lattice size from $L \approx 2.1$ fm to 2.8 fm. This is expected to be a conservative choice since these ensembles correspond to a rather small light-quark mass – and therefore finite size effects can be non-negligible. Also, these ensembles were obtained at the coarsest lattice spacing, where possible finite size effects (FSE) from the neutral pion mass should be larger.




Fig. 2. (a) Lattice spacing dependence of the mass-splittings $|M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2|/a^2$ – empty symbols – and $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ – filled symbols – at a reference mass $(M_{\pi^\pm} r_0)^2 \approx 0.55$. For a better visibility, the absolute value of the mass splitting is considered in the case of the empty symbols. (b) Pion mass-splitting, $(M_{\pi^0}^2 - M_{\pi^\pm}^2)/a^2$, normalised according to eq. (9) in order to relate it to the combination of Wilson LECs c_2 . The chiral extrapolation using a constant and a linear fit in $M_{\pi^\pm}^2$ is also shown. The lattice setup with the Iwasaki gauge action and $N_f = 2 + 1 + 1$ flavours of Wilson twisted mass fermions is considered.

As already mentioned, the determination of the Wilson LECs $W'_{6,8}$ from lattice data with $N_f = 2 + 1 + 1$ flavours assumes that the strange and the charm sea-quarks decouple sufficiently from the light-quark dynamics. The residual heavy quark mass dependence present in $W'_{6,8}$ can be studied by varying the strange and charm quark masses in the neighbourhood of their physical values. This effect is illustrated in Fig. 1 by the points labelled by “(s,c)” in the legend. We use the difference between the mass-splittings from ensembles A80.24 and A80.24s – which only differ by a change of the strange and charm quark masses – to estimate this systematic effect. We expect that this is a conservative choice because (a) the ensemble A80.24s has a strange quark mass which is very close to the physical point, (b) the change in the strange quark mass is largest for A80.24 and A80.24s and (c) the effect of strange sea-quarks should be larger than that of charm quarks.

After combining the previously discussed systematic uncertainties in quadrature, we obtain the following values for the mass-splittings for the case of a lattice setup with $N_f = 2 + 1 + 1$ Wtm fermions and the Iwasaki gauge action,

	$w'_8 r_0^4$	w'_8	$W'_8 (r_0^6 W_0^2)$
syst.	-2.9(4)	-[571(32) MeV] ⁴	-0.0138(22)
	$w'_6 r_0^4$	w'_6	$W'_6 (r_0^6 W_0^2)$
syst.	+1.7(7)	+ [502(58) MeV] ⁴	+0.0082(34)
	$c_2 r_0^4$	c_2	$-2 (2W'_6 + W'_8) (r_0^6 W_0^2)$
lin.	-1.1(2)	-[444(28) MeV] ⁴	-0.0050(10)
cst.	-2.3(1)	-[541(24) MeV] ⁴	-0.0111(10)

Table 2

Determination of the Wilson LECs $W'_{6,8}$ ($w'_{6,8}$) and c_2 from a lattice setup with $N_f = 2 + 1 + 1$ Wtm fermions and the Iwasaki gauge action. For the values quoted in physical units, the input $r_0 = 0.45(2)$ fm has been used. The values in the last column derive from eq. (5), where the input $r_0 f = 0.276(12)$ from [7] has been used. For the case of $W'_{6,8}$ the systematic error analysis described in the text has been incorporated in the overall uncertainty indicated by the label “syst.” in the table. For c_2 , in the last block, we quote separately the results of a constant and a linear chiral extrapolation in $M_{\pi^\pm}^2$ – as illustrated in Fig.2(b) – and the quoted errors are purely statistical.

$$\left(\frac{M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2}{a^2} \right) r_0^4 = -23.0 \pm 0.7 \pm 3.0, \quad (17)$$

$$\left(\frac{M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2}{2a^2} \right) r_0^4 = +13.8 \pm 0.6 \pm 5.6, \quad (18)$$

where the first error is statistical and the second systematic. The corresponding values of the Wilson LECs are collected in Tab. 2. As already anticipated, the results for $w'_{6,8}$ are precise enough to identify a definite sign for these LECs.

The combination of LECs c_2 , can be determined directly from the mass-splitting $M_{\pi^0}^2 - M_{\pi^\pm}^2$ as indicated in eq. (9). The measurements of this mass-splitting are illustrated in Fig. 2(b) where the results of a chiral extrapolation by using a constant and a linear fit in $M_{\pi^\pm}^2$ are also shown. For the case of c_2 , we provide in Tab. 2 the results from both these chiral extrapolations and quote in the individual numbers only the statistical error. The values arising from these extrapolations are both compatible with a negative sign of c_2 .

The $W\chi$ PT expressions at NLO relevant for c_2 have been derived in Ref [51]. In addition to the Wilson LECs appearing at LO and to the usual Gasser-Leutwyler LECs, other parameters also appear at NLO. A complete determination of these parameters is beyond the scope of this study. We postpone such an analysis to a future dedicated study of the $W\chi$ PT description of both the

pion mass and decay constant. First results for the Gasser-Leutwyler LECs, from fits based on continuum χ PT, have been presented in Refs. [7, 81].

The connected neutral pion mass, $M_{\pi(0,c)}$, is an important ingredient in order to isolate the individual values of the LECs $W'_{6,8}$. As already pointed out, the connected neutral pion can be interpreted as the pion of a mixed action with OS fermions. Such a mixed action has been used to determine observables in the Kaon sector [65, 82, 83]. We note that extensions of the analytical expressions to SU(3) $W\chi$ PT is currently not available in the literature. Contrary to the pion case, the absence of disconnected diagrams in correlations functions in the Kaon sector could possibly allow to consider quantities from which the Wilson LECs can be determined with good accuracy.

4.2 $N_f = 2$ Wtm fermions with tlSym gauge action

In this section, we again determine the LECs $W'_{6,8}$, but this time using $N_f = 2$ flavours of Wilson twisted mass fermions and the tree-level Symanzik improved gauge action [3, 63, 70]. We already anticipate that a smaller set of ensembles and of measurements of the relevant pion masses are available in this case, in comparison to the $N_f = 2 + 1 + 1$ case discussed previously. Therefore, the resulting determinations and comparisons might suffer from insufficient control of systematic effects. However, we think that already a qualitative comparison can provide useful information to parametrise the size of cutoff effects from different lattice setups.

The simulations considered in this work [63, 85] were performed at three values of the lattice gauge coupling $\beta = 3.90$, 4.05 and $\beta = 4.20$, corresponding to values of the lattice spacing $a \approx 0.08$ fm, 0.07 fm and 0.05 fm, respectively. The charged pion mass M_{π^\pm} approximately ranges from 310 MeV to 460 MeV. Physical spatial volumes range from $(1.3 \text{ fm})^3$ to $(2.6 \text{ fm})^3$ and ensembles which differ only by the lattice size have been considered in order to address the size of finite volume effects in the determination of the Wilson LECs.

The values of the pseudoscalar meson masses [63] for the $N_f = 2$ ensembles are collected in Table 3. The mass-splittings in eqs. (6)-(7) are illustrated in Fig. 3. In order to explore the systematic effects present in these determinations, we follow a similar path to that described for the case of $N_f = 2 + 1 + 1$ ensembles. The availability of ensembles differing only by the physical volume allows to address the size of FSE in the mass-splittings. This is illustrated in Fig. 4(a). At $\beta = 3.90$, a set of three ensembles with $L/a = 16$, 24 and 32 could be used for the case of $(M_{\pi^\pm}^2 - M_{\pi(0,c)}^2)/a^2$. With the current statistical uncertainties, no clear signs of FSE can be observed in the data. Furthermore, data from two different lattice spacings – $\beta = 3.90$ and 4.05 – agree within errors, indicating

β	L/a	$a\mu_\ell$	aM_{π^\pm}	$aM_{\pi^{(0,c)}}$	aM_{π^0}	r_0/a
3.90	32	0.0040	0.1338(02)	0.2080(30)	0.1100(080)	5.35(4)
	24	0.0040	0.1362(07)	0.2120(30)	0.1090(070)	
	16	0.0040	0.1596(30)	0.2226(95)	-	
	24	0.0064	0.1694(04)	-	0.1340(100)	
		0.0085	0.1940(05)	-	0.1690(110)	
16	0.0074	0.1963(17)	0.2541(55)	-		
4.05	32	0.0030	0.1038(06)	0.1500(30)	0.0900(060)	6.71(4)
	20	0.0030	0.1191(41)	0.1571(62)	-	
	32	0.0060	0.1432(06)	0.1800(20)	0.1230(060)	
4.20	24	0.0020	0.0941(31)	0.1157(61)	-	8.36(6)

Table 3

Determination of the charged M_{π^\pm} , neutral connected $M_{\pi^{(0,c)}}$ and neutral M_{π^0} pseudoscalar meson masses from simulations with $N_f = 2$ flavours of Wilson twisted mass fermions at maximal twist and the tree-level Symanzik improved gauge action [63]. The value of the Sommer scale r_0 is determined in the chiral limit [84].




Fig. 3. Determination of the mass-splittings (a) $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ in eq. (6) and (b) $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ in eq. (7) as a function of $M_{\pi^\pm}^2$. These quantities are in units of the chirally extrapolated Sommer scale r_0 . The mass-splittings are directly related to the Wilson LECs W'_8 and W'_6 . The lattice setup with the tree-level Symanzik improved (tlSym) gauge action and $N_f = 2$ flavours of Wilson twisted mass fermions is considered. Filled and empty symbols signal a change in the lattice size.




Fig. 4. (a) Finite volume effects on the mass-splittings $|M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2|/a^2$ – empty symbols – and $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ – filled symbols. Note that the absolute value of the mass-splitting is used in the case of empty symbols. (b) Pion mass-splitting, $(M_{\pi^0}^2 - M_{\pi^\pm}^2)/a^2$, normalised according to eq. (9) in order to relate it to the combination of Wilson LECs c_2 . The chiral extrapolation using a constant and a linear fit in $M_{\pi^\pm}^2$ is also shown. The lattice setup with the tree-level Symanzik improved (tlSym) gauge action and $N_f = 2$ flavours of Wilson twisted mass fermions is considered.

the absence of large residual lattice artifacts in these mass-splittings. However, the lack of sufficient data does not allow to address the mass dependence of the mass-splitting $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/a^2$ in a satisfactory way. In analogy to the $N_f = 2 + 1 + 1$ case, we include the deviation between a constant and a linear extrapolation in $M_{\pi^\pm}^2$ in the estimate of the systematic uncertainties. The central value is taken from the result of the linear fit.

For the case of a lattice setup with $N_f = 2$ Wtm fermions at maximal twist and the tlSym gauge action, we obtain the following values for the mass-splittings

$$\left(\frac{M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2}{a^2}\right) r_0^4 = -20.1 \pm 2.3 \pm 1.7, \quad (19)$$

$$\left(\frac{M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2}{2a^2}\right) r_0^4 = +8.4 \pm 3.3 \pm 5.5, \quad (20)$$

where the first error is statistical and the second systematic. The corresponding values of the Wilson LECs are collected in Tab. 4. The LEC W'_8 has recently been determined from a mixed action involving the same $N_f = 2$ lattice action in the sea sector as that described here, but with Neuberger overlap valence quark [44]. The value quoted in Ref. [44], $W'_8(r_0^6 W_0^2) = -0.0064(24)$, differs from the estimate in Tab. 4 at the 2-sigma level. However, we stress once more

$w'_8 r_0^4$	w'_8	$W'_8 (r_0^6 W_0^2)$
-2.5(4)	-[552(025) MeV] ⁴	-0.0119(17)
$w'_6 r_0^4$	w'_6	$W'_6 (r_0^6 W_0^2)$
+1.0(8)	+ [443(138) MeV] ⁴	+0.0049(38)

Table 4

Determination of the Wilson LECs $W'_{6,8}$ ($w'_{6,8}$) from a lattice setup with $N_f = 2$ Wtm fermions and the tlSym gauge action. For the values quoted in physical units, the input $r_0 = 0.45(2)$ fm has been used. The values in the last column derive from eq. (5), where the input $r_0 f = 0.275(6)$ from [63] has been used.

that our present $N_f = 2$ estimate does not include a complete assessment of the systematic errors.

The determination of c_2 from $N_f = 2$ ensembles is illustrated in Fig.4(b). More data would be needed to isolate the residual mass-dependence present in c_2 . For this reason we opt for quoting separately the results of a constant and a linear chiral extrapolation in $M_{\pi^\pm}^2$,

$$c_2 r_0^4 [\text{cst.}] = -3.1 \pm 0.4, \quad (21)$$

$$c_2 r_0^4 [\text{lin.}] = -0.5 \pm 1.5, \quad (22)$$

where the errors are statistical only. These values are consistent with those arising from the measurements of pseudoscalar meson masses in Ref. [63] and are also very similar to the result obtained in Ref. [50] for $K = -4c_2$ from twisted mass finite volume effects. All the lattice measurements favour a negative sign of c_2 . However, as already mentioned, more data would be needed to properly address the residual mass dependence.

We refer to Refs. [63, 84] for more details about the description of the pion mass and decay constant by means of χ PT expressions including discretisation effects.

5 Discussion

In this section we collect a few comments concerning the extraction of the Wilson LECs. The LECs W'_6 and c_2 depend on the neutral pion mass M_{π^0} . The statistical error in M_{π^0} is dominated by the contribution of disconnected

diagrams.² We observe that the precise form of the light-quark mass dependence of the mass-splittings related to W'_6 and c_2 – see e.g. Figs. 1(b) and 2(b) – cannot be addressed within the present uncertainties. As previously discussed this mass dependence can arise at NLO in the $W\chi$ PT expansion. We can, however, not exclude at this stage that this higher-order effects are negligible. This issue is particularly relevant for the case of $c_2 \propto -(2w'_6 + w'_8)$, where a partial cancellation of the effect of w'_6 and w'_8 is present. The presence of higher order effects in the determination of c_2 has also been discussed in Ref. [40]. We recall that the sign of c_2 controls the appearance of an Aoki ($c_2 > 0$) or of a Sharpe-Singleton ($c_2 < 0$) scenario for the phase structure of Wilson fermions.

From the previously discussed determinations, a comparison of the values of the Wilson LECs from the lattice actions (i) $N_f = 2 + 1 + 1$ flavours of Wtm fermions at maximal twist and Iwasaki gauge action and (ii) $N_f = 2$ flavours of Wtm quarks and tISym gauge action, suggests that $W'_{6,8}$ and c_2 do not vary significantly in between these two setups. Let us extend this observation by performing a comparison of the pion mass-splittings as determined from different lattice actions. We stress that a complete assessment of the overall uncertainty is not available for most of these measurements and therefore the comparison remains at the qualitative level.

Lattice simulations with $N_f = 2 + 1 + 1$ Wtm fermions and the Iwasaki gauge action, but including in addition one iteration of stout smearing – labelled 1-stout – have been reported in Ref. [7]. A qualitative comparison of the effect of the stout smearing on the size of the mass-splittings is shown in Fig. 5. Note that a single ensemble is used in the estimate of the mass-splittings for the case of stout smearing. The used smearing seems to help in reducing the magnitude of the splitting $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ while – given the current uncertainties – it does not introduce a significant change in $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/a^2$.

Fig. 5(a) also includes the estimates of the mass-splitting $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ as determined from quenched ensembles either with or without the presence of the Sheikholeslami-Wohlert term. Maximally twisted-mass fermions and the plaquette gauge action are used in both cases. The values of the mass-splitting are derived from studies available in the literature. For the case without the Sheikholeslami-Wohlert term, results are taken from Ref. [66]. We use the information from different lattice spacings and quark masses to estimate the size of the systematic effects in $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$. For the case in which the Sheikholeslami-Wohlert term is included – labelled c_{SW} – we follow Ref. [86],

² It is interesting to note that Table 1 indicates that the relative error on the neutral pion mass is roughly independent of the light-quark mass and that it decreases when increasing the volume. In practice, in the current simulation conditions, this implies that the measurements of M_{π^0} are statistically more precise for the ensembles with lighter quark masses.




Fig. 5. Comparison of the values of the mass-splittings (a) $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ and (b) $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ from different lattice setups (see text for details).

where the non-perturbative determination of c_{SW} was used.

For cases other than those involving the Sheikholeslami-Wohlert term or stout-smearing, Fig. 5(a) suggests that, with the current precision, the value of the mass-splitting $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ does not significantly depend on a simultaneous change of the number of flavours N_f and of the gauge action. Note however that we cannot exclude that a change only in N_f or only in the parameter b_1 of the gauge action, leads to a different conclusion. It would be very desirable, if further actions are investigated and the precision could be increased.

One important observation arising from the measurements in Ref. [86] is that the introduction of the Sheikholeslami-Wohlert term in quenched studies significantly reduces the size of $O(a^2)$ effects by lowering the value of the mass-splitting $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$. It would thus be interesting to study whether this result still holds for simulations with dynamical fermions and whether the value of $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ is also reduced in that case.

Conclusions

We have presented the determination of the Wilson LECs $W'_{6,8}$ and c_2 , parametrisng the size of $O(a^2)$ lattice artifacts in $W\chi\text{PT}$, from simulations with

a lattice action composed out of the Iwasaki gauge action and $N_f = 2 + 1 + 1$ flavours of Wilson twisted mass fermions at maximal twist. The values of $W'_{6,8}$ include a rather complete account of the systematic uncertainties. Our measurements satisfy the recently derived bounds [29, 34, 45], $W'_8 < 0$ and $W'_6 > 0$.

We have also explored the dependence of the mass-splittings $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$ and $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$ on the choice of the lattice action. From this qualitative comparison, it is tempting to conjecture that a lattice action made of dynamical twisted mass fermions including the Sheikholeslami-Wohlert and smearing might lead to a reduction of the mass-splitting $(M_{\pi^\pm}^2 - M_{\pi^{(0,c)}}^2)/a^2$. Further studies are needed to clarify this point and to extend it to the case of $(M_{\pi^{(0,c)}}^2 - M_{\pi^0}^2)/2a^2$.

A partial cancellation of the contributions from W'_6 and W'_8 implies that the residual mass-dependence of c_2 is more sensitive to higher order effects in the $W\chi$ PT expansion. While this potential reduction of c_2 would certainly be beneficial, the precise determination of its value cannot be achieved with the currently available data.

The determination of the Wilson LECs can help to quantify the size of $O(a^2)$ terms in a given lattice action. Knowing these LECs, in particular with better precision, can significantly contribute to design a lattice fermion action with small lattice artifacts, thus allowing to reach the continuum limit in a better controlled way. In addition, an independent calculation of the Wilson LECs, as carried through here, can be used in chiral perturbation theory fits of light meson observables by constraining these fits to lattice data. Thus we think that the study of the (connected and full) neutral and charged pion masses of this work can be beneficial for many other groups working with Wilson-like lattice fermions.

Acknowledgements

We thank Giancarlo Rossi for useful comments on the manuscript. The computer time for this project was made available to us by the John von Neumann-Institute for Computing (NIC) on the JUDGE and Jugene systems in Jülich and the IDRIS (CNRS) computing center in Orsay. In particular we thank U.-G. Meißner for granting us access on JUDGE. Falk Zimmermann cross-checked correlation functions for one of our ensembles, which we gratefully acknowledge. G. H. acknowledges the support by DFG (SFB 1044), the Spanish Ministry for Education and Science project FPA2009-09017, the Consolider-Ingenio 2010 Programme CPAN (CSD2007-00042), the Comunidad Autónoma de Madrid (HEPHACOS P-ESP-00346 and HEPHACOS S2009/ESP-1473)

and the European project STRONGnet (PITN-GA-2009-238353). K. J. was supported in part by the Cyprus Research Promotion Foundation under contract ΠΡΟΣΕΛΑΚΥΣΗ/ΕΜΠΕΙΡΟΣ/0311/16. This work has been supported in part by the DFG Sonderforschungsbereich/ Transregio SFB/TR9. Two of the authors (K. O. and C. U.) were supported by the Bonn-Cologne Graduate School (BCGS) of Physics and Astronomie. This project was supported in parts by the DFG in SFB/TR16 and CRC 110.

References

- [1] K. G. Wilson, Phys. Rev. **D10**, 2445 (1974).
- [2] L. Del Debbio, L. Giusti, M. Lüscher, R. Petronzio and N. Tantalo, JHEP **02**, 056 (2007), [arXiv:hep-lat/0610059](#).
- [3] **ETM** Collaboration, P. Boucaud *et al.*, Phys. Lett. **B650**, 304 (2007), [arXiv:hep-lat/0701012](#).
- [4] **Hadron Spectrum** Collaboration, H.-W. Lin *et al.*, Phys.Rev. **D79**, 034502 (2009), [arXiv:0810.3588 \[hep-lat\]](#).
- [5] **BMW** Collaboration, S. Dürr *et al.*, Science **322**, 1224 (2008), [arXiv:0906.3599 \[hep-lat\]](#).
- [6] **PACS-CS** Collaboration, S. Aoki *et al.*, Phys.Rev. **D81**, 074503 (2010), [arXiv:0911.2561 \[hep-lat\]](#).
- [7] **ETM** Collaboration, R. Baron *et al.*, JHEP **1006**, 111 (2010), [arXiv:1004.5284 \[hep-lat\]](#).
- [8] **BMW** Collaboration, S. Dürr *et al.*, JHEP **1108**, 148 (2011), [arXiv:1011.2711 \[hep-lat\]](#).
- [9] **QCDSF-UKQCD** Collaboration, W. Bietenholz *et al.*, Phys.Rev. **D84**, 054509 (2011), [arXiv:1102.5300 \[hep-lat\]](#).
- [10] S. R. Sharpe and J. Singleton, R., Phys. Rev. **D58**, 074501 (1998), [hep-lat/9804028](#).
- [11] S. Aoki, Phys.Rev. **D30**, 2653 (1984).
- [12] M. Creutz, [arXiv:hep-lat/9608024 \[hep-lat\]](#).
- [13] S. Aoki and A. Gocksch, Phys.Rev. **D45**, 3845 (1992).
- [14] T. Blum *et al.*, Phys. Rev. **D50**, 3377 (1994), [hep-lat/9404006](#).
- [15] S. Aoki, Prog. Theor. Phys. Suppl. **122**, 179 (1996), [hep-lat/9509008](#).
- [16] S. Aoki, T. Kaneda and A. Ukawa, Phys.Rev. **D56**, 1808 (1997), [arXiv:hep-lat/9612019 \[hep-lat\]](#).

- [17] **JLQCD** Collaboration, S. Aoki *et al.*, Nucl. Phys. Proc. Suppl. **106**, 263 (2002), [hep-lat/0110088](#).
- [18] E.-M. Ilgenfritz, W. Kerler, M. Müller-Preußker, A. Sternbeck and H. Stüben, Phys. Rev. **D69**, 074511 (2004), [hep-lat/0309057](#).
- [19] **JLQCD** Collaboration, S. Aoki *et al.*, Phys.Rev. **D72**, 054510 (2005), [arXiv:hep-lat/0409016](#) [[hep-lat](#)].
- [20] F. Farchioni *et al.*, Eur. Phys. J. **C39**, 421 (2005), [hep-lat/0406039](#).
- [21] F. Farchioni *et al.*, Eur. Phys. J. **C42**, 73 (2005), [hep-lat/0410031](#).
- [22] F. Farchioni *et al.*, Phys. Lett. **B624**, 324 (2005), [hep-lat/0506025](#).
- [23] F. Farchioni *et al.*, Eur. Phys. J. **C47**, 453 (2006), [hep-lat/0512017](#).
- [24] T. Chiarappa *et al.*, Eur.Phys.J. **C50**, 373 (2007), [arXiv:hep-lat/0606011](#) [[hep-lat](#)].
- [25] A. Deuzeman, U. Wenger and J. Wuilloud, [arXiv:1110.4002](#) [[hep-lat](#)].
- [26] P. Damgaard, U. Heller and K. Splittorff, [arXiv:1110.2851](#) [[hep-lat](#)].
- [27] P. Damgaard, U. Heller and K. Splittorff, [arXiv:1206.4786](#) [[hep-lat](#)].
- [28] P. H. Damgaard, U. M. Heller and K. Splittorff, [arXiv:1301.3099](#) [[hep-lat](#)].
- [29] P. Damgaard, K. Splittorff and J. Verbaarschot, Phys.Rev.Lett. **105**, 162002 (2010), [arXiv:1001.2937](#) [[hep-th](#)].
- [30] G. Akemann, P. Damgaard, K. Splittorff and J. Verbaarschot, Phys.Rev. **D83**, 085014 (2011), [arXiv:1012.0752](#) [[hep-lat](#)].
- [31] K. Splittorff and J. Verbaarschot, Phys.Rev. **D84**, 065031 (2011), [arXiv:1105.6229](#) [[hep-lat](#)].
- [32] K. Splittorff and J. Verbaarschot, PoS **LATTICE2011**, 113 (2011), [arXiv:1112.0377](#) [[hep-lat](#)].
- [33] K. Splittorff and J. Verbaarschot, Phys.Rev. **D85**, 105008 (2012), [arXiv:1201.1361](#) [[hep-lat](#)].
- [34] M. Kieburg, K. Splittorff and J. Verbaarschot, Phys.Rev. **D85**, 094011 (2012), [arXiv:1202.0620](#) [[hep-lat](#)].
- [35] K. Splittorff, PoS **LATTICE2012**, 018 (2012), [arXiv:1211.1803](#) [[hep-lat](#)].
- [36] S. Necco and A. Shindler, JHEP **1104**, 031 (2011), [arXiv:1101.1778](#) [[hep-lat](#)].
- [37] S. Necco and A. Shindler, PoS **LATTICE2011**, 250 (2011), [arXiv:1108.1950](#) [[hep-lat](#)].
- [38] S. Necco and A. Shindler, [arXiv:1302.5595](#) [[hep-lat](#)].

- [39] S. Aoki, O. Bär and B. Biedermann, Phys.Rev. **D78**, 114501 (2008), arXiv:0806.4863 [hep-lat].
- [40] F. Bernardoni, J. Bulava and R. Sommer, PoS **LATTICE2011**, 095 (2011), arXiv:1111.4351 [hep-lat].
- [41] M. T. Hansen and S. R. Sharpe, Phys.Rev. **D85**, 054504 (2012), arXiv:1112.3998 [hep-lat].
- [42] O. Bär, G. Rupak and N. Shoresh, Phys. Rev. **D67**, 114505 (2003), arXiv:hep-lat/0210050.
- [43] O. Bär, G. Rupak and N. Shoresh, Phys. Rev. **D70**, 034508 (2004), arXiv:hep-lat/0306021.
- [44] K. Cichy, V. Drach, E. Garcia-Ramos, G. Herdoiza and K. Jansen, Nucl.Phys. **B869**, 131 (2013), arXiv:1211.1605 [hep-lat].
- [45] M. T. Hansen and S. R. Sharpe, Phys.Rev. **D85**, 014503 (2012), arXiv:1111.2404 [hep-lat].
- [46] M. Golterman, arXiv:0912.4042 [hep-lat].
- [47] G. Münster, JHEP **09**, 035 (2004), hep-lat/0407006.
- [48] L. Scorzato, Eur. Phys. J. **C37**, 445 (2004), hep-lat/0407023.
- [49] S. R. Sharpe and J. M. S. Wu, Phys. Rev. **D70**, 094029 (2004), hep-lat/0407025.
- [50] G. Colangelo, U. Wenger and J. M. Wu, Phys.Rev. **D82**, 034502 (2010), arXiv:1003.0847 [hep-lat].
- [51] O. Bär, Phys.Rev. **D82**, 094505 (2010), arXiv:1008.0784 [hep-lat].
- [52] S. Ueda and S. Aoki, arXiv:1109.0073 [hep-lat].
- [53] G. Münster and T. Sudmann, JHEP **1104**, 116 (2011), arXiv:1103.1494 [hep-lat].
- [54] **ALPHA** Collaboration, R. Frezzotti, P. A. Grassi, S. Sint and P. Weisz, JHEP **08**, 058 (2001), hep-lat/0101001.
- [55] R. Frezzotti and G. C. Rossi, JHEP **08**, 007 (2004), hep-lat/0306014.
- [56] $\chi_{\text{L}}^{\text{F}}$ Collaboration, K. Jansen, A. Shindler, C. Urbach and I. Wetzorke, Phys. Lett. **B586**, 432 (2004), hep-lat/0312013.
- [57] $\chi_{\text{L}}^{\text{F}}$ Collaboration, K. Jansen, M. Papinutto, A. Shindler, C. Urbach and I. Wetzorke, Phys. Lett. **B619**, 184 (2005), hep-lat/0503031.
- [58] $\chi_{\text{L}}^{\text{F}}$ Collaboration, K. Jansen, M. Papinutto, A. Shindler, C. Urbach and I. Wetzorke, JHEP **09**, 071 (2005), hep-lat/0507010.
- [59] A. M. Abdel-Rehim, R. Lewis and R. M. Woloshyn, Phys. Rev. **D71**, 094505 (2005), hep-lat/0503007.

- [60] **ETM** Collaboration, C. Urbach, PoS **LAT2007**, 022 (2007), [arXiv:0710.1517 \[hep-lat\]](#).
- [61] **ETM** Collaboration, P. Dimopoulos, R. Frezzotti, G. Herdoiza, C. Urbach and U. Wenger, PoS **LAT2007**, 102 (2007), [arXiv:0710.2498 \[hep-lat\]](#).
- [62] **ETM** Collaboration, C. Alexandrou *et al.*, Phys. Rev. **D78**, 014509 (2008), [arXiv:0803.3190 \[hep-lat\]](#).
- [63] **ETM** Collaboration, R. Baron *et al.*, JHEP **1008**, 097 (2010), [arXiv:0911.5061 \[hep-lat\]](#).
- [64] **ETM** Collaboration, V. Drach, K. Jansen, J. Carbonell, M. Papinutto and C. Alexandrou, PoS **LATTICE2010**, 101 (2010), [arXiv:1012.3861 \[hep-lat\]](#).
- [65] G. Herdoiza, PoS **LATTICE2010**, 010 (2010), [arXiv:1103.1523 \[hep-lat\]](#).
- [66] $\chi_{\mathbf{L}}^{\mathbf{F}}$ Collaboration, K. Jansen *et al.*, Phys. Lett. **B624**, 334 (2005), [hep-lat/0507032](#).
- [67] R. Frezzotti and G. Rossi, PoS **LAT2007**, 277 (2007), [arXiv:0710.2492 \[hep-lat\]](#).
- [68] **ETM** Collaboration, P. Dimopoulos, R. Frezzotti, C. Michael, G. Rossi and C. Urbach, Phys.Rev. **D81**, 034509 (2010), [arXiv:0908.0451 \[hep-lat\]](#).
- [69] **ETM** Collaboration, C. Michael and C. Urbach, PoS **LAT2007**, 122 (2007), [arXiv:0709.4564 \[hep-lat\]](#).
- [70] **ETM** Collaboration, P. Boucaud *et al.*, Comput.Phys.Commun. **179**, 695 (2008), [arXiv:0803.0224 \[hep-lat\]](#).
- [71] R. Frezzotti and G. C. Rossi, JHEP **10**, 070 (2004), [hep-lat/0407002](#).
- [72] R. Frezzotti and G. C. Rossi, Nucl. Phys. Proc. Suppl. **128**, 193 (2004), [hep-lat/0311008](#).
- [73] K. Osterwalder and E. Seiler, Ann. Phys. **110**, 440 (1978).
- [74] C. Morningstar and M. J. Peardon, Phys. Rev. **D69**, 054501 (2004), [arXiv:hep-lat/0311018](#).
- [75] B. Sheikholeslami and R. Wohlert, Nucl. Phys. **B259**, 572 (1985).
- [76] P. Weisz, Nucl. Phys. **B212**, 1 (1983).
- [77] P. Weisz and R. Wohlert, Nucl. Phys. **B236**, 397 (1984).
- [78] Y. Iwasaki, Nucl. Phys. **B258**, 141 (1985).
- [79] Y. Iwasaki, K. Kanaya, T. Kaneko and T. Yoshie, Phys.Rev. **D56**, 151 (1997), [arXiv:hep-lat/9610023 \[hep-lat\]](#).
- [80] Y. Iwasaki, [arXiv:1111.7054 \[hep-lat\]](#).

- [81] **ETM** Collaboration, R. Baron *et al.*, PoS **LATTICE2010**, 123 (2010), [arXiv:1101.0518 \[hep-lat\]](#).
- [82] **ETM** Collaboration, M. Constantinou *et al.*, Phys.Rev. **D83**, 014505 (2011), [arXiv:1009.5606 \[hep-lat\]](#).
- [83] **ETM** Collaboration, F. Farchioni *et al.*, PoS **LATTICE2010**, 128 (2010), [arXiv:1012.0200 \[hep-lat\]](#).
- [84] **ETM** Collaboration, B. Blossier *et al.*, Phys.Rev. **D82**, 114513 (2010), [arXiv:1010.3659 \[hep-lat\]](#).
- [85] K. Cichy, G. Herdoiza and K. Jansen, Nucl.Phys. **B847**, 179 (2011), [arXiv:1012.4412 \[hep-lat\]](#).
- [86] P. Dimopoulos, H. Simma and A. Vladikas, JHEP **0907**, 007 (2009), [arXiv:0902.1074 \[hep-lat\]](#).