

Effects of Acute and Chronic Gamma Irradiation on *in vitro* Growth of *Stevia rebaudiana* Bertoni.

Norazlina Noordin, Rusli Ibrahim, Nur Hidayah Mohd Sajahan, Salmah Moosa and Sobri Hussein

Agrotechnology and Biosciences Division, Malaysian Nuclear Agency,
Ministry of Science, Technology and Innovation Malaysia (MOSTI),
Bangi, 43000 Kajang, Selangor
azlina@nuclearmalaysia.gov.my

Abstract

Stevia rebaudiana Bertoni is a perennial herb that belongs to the family of Asteraceae. It is a natural sweetener plant known as sweet leaf, which is estimated to be 300 times sweeter than cane sugar. In this study, micropropagation and *in vitro* mutagenesis of this natural herb was successfully conducted. It was found that shoot tips on MS medium supplemented with 1 mg/l Kinetin showed the highest shoot induction and multiplication after 3 weeks of culture ($5.50 \pm 1.95a$). Radiosensitivity test was conducted to identify the LD50 for *in vitro* stevia shoots and to select effective doses to be used for the *in vitro* mutagenesis. Shoot tips were irradiated with acute and chronic gamma radiation at 0, 10.00, 20.00, 30.00, 40.00, 60.00, and 80.00 Gy. At 60 Gy and 80 Gy, the shoot tips demonstrated 0% survival, all were killed. In this study, LD 50 for the stevia (the dose that killed 50% of the irradiated explants) was at 29 Gy for acute irradiation and was at 45 Gy for chronic irradiation. The effective doses were selected at 10, 20, 30 and 40 Gy. These selected doses were applied for the *in vitro* mutagenesis of the stevia shoots.

Keywords: *Stevia rebaudiana*, micropropagation, *in vitro* mutagenesis, gamma irradiation

INTRODUCTION

Stevia is herbaceous perennial shrub originated from the highlands of Paraguay and sections of Argentina and Brazil. Stevia was discovered by Antonio Bertoni, a South American Natural Scientist, in 1887. This plant is a natural sweetener and famously known as “Sweet Weed”, “Sweet Leaf”, “Sweet Herbs” and “Honey Leaf”, which is estimated to be 300 times sweeter than cane sugar. *Stevia rebaudiana* Bert belongs to the family Asteraceae, one of 154 members of the genus Stevia, which produces sweet steviol glycosides (Robinson, 1930; Soejarto *et al.*, 1982). The leaves of stevia are the source of diterpene glycosides, stevioside and rebaudioside (Yoshida, 1986). Stevioside is regarded as a valuable natural sweetening agent because of its relatively good taste and chemical stability (Yamada *et al.*, 1997). Stevioside is of special interest to diabetic persons with hyperglycemia and the diet conscious (Arpita *et al.*, 2011). In Japan alone, an estimated 50 tons of stevioside is used annually with sales valued in order of \$220 million Canadian (Brandle and Rosa, 1992). Now, stevia has been introduced as a crop in a number of countries including Brazil, Korea, Japan, Mexico, United States, Indonesia, Tanzania and Canada (Shock, 1982; Saxena and Ming, 1988; Brandle and Rosa, 1992; Fors, 1995) for food and pharmaceuticals products. Currently *S. rebaudiana* production is centered in China with major market in Japan (Kinghorn and Soejarto, 1985). The product also can be added to tea and coffee, cooked or baked goods, processed foods, pickles, fruit juices, tobacco products, confectionary goods, jams and jellies, candies, yogurts, pastries, chewing gum and sherbets beverages (Arpita *et al.*, 2011).

Seeds of stevia show a very low germination percentage and vegetative propagation is limited by lower number of individuals (Sakaguchi *et al.*, 1982). Tissue culture is the only rapid process for the mass propagation of stevia and there have been few reports of *in vitro* growth of stevia (Miyagaya *et al.*, 1986), *in vitro* micropropagation from shoot tip and leaf (Uddin *et al.*, 2006).

Mutation techniques using radiation have been widely applied to improve crop yield, quality and disease and pest resistance. They have produced many germplasm with novel and desired traits (Maluszynski *et al.* 1995). Many mutants were developed released worldwide in many crop plants (Ahloowalia *et al.* 2004). Gamma rays were reported to be the most efficient ionizing radiation of creating mutants in plants as they can induce high mutation rate. They could also modify physiological characteristics to create new mutants with improved properties that can produce higher amounts of commercially essential metabolites, developing varieties that are agriculturally and economically significant and contain high productivity potential.

At present, there is no report on the use of radiation to induce mutation for the improvement of physiological traits and enhancement of secondary metabolites, steviol glycosides in *Stevia rebaudiana*. However, gamma irradiation had been applied on other herbal and medicinal plants for the same purpose. Sina *et al.* (2011) found that acute gamma rays at doses of 40 and 60Gy had a positive effect on the stimulation of flavonoid in two accessions of *Centella asiatica* (CA03 and CA23). Biochemical tests revealed that the irradiated accessions displayed higher total flavonoid content than the non-irradiated ones. Moreover, the irradiated plants were found to have the highest flavonoid contents in their eighth week of growth.

Therefore, the main objectives of this project are to study the effects of chronic gamma irradiation on growth and multiplication rate of *in vitro* shoots and to study the potential of gamma rays for the stimulation and enhancement of steviol glycosides in *Stevia rebaudiana*. Hopefully with the introduction of a new variety of *Stevia rebaudiana* with enhanced production of steviol glycosides can create an improvement and utilization of this natural sweetener to develop a sustainable Stevia industry in Malaysia.

MATERIALS AND METHODS

Collection of explants and surface sterilization:

Young, actively growing shoot tips and nodal segments were collected from stevia seedlings that were maintained in the glasshouse of Malaysian Nuclear Agency. The collected explants of 2-3 cm were put under running tap water to remove any traces of soil and dirt and later shake with systemic fungicide 5% (w/v). In laminar air flow which provided a strict aseptic condition, the cleaned explants were surface sterilized using 5% (v/v) commercial sodium hypochlorite with a few drops of Tween 20 for 15 min. This step was repeated twice and then rinsed 4 times with sterile distilled water. Excess water adhering to the explants was air-dried then the explants were ready to be inoculated into culture media.

Shoot induction and multiplication

Sterilized shoot tips and nodal segment (with a single axillary bud) were cultured onto semi-solid MS Medium (Murashige and Skoog, 1962) supplemented with 6- benzylamino purine (BAP) or 6-furfurylaminopurine (kinetin) at concentrations ranging from 0, 0.5, 1.0 and 1.5 mg/l. These plant growth regulators (PGRs) were added singly into the MS medium together with 3% (w/v) sucrose and 2.5 g/l gelrite. The pH of the medium was adjusted at 5.8 with 0.1M NaOH before autoclaving at 121°C for 15 min. The cultures were incubated in the incubation room at 24± 2°C with 16 h photoperiod. Observation on new shoots induction and shoot multiplication was done weekly. The length of plant and number of new shoots formed were monitored weekly for one month. The data was analyzed by SPSS statistical analysis version 19 using one way ANOVA with Duncan test.

Subcultures were done every 30 days interval. Nodal segments and shoot tips from the proliferated shoots were subcultured for further multiple shoot induction. The regenerated multiple shoots were cut and individual shoots were placed in semi-solid MS without PGR for elongation and root induction.

Table 1 Different treatment for shoots formation of *Stevia rebaudiana*

Treatment		Treatment	
1	MSO (shoot tip)	8	MS + 1.5mg/L Kin (nodal)
2	MSO (nodal)	9	MS + 0.5 mg/L BAP (shoot tip)
3	MS + 0.5 mg/L kin (shoot tip)	10	MS + 0.5 mg/L BAP (nodal)
4	MS + 0.5 mg/L Kin (nodal)	11	MS + 1.0 mg/L BAP (shoot tip)
5	MS + 1.0 mg/L Kin (shoot tip)	12	MS + 1.0 mg/L BAP (nodal)
6	MS + 1.0 mg/L Kin (nodal)	13	MS + 1.5 mg/L BAP (shoot tip)
7	MS + 1.5 mg/L Kin (shoot tip)	14	MS + 1.5 mg/L BAP (nodal)

In vitro mutagenesis with acute and chronic gamma irradiation

The *in vitro* grown and healthy looking stevia shoot tips were cultured onto semi-solid MS Medium supplemented with 1.0 mg/l kinetin. After 7 days of culture, these explants were irradiated with acute gamma radiation at 0, 10, 20, 30, 40, 60, and 80 Gy using Gamma Cell facility and chronic gamma irradiation at 0, 10, 20, 30, 40, 60, and 80 Gy using Gamma Greenhouse facility. The irradiated explants were incubated in the incubation room at 24± 2°C with 16 h photoperiod. The survival rate, length of plant, number of new shoots formed, and number of leaves formed were recorded weekly for a period of 30 days. The data obtained was analyzed by SPSS statistical analysis version 19 using one way ANOVA with Duncan test.

RESULTS AND DISCUSSION

Micropropagation of *S. rebaudiana*

Table 2 Effects of 14 different treatments on *stevia rebaudiana* growth after 4 weeks of culture

Treatment	Length of plantlet (cm±SD)	No of new shoot formed (mean±SD)
1	9.97±3.00a	1.24±0.59f
2	8.06±3.26b	1.70±0.54ef
3	5.89±1.93c	5.32±1.83ab
4	6.18±1.30c	4.92±1.38ab
5	5.95±1.15c	5.50±1.95a
6	5.64±1.32cd	4.70±1.43b
7	1.03±0.54gh	1.40±0.70f
8	0.61±0.44h	1.62±0.70ef
9	5.13±1.18de	3.86±1.39c
10	4.96±2.90e	5.24±2.44ab
11	4.50±0.94ef	2.70±1.75d
12	4.23±0.87f	2.18±1.65de
13	0.92±0.43gh	1.26±2.18f
14	1.53±0.83g	1.62±1.82ef

The above results showed significant difference for the length of plant and number of shoots for each treatment. In this study, explants cultured on MS media without hormone gave the highest length of plantlet but having lowest number of new shoots formed. It was observed that, the best treatment for stevia growth and shoot multiplication was in MS media supplemented with low concentration of kinetin (0.5-1.0 mg/l) for both shoot tips and nodal explants. After 3 weeks of culture in kinetin, it can be seen that multiple shoots formed into complete plantlets with vigor stem, larger, greener and normal looking leaves, no callus formation was observed too. Meanwhile, for BAP treatments, number of new shoot produced was lower and from the observation, stevia shoots produced in BAP demonstrated curly and abnormal looking phenotype. Callus growth was also observed at the base of the shoots. Comparatively, it was found that MS media supplemented with 1.0 mg/L kinetin is the most suitable and workable treatment to be used for shoot induction and multiplication ($5.50 \pm 1.95a$) and shoot tips showed better potential to be used as explants compared to the nodal.




Figure 1 Comparison of shoot multiplication in Kinetin and BAP

Effects of acute and chronic gamma irradiation

Using acute gamma irradiation

Table 3 Effects of acute gamma irradiation on *stevia rebaudiana* growth after 4 weeks of culture

Dose (Gy)	Survival rate (%±SD)	Length of plant (cm±SD)	No of new shoots formed (mean±SD)	No of leaves formed (mean±SD)
0	100a	5.74±2.43a	5.00±1.98a	26.96±7.92a
10.0	100a	4.96±1.49b	5.06±1.98a	24.13±9.84a
20.0	73.33±44.98b	1.35±1.12c	1.40±1.10b	8.37±5.16b
30.0	46.47±50.74c	1.20±1.49c	0.93±1.23b	4.77±5.78c
40.0	40.00±49.83c	0.74±1.35cd	0.90±1.63b	4.40±6.99c
60.0	0	0	0	0
80.0	0	0	0	0

The data are from 5 replications and were analyze by one way ANOVA statistical analysis. The numbers followed by different letters shows significant difference at $p < 0.05$ by Duncan test.

Based on table 3, the results showed significant difference for the percentage of the survival amongst shoots irradiated with lower and higher doses of gamma radiation. All non- irradiated shoots (control/0Gy) and shoots irradiated at 10.0 Gy showed 100% survival rate with the highest number of new shoots formed, 5.00 ± 1.98 and 5.06 ± 1.98 respectively. It was observed that, the survival rate of the shoot tips declined with the increasing dose of gamma radiation. At 60 Gy and 80 Gy, the shoot tips demonstrated 0% survival, all were killed.




Figure 2 Effects of acute gamma radiation on plantlets growth after 4 weeks of culture in MS + 1 mg/L Kinetin




Figure 3 LD₅₀ of *Stevia rebaudiana* after 4 weeks of culture irradiated with acute gamma

From the LD50 results, it was clearly demonstrated that with the increasing gamma dose, the survival rate of stevia shoots showed significant declining in the survival rate. Based on the figure above, LD 50 for the stevia (the dose that killed 50% of the irradiated explants) was at 29 Gy. From this study, the effective doses for acute irradiation were selected at 10, 20 and 30 Gy. These three selected doses were applied for the *in vitro* mutagenesis of the stevia shoots.

Using chronic gamma irradiation

This radiosensitivity test was conducted as to identify the LD50 for *in vitro* stevia shoots and to select effective doses to be used for the *in vitro* mutagenesis.. Results from table 4 showed significant difference for the percentage of the survival amongst shoots irradiated with lower and higher doses of chronic gamma radiation. All non-irradiated shoots (control/0Gy) and shoots irradiated at 10.0 and 20.0 Gy showed higher survival rate with the highest length of plant, 5.55±2.24, 6.26±2.74 and 6.33±1.84 respectively. Non irradiated shoots produced higher number of new shoot compared to irradiated shoots. It was observed that, the survival rate of the shoot tips declined with the increasing dose of gamma radiation. At 60, and 80 Gy the shoot tips demonstrated 0% survival, all were killed.

Table 4 Effect of different doses of chronic gamma irradiation on *Stevia rebaudiana* Bertoni growth after 4 week of culture

dose	Survival rate (%±SD)	Length of plant (cm±SD)	No of new shoot form (mean±SD)	No of leave(mean±SD)
0	100a	5.55±2.24a	4.43±2.09a	25.83±10.10a
10.0	92.50±26.67a	6.26±2.74a	3.33±1.828b	16.53±7.37b
20.0	100a	6.33±1.84a	3.00±1.88b	19.18±.59b
30.0	80.00±40.50b	4.37±2.89b	1.90±1.70c	12.10±7.53c
40.0	47.50±50.57c	2.00±2.31c	1.45±1.89d	5.45±6.73d
60.0	0	0	0	0
80.0	0	0	0	0

Data was taken from 5 replicates with twenty explants. Significance p<0.05 using Duncan's Multiple Range Test. SD= Standard Deviation.




Figure 4 LD₅₀ of *Stevia rebaudiana* after 4 weeks of culture irradiated with chronic gamma

From the LD50 results, it was clearly demonstrated that with the increasing gamma dose, the survival rate of stevia shoots showed significant declining. Based on the figure above, LD 50 for the stevia (the dose that killed 50% of the irradiated explants) was at 45 Gy. From this study, the effective doses were selected at 10, 20, 30 and 40 Gy. These four selected doses were applied for the *in vitro* mutagenesis of the stevia shoots. Shoot tip explants were irradiated with these doses and regenerated shoots were sub-cultured up to M1V4 generation to reduce the effect of chimeras.




Figure 1 Effect of acute gamma radiation on plantlets growth after 4 weeks of culture in MS + 1 mg/L Kinetin

CONCLUSION

The micropropagation protocol of *Stevia rebaudiana* Bertoni has been successfully optimized and established. This study suggested that 1.0mg/L Kinetin is the optimal PGR for stevia shoot multiplication and MS basal medium played a good role in promoting the elongation and root induction of stevia plantlets. The radiosensitivity test for *Stevia rebaudiana* Bertoni has been successfully optimized and established for both acute and chronic gamma irradiation. The LD50 for stevia shoots was successfully determined and the effective doses were selected at 10, 20, 30 and 40 Gy for chronic and 10, 20, 30 Gy for acute. These selected doses were applied for the *in vitro* mutagenesis of the stevia shoots and were mass propagated for further screening of desired traits.

ACKNOWLEDMENT

The authors would like to thank the management of Malaysian Nuclear Agency for their continuous assistance, advices and supports on this project. Special thanks is also extended to the Ministry of Science, Technology and Innovation (MOSTI) for providing fund for this project under the Science Fund grant 02-03-01-SF0163: *Effects of chronic irradiation on growth and multiplication rate with reference to enhanced production of steviol glycoside in Stevia rebaudiana Bertoni.*

REFERENCES

- Ahloowalia BS, Maluszynski M, Nichterlein K (2004) Global impact of mutation-derived varieties. *Euphytica* 135: 187-204.
- Arpita D., Gantait S. and Mandal N. (2011) Micropropagation of an elite medicinal plant: *Stevia rebaudiana* Bert. *Int. J Agric. Resech.* 6(1):40-48
- Brandle JE, Rosa N (1992). Heritability for yield, leaf- stem ratio and stevioside content estimated from a landrace cultivar of *Stevia rebaudiana*. *Canadian Journal of Plant Science.* 72: 1263-1266.
- Brandle, J. E., A. N. Starratt and M. Gijzen (1998). *Stevia rebaudiana*: Its agricultural, biological and chemical properties. *Canad. J. Plant Sci.* 78:527-536
- Fors A (1995). A new character in the sweetener scenario. *Sugar J*,58:30.
- Kinghorn AD, Soejarto DD (1985). Current status of stevioside as asweetening agent for human use. In Wagner, Hikino H, Fransworth NR. (eds).Economic and medical plant research. Academic Press,London.
- Maluszynski M, Ahloowalia BS, Sigurbjornsson B (1995) Application of *in vivo* and *in vitro* mutation techniques for crop improvement. *Euphytica* 85: 303-315
- Murashige, T. and F. Skoog (1962). A revised medium for rapid growth and bioassay with tobacco tissue culture. *Physiol. Plant.* 15:473-497
- Robinson BL (1930). Contributions from the Grey Herbarium of Harvard University. *The Grey Herbarium of Harvard University, Cambridge.*
- Saxena NC, Ming IS (1988). Preliminary harvesting characteristics of Stevia. *Phys. Prop. Agric. Mat. Prod.* 3: 299-303.
- Sina Siavash Moghaddam, Hawa Jaafar, Rusli Ibrahim, Asmah Rahmat, Maheran Abdul Aziz, Elizabeth Philip (2011). Effects of Acute Gamma Irradiation on Physiological Traits and Flavonoid Accumulation of *Centella asiatica*. *Molecules* 2011, 16, 4994-5007
- Soejarto DD, Kinghorn AD, Fransworth NR (1982). Potential sweetening agents of plant origin. *J. Nat. Prod.* 45: 590-599.
- Shock CC (1982). Experimental cultivation of Rebaudia's stevia in California. University of California Agronomy Progress Report No.122
- Yoshida, S. (1986). Studies on the production of sweet substances in *Stevia rebaudiana*: I. Simple determination of sweet glucoside in *Stevia* plant by thin layer chromate-scanner and their accumulation patterns with plant growth. *Jap. J. Crop. Sci.* 55(2):189-195